



Gamma-rays from CR sources

Michael Kachelrieß

NTNU, Trondheim



TeV gamma-rays from UHECR sources



TeV gamma-rays from UHECR sources

- during propagation: “cosmogenic” photons
- in sources:



TeV gamma-rays from UHECR sources

- during propagation: “cosmogenic” photons
- in sources:
 - ▶ galactic CR sources
 - ▶ GRB:
 - ★ large redshift
 - ★ time-delay UHECR \leftrightarrow photons makes direct correlation impossible
 - ★ for small EGMF: auto-correlation $n_{\text{GRB}} \sim \tau \dot{n}_{\text{GRB}} < n_{\text{AGN}}$



TeV gamma-rays from UHECR sources

- during propagation: “cosmogenic” photons
- in sources:
 - ▶ galactic CR sources
 - ▶ GRB:
 - ★ large redshift
 - ★ time-delay UHECR \leftrightarrow photons makes direct correlation impossible
 - ★ for small EGMF: auto-correlation $n_{\text{GRB}} \sim \tau \dot{n}_{\text{GRB}} < n_{\text{AGN}}$
 - ▶ AGN:
 - ★ jets: small densities, B
 - ★ core: high B , large UV and IR densities, $\tau_{p\gamma} \sim 1$



HESS observations of M87:



HESS observations of M87:



HESS & Veritas observations of M87:



Outline of the talk

- 1 Introduction
- 2 Gamma-rays produced in UHECR sources
 - ▶ How do we get multi TeV gamma-rays out of AGN cores: electromagnetic cascades in UHECR sources
- 3 Cosmogenic fluxes:
 - ▶ Cosmogenic neutrino limits from Fermi-LAT
 - ▶ Cosmogenic photons: diffuse flux
 - ▶ Secondary photons from CR point sources
- 4 Lower limit on EGMF using gamma-rays
- 5 Summary

Outline of the talk

- 1 Introduction
- 2 Gamma-rays produced in UHECR sources
 - ▶ How do we get multi TeV gamma-rays out of AGN cores: electromagnetic cascades in UHECR sources
- 3 Cosmogenic fluxes:
 - ▶ Cosmogenic neutrino limits from Fermi-LAT
 - ▶ Cosmogenic photons: diffuse flux
 - ▶ Secondary photons from CR point sources
- 4 Lower limit on EGMF using gamma-rays
- 5 Summary

Outline of the talk

- 1 Introduction
- 2 Gamma-rays produced in UHECR sources
 - ▶ How do we get multi TeV gamma-rays out of AGN cores: electromagnetic cascades in UHECR sources
- 3 Cosmogenic fluxes:
 - ▶ Cosmogenic neutrino limits from Fermi-LAT
 - ▶ Cosmogenic photons: diffuse flux
 - ▶ Secondary photons from CR point sources
- 4 Lower limit on EGMF using gamma-rays
- 5 Summary

Outline of the talk

- 1 Introduction
- 2 Gamma-rays produced in UHECR sources
 - ▶ How do we get multi TeV gamma-rays out of AGN cores: electromagnetic cascades in UHECR sources
- 3 Cosmogenic fluxes:
 - ▶ Cosmogenic neutrino limits from Fermi-LAT
 - ▶ Cosmogenic photons: diffuse flux
 - ▶ Secondary photons from CR point sources
- 4 Lower limit on EGMF using gamma-rays
- 5 Summary


Multi-messenger astronomy with Cen A?

- + 2 events correlated with Cen A within 3.1°
- + more events close-by
- + general correlation with AGN
- confusion with LSS?
- no confirmation by HiRes
- tension to PAO chemical composition
- E_{\max} for most AGN (incl. Cen A) high enough?


[Gorbunov et al. '07, Fargione '08, Rachen '08]

correlations with AGN:

- independent/additional evidence?
 - Cen A closest AGN
- ⇒ good test case for multi-messenger astronomy: accompanying γ -ray and neutrino fluxes?


Results for acceleration close to the core: $\alpha = 1.2$ 

Results for acceleration close to the core: $\alpha = 2$




- promising test case for HESS / FERMI
- γ -ray spectrum rather insensitive to α

HESS observations of Cen A




- no variability

HESS observations of Cen A




- no variability
- consistent with point source

HESS observations of Cen A




- no variability
- consistent with point source
- HE emission from central region ($1' \simeq 1.1 \text{ kpc}$)

Comparison to recent HESS and FERMI observations




Comparison to recent HESS and FERMI observations



- shape and normalization okay
- TeV γ -ray and neutrino source


Regenerating TeV photons: a) (isotropic) source

- injection spectrum $F_\gamma(E) \propto 1/E^2$




Regenerating TeV photons: a) (isotropic) source

- : thin above 10^{16} eV, ultra-rel. regime




Regenerating TeV photons: b) on EBL

- photons above 10^{14} eV cascade on EBL




Regenerating TeV photons: b) on EBL

- photons above 10^{14} eV cascade on EBL : fill up GeV–TeV range



Regenerating TeV photons: b) on EBL


- photons above 10^{14} eV cascade on EBL : fill up GeV–TeV range




main criteria: L_{UHE} , not shape

- evidence for acceleration to UHECRs?
- however: **anisotropic UV field complicates picture already in source**


Regenerating TeV photons: collinear regime




Regenerating TeV photons: decaying muons



Regenerating TeV photons: adding IR




adding IR from a compact source:



Cascade limit for cosmogenic neutrinos

- **Photon** and **neutrino** production relatively **tight connected**:
 - ▶ protons:



Cascade limit for cosmogenic neutrinos

- **Photon** and **neutrino** production relatively tight **connected**:

- ▶ protons:

$$p + \gamma_{3K} \rightarrow \begin{cases} p + \pi^0 \rightarrow p + 2\gamma \\ n + \pi^+ \rightarrow p + 2e^\pm + 4\nu \end{cases}$$

- ▶ nuclei: $A + \gamma_{3K} \rightarrow (A - 1) + n \rightarrow (A - 1) + p + e^- + \nu_e$
- ▶ **connection to UHECRs looser**

Cascade limit for cosmogenic neutrinos

- Photon and neutrino production relatively tight connected:

- ▶ protons:


$$p + \gamma_{3K} \rightarrow \begin{cases} p + \pi^0 \rightarrow p + 2\gamma \\ n + \pi^+ \rightarrow p + 2e^\pm + 4\nu \end{cases}$$

- ▶ nuclei: $A + \gamma_{3K} \rightarrow (A - 1) + n \rightarrow (A - 1) + p + e^- + \nu_e$

- **cascade limit:**

[Berezinsky, Smirnov '75]

all energy in γ and e^\pm cascades below ~ 100 GeV



Cascade limit for cosmogenic neutrinos

- Photon and neutrino production relatively tight connected:

- ▶ protons:

$$p + \gamma_{3K} \rightarrow \begin{cases} p + \pi^0 \rightarrow p + 2\gamma \\ n + \pi^+ \rightarrow p + 2e^\pm + 4\nu \end{cases}$$

- ▶ nuclei: $A + \gamma_{3K} \rightarrow (A - 1) + n \rightarrow (A - 1) + p + e^- + \nu_e$

- **cascade limit:**

[Berezinsky, Smirnov '75]

all energy in γ and e^\pm cascades below ~ 100 GeV

$$J_\gamma(E) = \begin{cases} K(E/\varepsilon_X)^{-3/2} & \text{at } E \leq \varepsilon_X \\ K(E/\varepsilon_X)^{-2} & \text{at } \varepsilon_X \leq E \leq \varepsilon_a \\ 0 & \text{at } E > \varepsilon_a \end{cases}$$

Cascade limit for cosmogenic neutrinos

- Photon and neutrino production relatively tight connected:

- ▶ protons:

$$p + \gamma_{3K} \rightarrow \begin{cases} p + \pi^0 \rightarrow p + 2\gamma \\ n + \pi^+ \rightarrow p + 2e^\pm + 4\nu \end{cases}$$

- ▶ nuclei: $A + \gamma_{3K} \rightarrow (A - 1) + n \rightarrow (A - 1) + p + e^- + \nu_e$

- cascade limit:


[Berezinsky, Smirnov '75]

all energy in γ and e^\pm cascades below ~ 100 GeV


- Fermi-LAT measurement of EGRB:

$$\begin{aligned} \omega_{\text{cas}} &> \frac{4\pi}{c} \int_{E_0}^{\infty} dE EI_\nu(E) \geq \frac{4\pi}{c} E_0 I_\nu(> E_0) \\ &\lesssim 6 \cdot 10^{-7} \text{ eV/cm}^3 \end{aligned}$$

Comparison of Monte Carlo and analytical estimate




Comparison of Monte Carlo and analytical estimate




Fermi-LAT vs. UHECR data:

[Berezinsky et al. '10]



Fermi-LAT vs. UHECR data:


[Berezinsky et al. '10]



integrating $EJ(E)$ gives bound $\omega_{\text{cas}} \lesssim 6 \cdot 10^{-7} \text{ eV/cm}^3$


Cascade limit for cosmogenic neutrinos

[Berezinsky et al. '10]



Cascade limit for cosmogenic neutrinos


[Ahlers et al. '10]



Cascade limit for cosmogenic neutrinos

Main difference: expected IceCube sensitivity


- end of exp. sensitivity for $1/E^2$ flux: 10^{17} vs. 10^{19} eV
- overall sensitivity of IceCube



Cascade limit for cosmogenic neutrinos


Main difference: expected IceCube sensitivity

- end of exp. sensitivity for $1/E^2$ flux: 10^{17} vs. 10^{19} eV
- overall sensitivity of IceCube




All plots for proton primaries

- for nuclei reduced neutrino fluxes...




Cosmogenic photons

[Hooper, Taylor, Sarkar '10]



Cosmogenic photons


[Hooper, Taylor, Sarkar '10]



- $A > 1$: photons suppressed by factor 10 compared to protons
- even proton: requires sensitivity improved by ~ 100
- large E_{\max} and small α helps...


Cosmogenic photons

[Hooper, Taylor, Sarkar '10]



Cosmogenic photons: dependence on α


[Gelmini, Kalashev, Semikoz '05]



Secondary photons from CR point sources

- for $d < \text{few} \times 10 \text{ Mpc}$: UHE photons survive

[Taylor et al. '09]




Secondary photons from CR point sources

- for $d < \text{few} \times 10 \text{ Mpc}$: UHE photons survive [Taylor et al. '09]
- non-observation is no constraint, but observation identifies close source


Secondary photons from CR point sources

- for $d < \text{few} \times 10 \text{ Mpc}$: UHE photons survive [Taylor et al. '09]
- **larger distance:** UHE photons cascade down in TeV range [Essey et al. '10]
 - ▶ universal shape, depending only on EBL and z



Secondary photons from CR point sources

- for $d < \text{few} \times 10 \text{ Mpc}$: UHE photons survive [Taylor et al. '09]
- larger distance: UHE photons cascade down in TeV range [Essey et al. '10]
 - ▶ universal shape, depending only on EBL and z




- ▶ **large luminosities** $L_{\text{eff}} \sim 10^{47} - 10^{49} \text{ erg/s}$ in UHECR
- ▶ **relatively small EGMFs** in voids, $B \lesssim 10^{-14} \text{ G}$

Gamma-rays and extragalactic magnetic fields (EGMF)

- Origin of **seed** for EGMF is **mysterious**
- Seed required as input for EGMF simulations

Gamma-rays and extragalactic magnetic fields (EGMF)

- Origin of seed for EGMF is mysterious
- Seed required as input for EGMF simulations



Gamma-rays and extragalactic magnetic fields (EGMF)

- Origin of seed for EGMF is mysterious
- Seed required as input for EGMF simulations
- Observations only in clusters,
 - ▶ synchrotron halo: $\Rightarrow B \sim (0.1 - 1) \mu\text{G}$
 - ▶ Faraday rotation: $\Rightarrow B \sim (1 - 10) \mu\text{G}$

Gamma-rays and extragalactic magnetic fields (EGMF)


- Origin of seed for EGMF is mysterious
- Seed required as input for EGMF simulations
- Observations only in clusters,
 - ▶ synchrotron halo: $\Rightarrow B \sim (0.1 - 1) \mu\text{G}$
 - ▶ Faraday rotation: $\Rightarrow B \sim (1 - 10) \mu\text{G}$
- Aharonian, Coppi, Völk '94: **Pair halos** around AGNs

Gamma-rays and extragalactic magnetic fields (EGMF)

- Origin of seed for EGMF is mysterious
- Seed required as input for EGMF simulations
- Observations only in clusters,
 - ▶ synchrotron halo: $\Rightarrow B \sim (0.1 - 1) \mu\text{G}$
 - ▶ Faraday rotation: $\Rightarrow B \sim (1 - 10) \mu\text{G}$
- Aharonian, Coppi, Völk '94: Pair halos around AGNs
- Plaga '95: **EGMFs deflect and delay cascade electrons**
 \Rightarrow search for delayed “echoes” of multi-TeV AGN flares/GRBs

Observer misaligned with jet:


[Neronov et al. '10]



- probability for misalignment $p \propto \vartheta_{\text{obs}} \Rightarrow$ most blazars viewed with $\vartheta_{\text{obs}} \sim \vartheta_{\text{jet}}$

Observer misaligned with jet:

[Neronov et al. '10]




- probability for misalignment $p \propto \vartheta_{\text{obs}} \Rightarrow$ most blazars viewed with $\vartheta_{\text{obs}} \sim \vartheta_{\text{jet}}$

\Rightarrow halos are not symmetric

Observer misaligned with jet:


[Neronov et al. '10]




- probability for misalignment $p \propto \vartheta_{\text{obs}} \Rightarrow$ most blazars viewed with $\vartheta_{\text{obs}} \sim \vartheta_{\text{jet}}$
- \Rightarrow halos are not symmetric
- \Rightarrow **time-delay** is function of ϑ ,

$$T_{\text{delay}}(\vartheta) \sim 3 \times 10^6 \text{yr} \left[\frac{(\vartheta_{\text{obs}} + \Theta_{\text{jet}})}{5^\circ} \right] \left[\frac{\vartheta}{5^\circ} \right]$$


Asymmetric halos around TeV blazars (“GeV jets”):



“GeV jets”: B dependence




“GeV jets”: time dependence of flares




Lower limit on EGMF:

- choose blazar: **large z** , stationary, **low GeV**, high multi-TeV emission



Lower limit on EGMF:


- choose blazar: large z , stationary, low GeV, high multi-TeV emission



- TeV photons cascade down:
 - ▶ small EGMF: fill up GeV range
 - ▶ “large” EGMF: deflected outside, isotropized


Lower limit on EGMF:

[F. Tavecchio et al. '10, A. Neronov, I. Vovk '10]



Lower limit on EGMF:


[F. Tavecchio et al. '10, A. Neronov, I. Vovk '10]



- $B \gtrsim 10^{-15} \text{ G}$
- some dependence on ϑ_{jet}

Lower limit on EGMF:


[F. Tavecchio et al. '10, A. Neronov, I. Vovk '10]



- $B \gtrsim 10^{-15} \text{ G}$
- some dependence on ϑ_{jet}
- no simulation of elmag. cascade with B
- what happens for **structured B** ?

Lower limit on EGMF:


[MK, Ostapchenko, Tomàs]



Lower limit on filling factor:


[MK, Ostapchenko, Tomàs '10]

- model filaments by a top-hat:




Lower limit on filling factor:

[MK, Ostapchenko, Tomàs '10]



Lower limit on filling factor:

[MK, Ostapchenko, Tomàs '10]




linear filling factor $\gtrsim 50\%$

- mainly 3-step cascade: $\gamma \rightarrow e^\pm \rightarrow \gamma$
- photon mean free path $D_\gamma(E) \sim 1000\text{--}50 \text{ Mpc}$
- electron mean free path $D_e(E) \sim \text{few kpc}$

$\log_{10}(E/eV)$

Lower limit on filling factor:

[MK, Ostapchenko, Tomàs '10]



linear filling factor $\gtrsim 50\%$

- mainly 3-step cascade: $\gamma \rightarrow e^\pm \rightarrow \gamma$
 - photon mean free path $D_\gamma(E) \sim 1000\text{--}50 \text{ Mpc}$
 - electron mean free path $D_e(E) \sim \text{few kpc}$
- \Rightarrow electrons are created “everywhere” and feel B only close to interaction point

$\log_{10}(E/\text{eV})$

EGMF in voids already observed?


[*Ando, Kusenko '10*]

- **stack** 170 brightest **AGN**

EGMF in voids already observed?

[Ando, Kusenko '10]


- stack 170 brightest AGN
- search for excess over PSF



EGMF in voids already observed?

[Ando, Kusenko '10]


- stack 170 brightest AGN
- search for **excess** over PSF



EGMF in voids already observed?

[Ando, Kusenko '10]

- stack 170 brightest AGN
- search for excess over PSF



⇒ explained by


$$B \sim 10^{-15} \text{ G } (\lambda_B / 1 \text{ kpc})^{-1/2}$$

and $\lambda_B < 10 - 100 \text{ kpc}$

EGMF in voids already observed?

[Ando, Kusenko '10]

- stack 170 brightest AGN
- search for excess over PSF




- lower limit $B \gtrsim 5 \times 10^{-15}$ G requires $\lambda_B \lesssim 0.1$ kpc

EGMF already observed? Probably not...

[Neronov et al. '10]


- point source Crab shows the same “halo” as stacked AGN:



EGMF already observed? Probably not...

[Neronov et al. '10]

- point source Crab shows the same “halo” as stacked AGN:



- tail of PSF wrong (?), difference between “front” and “back” photons

Summary

- multi-TeV photons from AGN cores require
 - ▶ photons in KN regime
 - ▶ HE muons

⇒ hadronic models

Summary

- multi-TeV photons from AGN cores require
 - ▶ photons in KN regime
 - ▶ HE muons

⇒ hadronic models

- secondary photons:
 - ▶ chemical composition
 - ▶ TeV sources ??

Summary

- multi-TeV photons from AGN cores require
 - ▶ photons in KN regime
 - ▶ HE muons

⇒ hadronic models

- secondary photons:
 - ▶ chemical composition
 - ▶ TeV sources ??
- **cascade limit** from Fermi data reduced by factor ~ 7

Summary

- multi-TeV photons from AGN cores require
 - ▶ photons in KN regime
 - ▶ HE muons

⇒ hadronic models

- secondary photons:
 - ▶ chemical composition
 - ▶ TeV sources ??
- cascade limit from Fermi data reduced by factor ~ 7
- lower limit on EGMF in voids $B \gtrsim 10^{-15} \text{ G}$