





DE LA RECHERCHE À L'INDUSTRIE

cea



Fundamental Interactions Seminar | Hervé MOUTARDE

May 30th, 2016

Quantum Chromodynamics as a paradigm.

The *theory* (and not an *effective theory*) of the strong interaction.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Facts

- **Restricted number** of parameters.
- Mathematically **consistent**.
- **Large** scope.
- Validated up to **large energy** $\lesssim 13$ TeV.
- Accurate **algorithmic** answer.

Open questions

- **Confinement**.
- **Chiral symmetry** breaking.
- Existence of a **mass gap**.



Quantum Chromodynamics as a paradigm.

The *theory* (and not an *effective theory*) of the strong interaction.

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning


Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Facts

- **Restricted number** of parameters.
- Mathematically **consistent**.
- **Large** scope.
- Validated up to **large energy** $\lesssim 13$ TeV.
- Accurate **algorithmic** answer.



Open questions

- **Confinement**.
- **Chiral symmetry** breaking.
- Existence of a **mass gap**.

Quantum Chromodynamics as a paradigm.

The theory (and not an *effective theory*) of the strong interaction.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning


Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Facts

- Restricted number of parameters.
- Mathematically **consistent**.
- Large scope.
- Validated up to large energy $\lesssim 13$ TeV.
- Accurate **algorithmic** answer.



Open questions

- Confinement.
- Chiral symmetry breaking.
- Existence of a **mass gap**.

Quantum Chromodynamics as a paradigm.

The theory (and not an *effective theory*) of the strong interaction.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning


Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Facts

- Restricted number of parameters.
- Mathematically **consistent**.
- **Large scope**.
- Validated up to **large energy** $\lesssim 13$ TeV.
- Accurate **algorithmic** answer.



Open questions

- **Confinement**.
- **Chiral symmetry** breaking.
- Existence of a **mass gap**.

Quantum Chromodynamics as a paradigm.

The *theory* (and not an *effective theory*) of the strong interaction.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Facts

- **Restricted number** of parameters.
- Mathematically **consistent**.
- **Large scope**.
- **Validated up to large energy** $\lesssim 13$ TeV.
- Accurate **algorithmic** answer.

Open questions

- **Confinement**.
- **Chiral symmetry** breaking.
- Existence of a **mass gap**.



Quantum Chromodynamics as a paradigm.

The **theory** (and not an *effective theory*) of the strong interaction.

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Facts

- **Restricted number** of parameters.
- Mathematically **consistent**.
- **Large** scope.
- Validated up to **large energy** $\lesssim 13$ TeV.
- **Accurate algorithmic** answer.

Open questions

- **Confinement**.
- **Chiral symmetry** breaking.
- Existence of a **mass gap**.



Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Facts

- **Restricted number** of parameters.
- Mathematically **consistent**.
- **Large** scope.
- Validated up to **large energy** $\lesssim 13$ TeV.
- Accurate **algorithmic** answer.

Open questions

- **Confinement**.
- **Chiral symmetry** breaking.
- Existence of a **mass gap**.



Quantum Chromodynamics as a paradigm.

The *theory* (and not an *effective theory*) of the strong interaction.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning


Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Facts

- **Restricted number** of parameters.
- Mathematically **consistent**.
- **Large** scope.
- Validated up to **large energy** $\lesssim 13$ TeV.
- Accurate **algorithmic** answer.



Open questions

- **Confinement**.
- **Chiral symmetry** breaking.
- Existence of a **mass gap**.

Quantum Chromodynamics as a paradigm.

The *theory* (and not an *effective theory*) of the strong interaction.

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Facts

- **Restricted number** of parameters.
- Mathematically **consistent**.
- **Large** scope.
- Validated up to **large energy** $\lesssim 13$ TeV.
- Accurate **algorithmic** answer.

Open questions

- **Confinement**.
- **Chiral symmetry** breaking.
- Existence of a **mass gap**.



Quantum Chromodynamics as a paradigm.

The *theory* (and not an *effective theory*) of the strong interaction.

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

No observed free color charges (PDG 2009)

FREE QUARK SEARCHES

The basis for much of the theory of particle scattering and hadron spectroscopy is the construction of the hadrons from a set of fractionally charged constituents (quarks). **A central but unproven hypothesis of this theory**, Quantum Chromodynamics, is that quarks cannot be observed as free particles but are confined to mesons and baryons.

Experiments show that it is at best difficult to “unglue” quarks. Accelerator searches at increasing energies have produced **no evidence for free quarks**, while only a few cosmic-ray and matter searches have produced uncorroborated events.



Open questions

- Confinement.
- Chiral symmetry breaking.
- Existence of a mass gap.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning


Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

From quarks to hadrons

- What are the relevant degrees of freedom?
- What are the effective forces between them?



Open questions

- **Confinement.**
- **Chiral symmetry breaking.**
- Existence of a **mass gap**.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Clay Millenium Prize (Jaffe and Witten)

QUANTUM YANG-MILLS THEORY

5

Finally, QFT is the jumping-off point for a quest that may prove central in 21st century physics—the effort to unify gravity and quantum mechanics, perhaps in string theory. For mathematicians to participate in this quest, or even to understand the possible results, QFT must be developed further as a branch of mathematics. It is important not only to understand the solution of specific problems arising from physics, but also to set such results within a new mathematical framework. One hopes that this framework will provide a unified development of several fields of mathematics and physics, and that it will also provide an arena for the development of new mathematics and physics.

For these reasons the Scientific Advisory Board of CMI has chosen a Millennium problem about quantum gauge theories. Solution of the problem requires both understanding one of the deep unsolved physics mysteries, the existence of a mass gap, and also producing a mathematically complete example of quantum gauge field theory in four-dimensional space-time.



Open questions

- Confinement.
- Chiral symmetry breaking.
- Existence of a mass gap.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Clay Millenium Prize (Jaffe and Witten)

QUANTUM YANG-MILLS THEORY

5


Finally, QFT is the jumping-off point for a quest that may prove central in 21st century physics—the effort to unify gravity and quantum mechanics, perhaps in string theory. For mathematicians to participate in this quest, or even to understand the possible results, QFT must be developed further as a branch of mathematics. It is important not only to understand the solution of specific problems



"I think you should be more explicit here in step two."

within a new mathematical framework. A unified development of several theories will also provide an arena for the

of CMI has chosen a Millennium Prize problem. The solution of the problem requires both mathematics, the existence of a mass gap is an example of quantum gauge




Open questions

- Confinement.
- Chiral symmetry breaking.
- Existence of a mass gap.


Nucleon Reverse Engineering

- Lattice QCD clearly shows that the mass of hadrons is generated by the **interaction**, not by the quark masses.



Durr et al., Science **322**, 1224 (2008)

- Can we **map** the *location of mass* inside a hadron?



QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap

Radon transform

Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

How can we recover the well-known characteristics of the nucleon from the properties of its **colored building blocks**?

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition


Overlap

Radon transform

Covariant extension

Conclusion


Appendix



Nucleon Reverse Engineering

How can we recover the well-known characteristics of the nucleon from the properties of its **colored building blocks**?

Mass?



QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap

Radon transform

Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

How can we recover the well-known characteristics of the nucleon from the properties of its **colored building blocks**?

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap


Radon transform

Covariant extension

Conclusion

Appendix

Mass?
Spin?



Nucleon Reverse Engineering

How can we recover the well-known characteristics of the nucleon from the properties of its **colored building blocks**?

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap


Radon transform

Covariant extension

Conclusion

Appendix

Mass?
Spin?
Charge?



Nucleon Reverse Engineering

How can we recover the well-known characteristics of the nucleon from the properties of its **colored building blocks**?

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap

Radon transform


Covariant extension

Conclusion

Appendix

Mass?
Spin?
Charge?

...



Nucleon Reverse Engineering

How can we recover the well-known characteristics of the nucleon from the properties of its **colored building blocks**?

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap

Radon transform

Covariant extension


Conclusion

Appendix

Mass?
Spin?
Charge?

...

What are the relevant **effective degrees of freedom** and **effective interaction** at large distance?



Nucleon Reverse Engineering

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap

Radon transform


Covariant extension

Conclusion

Appendix


Structuring questions for the hadron physics community

- QCD mechanisms behind the origin of mass in the visible universe?
- Cartography of interactions giving its mass to the nucleon?
- Pressure and density profiles of the nucleon as a continuous medium?
- Localization of quarks and gluons inside the nucleon?




Nucleon Reverse Engineering


Perturbative QCD



Asymptotic freedom



Nonperturbative QCD



QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap

Radon transform

Covariant extension

Conclusion

Appendix

Perturbative AND nonperturbative QCD at work

- Define **universal** objects describing 3D nucleon structure:
Generalized Parton Distributions (GPD).
- Relate GPDs to measurements using **factorization**:
Virtual Compton Scattering (DVCS, TCS),
Deeply Virtual Meson production (DVMP).
- Get **experimental knowledge** of nucleon structure.

- Correlation of the **longitudinal momentum** and the **transverse position** of a parton in a hadron.
- DVCS recognized as the cleanest channel to access GPDs.

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension


Conclusion

Appendix

Deeply Virtual Compton Scattering (DVCS)



Transverse center of momentum R_\perp
 $R_\perp = \sum_i x_i r_{\perp i}$



Nucleon Reverse Engineering

- Correlation of the **longitudinal momentum** and the **transverse position** of a parton in a hadron.
- DVCS recognized as the cleanest channel to access GPDs.

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap



Radon transform

Covariant extension

Conclusion

Appendix

Deeply Virtual Compton Scattering (DVCS)



- Correlation of the **longitudinal momentum** and the **transverse position** of a parton in a hadron.
- DVCS recognized as the cleanest channel to access GPDs.

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team



Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Deeply Virtual Compton Scattering (DVCS)



Nucleon Reverse Engineering

- Correlation of the **longitudinal momentum** and the **transverse position** of a parton in a hadron.
- DVCS recognized as the cleanest channel to access GPDs.

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team



Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Deeply Virtual Compton Scattering (DVCS)



- 24 GPDs** $F^i(x, \xi, t, \mu_F)$ for each parton type $i = g, u, d, \dots$ for leading and sub-leading twists.


- **Probabilistic interpretation** of Fourier transform of $\text{GPD}(x, \xi = 0, t)$ in **transverse plane**.

$$\rho(x, b_{\perp}, \lambda, \lambda_N) = \frac{1}{2} \left[H(x, 0, b_{\perp}^2) + \frac{b_{\perp}^j \epsilon_{ji} S_{\perp}^i}{M} \frac{\partial E}{\partial b_{\perp}^2}(x, 0, b_{\perp}^2) + \lambda \lambda_N \tilde{H}(x, 0, b_{\perp}^2) \right]$$

- Notations : quark helicity λ , nucleon longitudinal polarization λ_N and nucleon transverse spin S_{\perp} .

Burkardt, Phys. Rev. **D62**, 071503 (2000)

Can we obtain this picture from exclusive measurements?



Weiss, AIP Conf. Proc. **1149**, 150 (2009)

Nucleon
Reverse
Engineering

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap

Radon transform

Covariant extension

Conclusion

Appendix

**Nucleon
Reverse
Engineering**

- Most general structure of matrix element of energy momentum tensor between nucleon states:

$$\left\langle N, P + \frac{\Delta}{2} \left| T^{\mu\nu} \right| N, P - \frac{\Delta}{2} \right\rangle = \bar{u} \left(P + \frac{\Delta}{2} \right) \left[A(t) \gamma^{(\mu} P^{\nu)} + B(t) P^{(\mu} i \sigma^{\nu)\lambda} \frac{\Delta_\lambda}{2M} + \frac{C(t)}{M} (\Delta^\mu \Delta^\nu - \Delta^2 \eta^{\mu\nu}) \right] u \left(P - \frac{\Delta}{2} \right)$$

with $t = \Delta^2$.

- Key observation: **link between GPDs and gravitational form factors**

$$\int dx x H^q(x, \xi, t) = A^q(t) + 4\xi^2 C^q(t)$$

$$\int dx x E^q(x, \xi, t) = B^q(t) - 4\xi^2 C^q(t)$$

Ji, Phys. Rev. Lett. **78**, 610 (1997)

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap

Radon transform

Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

■ Spin sum rule:

$$\int dx x (H^q(x, \xi, 0) + E^q(x, \xi, 0)) = A^q(0) + B^q(0) = 2J^q$$

Ji, Phys. Rev. Lett. **78**, 610 (1997)

■ Shear and pressure of a hadron considered as a continuous medium:

$$\langle N | T^{ij}(\vec{r}) | N \rangle = s(r) \left(\frac{r^i r^j}{r^2} - \frac{1}{3} \delta^{ij} \right) + p(r) \delta^{ij}$$

Polyakov and Shuvaev, hep-ph/0207153

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap

Radon transform

Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

QCD

Mass without mass

Nucleon structure

Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition

Overlap

Radon transform

Covariant extension

Conclusion


Appendix

1 Status of 3D imaging: phenomenological relevance of the field.

2 Building the tools: preparing for the high precision era.

3 Learning from GPDs: steps towards new GPD models.

How can we make this picture? What do we learn from it?



Phenomenological status of nucleon 3D imaging

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access

DVCS Kinematics
Universality tests
Towards 3D images

Building



Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Exclusive processes of current interest (1/2). Factorization and universality.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access

DVCS Kinematics
Universality tests
Towards 3D images

Building



Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Exclusive processes of current interest (1/2). Factorization and universality.

**Nucleon
Reverse
Engineering**

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access

DVCS Kinematics
Universality tests
Towards 3D images

Building




Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Exclusive processes of current interest (1/2). Factorization and universality.

**Nucleon
Reverse
Engineering**

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access

DVCS Kinematics
Universality tests
Towards 3D images

Building





Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Exclusive processes of current interest (1/2). Factorization and universality.

**Nucleon
Reverse
Engineering**

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access

DVCS Kinematics
Universality tests
Towards 3D images

Building





Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Exclusive processes of current interest (1/2). Factorization and universality.

**Nucleon
Reverse
Engineering**

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access

DVCS Kinematics
Universality tests
Towards 3D images

Building





Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access

DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Bjorken regime : large Q^2 and fixed $xB \simeq 2\xi/(1 + \xi)$

- Partonic interpretation relies on **factorization theorems**.
- All-order proofs for DVCS, TCS and some DVMP.
- GPDs depend on a (arbitrary) factorization scale μ_F .
- **Consistency** requires the study of **different channels**.

- GPDs enter DVCS through **Compton Form Factors** :

$$\mathcal{F}(\xi, t, Q^2) = \int_{-1}^1 dx C\left(x, \xi, \alpha_S(\mu_F), \frac{Q}{\mu_F}\right) F(x, \xi, t, \mu_F)$$

for a given GPD F .

- CFF \mathcal{F} is a **complex function**.

Need for global fits of world data.

Different facilities will probe different kinematic domains.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix


Experimental data collected at
3 facilities




Need for global fits of world data.

Different facilities will probe different kinematic domains.

Valence quarks



Experimental data collected at 3 facilities



Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Need for global fits of world data.

Different facilities will probe different kinematic domains.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access

DVCS Kinematics

Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Valence quarks



Experimental data collected at 3 facilities




DESY •

CERN •

Thomas
Jefferson
National
Laboratory


Sea quarks




Need for global fits of world data.

Different facilities will probe different kinematic domains.

Valence quarks



Experimental data collected at 3 facilities, soon 4: EIC !




DESY •

CERN •

Thomas
Jefferson
National
Laboratory

Sea quarks



Gluons

NSAC, Long Range Plan 2015:
"We recommend [...] EIC as the highest
priority for new facility construction"

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon
Reverse
Engineering



- Study the **harmonic structure** of $ep \rightarrow ep\gamma$ amplitude.

Diehl *et al.*,

Phys. Lett. **B411**, 193 (1997)

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Experiment	Kinematics		
	x_B	Q^2 [GeV ²]	t [GeV ²]
HERA	0.001	8.00	-0.30
COMPASS	0.05	2.00	-0.20
HERMES	0.09	2.50	-0.12
CLAS	0.19	1.25	-0.19
HALL A	0.36	2.30	-0.23

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Beam Charge Asymmetry, HERMES



Kroll *et al.*, Eur. Phys. J. **C73**, 2278 (2013)

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics

Universality tests

Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Beam Spin Asymmetry, CLAS



Kroll *et al.*, Eur. Phys. J. **C73**, 2278 (2013)

**Nucleon
Reverse
Engineering**

QCD

- Mass without mass
- Nucleon structure
- Content of GPDs

Imaging

Experimental access
DVCS Kinematics

Universality tests

Towards 3D images

Building

- Computing chain
- Examples
- Team

Learning

- Definition
- Overlap
- Radon transform
- Covariant extension

Conclusion

Appendix

- **Dominance** of twist-2 and **validity** of a GPD analysis of DVCS data.
- $Im\mathcal{H}$ **best determined**. Large uncertainties on $Re\mathcal{H}$.
- However sizable **higher twist contamination** for DVCS measurements.
- Already some indications about the **invalidity** of the H -dominance hypothesis with **unpolarized data**.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests

Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension


Conclusion

Appendix

1. Experimental data fits



2. GPD extraction



3. Nucleon imaging

Images from Guidal et al.,
Rept. Prog. Phys. 76 (2013) 066202


Reaching for the Horizon

The 2015 Long Range Plan for Nuclear Science

Sidebar 2.2: The First 3D Pictures of the Nucleon

A computed tomography (CT) scan can help physicians pinpoint minute cancer tumors, diagnose tiny broken bones, and spot the early signs of osteoporosis. Now physicists are using the principles behind the procedure to peer at the inner workings of the proton. This breakthrough is made possible by a relatively new concept in nuclear physics called generalized parton distributions.

An intense beam of high-energy electrons can be used



Nucleon Reverse Engineering

1 Extract $H(x, \xi, t, \mu_F^{\text{ref}})$ from experimental data.

2 Extrapolate to vanishing skewness $H(x, 0, t, \mu_F^{\text{ref}})$.

3 Extrapolate $H(x, 0, t, \mu_F^{\text{ref}})$ up to infinite t .

4 Compute 2D Fourier transform in transverse plane:

$$H(x, b_{\perp}) = \int_0^{+\infty} \frac{d|\Delta_{\perp}|}{2\pi} |\Delta_{\perp}| J_0(|b_{\perp}||\Delta_{\perp}|) H(x, 0, -\Delta_{\perp}^2)$$

5 Propagate uncertainties.

6 Control extrapolations with an accuracy matching that of experimental data with **sound** GPD models.

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests

Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

**Nucleon
Reverse
Engineering**

- Evaluation of the impact of higher order effects.

QCD

- Mass without mass
- Nucleon structure
- Content of GPDs

- Evaluation of the impact of target mass and finite- t corrections.

Imaging

- Experimental access
- DVCS Kinematics
- Universality tests

Towards 3D images

- Extrapolations with **GPD** models.

Building

- Computing chain
- Examples
- Team

Learning

- Definition
- Overlap
- Radon transform
- Covariant extension

- Evaluation of the contribution of **higher twist** GPDs.
- DVMP: sensitivity to **DA models**.

Conclusion

Appendix

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests

Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests

Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Building the tools for high precision: the PARTONS project



PARtonic
Tomography
Of
Nucleon
Software

Nucleon Reverse Engineering

Experimental
data and
phenomenology

Full processes

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Computation
of amplitudes

Small distance
contributions

Building

Computing chain

Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

First
principles and
fundamental
parameters

Large distance
contributions

Conclusion

Appendix

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Experimental data and phenomenology

Full processes

Computation of amplitudes

Small distance contributions

First principles and fundamental parameters

Large distance contributions

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension


Conclusion

Appendix

Experimental data and phenomenology

Computation of amplitudes

First principles and fundamental parameters



Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension


Conclusion

Appendix

Experimental data and phenomenology

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Experimental data and phenomenology

Need for modularity

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

- Perturbative approximations.
- Physical models.
- Fits.
- Numerical methods.
- Accuracy and speed.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Experimental data and phenomenology

Need for modularity

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

- Perturbative approximations.
- **Physical models.**
- Fits.
- Numerical methods.
- Accuracy and speed.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Experimental data and phenomenology

Need for modularity

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

- Perturbative approximations.
- Physical models.
- Fits.
- Numerical methods.
- Accuracy and speed.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Experimental data and phenomenology

Need for modularity

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

- Perturbative approximations.
- Physical models.
- Fits.
- Numerical methods.
- Accuracy and speed.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

Experimental data and phenomenology

Need for modularity

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

- Perturbative approximations.
- Physical models.
- Fits.
- Numerical methods.
- Accuracy and speed.

Nucleon Reverse Engineering

- 3 stages:
 - 1 Design.
 - 2 Integration and validation.
 - 3 Benchmarking and production.

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

- Flexible software architecture.

B. Berthou *et al.*, *PARTONS: a computing platform for the phenomenology of Generalized Parton Distributions*
arXiv:1512.06174, to appear in *Eur. Phys. J. C*.

▶ See more on software architecture.

- 1 new physical development = 1 new module.
- *Aggregate* **knowledge** and **know-how**:
 - Models
 - Measurements
 - Numerical techniques
 - Validation

- What *can* be automated *will be* automated.

Nucleon Reverse Engineering

Coefficient functions, from EIC to Jefferson Lab kinematics

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain


Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



■ *Work in progress!*

■ Validation and preparation of **nonregression** tools.

■ **Flexibility** at work: **physical** models and **numerical** techniques.

■ $\simeq 10^5$ GPDs computed in $\lesssim 1'$.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

```

_____ gpdExample() _____
1 // Lots of includes
2 #include <src/Partons.h>
3 ...
4 // Retrieve GPD service
5 GPDService* pGPDService = Partons::getInstance()->getServiceObjectRegistry
   ()->getGPDService();
6 // Load GPD module with the BaseModuleFactory
7 GPDModule* pGK11Model = Partons::getInstance()->getModuleObjectFactory
   ()->newGPDModule(GK11Model::classId);
8 // Create a GPDKinematic(x, xi, t, MuF, MuR) to compute
9 GPDKinematic gpdKinematic(0.1, 0.00050025, -0.3, 8., 8.);
10 // Compute data and store results
11 GPDResult gpdResult = pGPDService->
   computeGPDModelRestrictedByGPDType(gpdKinematic, pGK11Model,
   GPDType::ALL);
12 // Print results
13 std::cout << gpdResult.toString() << std::endl;
14
15 delete pGK11Model;
16 pGK11Model = 0;

```

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

```

_____ computeOneGPD.xml _____
1 <?xml version="1.0" encoding="UTF-8" standalone="yes" ?>
2 <scenario id="01" date="" description="Example of computation of one GPD
   model (GK11) without evolution">
3     <!-- Select type of computation -->
4     <task service="GPDSservice" method="computeGPDModel" >
5         <!-- Specify kinematics -->
6         <GPDKinematic>
7             <param name="x" value="0.1" />
8             <param name="xi" value="1.00050025" />
9             <param name="t" value="-0.3" />
10            <param name="MuF2" value="8" />
11            <param name="MuR2" value="8" />
12        </GPDKinematic>
13        <!-- Choose GPD model and set parameters -->
14        <GPDModule>
15            <param name="id" value="GK11Model" />
16        </GPDModule>
17    </task>
18 </scenario>

```

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

```

_____ computeOneGPD
1 <?xml version="1.0" encoding="UTF-8" stand
2 <scenario id="01" date="" description="Exam
   _model_(GK11)_without_evolution">
3     <!-- Select type of computation -->
4     <task service="GPDSservice" method=
5       <!-- Specify kinematics -->
6       <GPDKinematic>
7         <param name="x" valu
8         <param name="xi" va
9         <param name="t" valu
10        <param name="MuF2"
11        <param name="MuR2"
12      </GPDKinematic>
13    <!-- Choose GPD model and
14    <GPDModule>
15      <param name="id" va
16    </GPDModule>
17  </task>
18 </scenario>

```

$$H^u = 0.822557$$

$$H^{u(+)} = 0.165636$$

$$H^{u(-)} = 1.47948$$

$$H^d = 0.421431$$

$$H^{d(+)} = 0.0805182$$

$$H^{d(-)} = 0.762344$$

$$H^s = 0.00883408$$

$$H^{s(+)} = 0.0176682$$

$$H^{s(-)} = 0$$

$$H^g = 0.385611$$

$$\text{and } E, \tilde{H}, \tilde{E}, \dots$$

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

```

computeOneCFF.xml
1 <?xml version="1.0" encoding="UTF-8" standalone="yes" ?>
2 <scenario id="03" date="" description="Example of one
   convol_coeff_function_model (DVCSConvCoeff) with GPD_model (GK11)">
3   <task service="DVCSConvCoeffFunctionService" method="
computeWithGPDModel"
4     <DVCSConvCoeffFunctionKinematic>
5       <param name="xi" value="0.5" />
6       <param name="t" value="-0.1346" />
7       <param name="Q2" value="1.5557" />
8       <param name="MuF2" value="4" />
9       <param name="MuR2" value="4" />
10    </DVCSConvCoeffFunctionKinematic>
11    <GPDMModule>
12      <param name="id" value="GK11Model" />
13    </GPDMModule>
14    <DVCSConvCoeffFunctionModule>
15      <param name="id" value="DVCSConvCoeffModel" />
16      <param name="qcd_order_type" value="LO" />
17    </DVCSConvCoeffFunctionModule>
18  </task>
19 </scenario>

```

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

computeOneCFF.xml

```

1 <?xml version="1.0" encoding="UTF-8" standalone="yes" ?>
2 <scenario id="03" date="" description="Example of computation of one
   convol coeff function model (DVCS CFF) with GPD model (GK11)">
3   <task service="DVCSConvolCoeffFunctionService" method="
   computeWithGPDModel"
4     <DVCSConvolCoeffFunctionKinematic>
5       <param name="xi" value="0.5" />
6       <param name="t" value="-0.1346" />
7       <param name="Q2" value="1.5557" />
8       <param name="MuF2" value="4" />
9       <param name="MuR2" value="4" />
10    </DVCSConvolCoeffFunctionKinematic>
11    <GPDMModule>
12      <param name="id" value="GK11Model" />
13    </GPDMModule>
14    <DVCSConvolCoeffFunction>
15      <param name="xi" value="0.5" />
16      <param name="t" value="-0.1346" />
17    </DVCSConvolCoeffFunction>
18  </task>
19 </scenario>

```

$$\begin{aligned}
 \mathcal{H} &= 1.47722 + 1.76698 i \\
 \mathcal{E} &= 0.12279 + 0.512312 i \\
 \tilde{\mathcal{H}} &= 1.54911 + 0.953728 i \\
 \tilde{\mathcal{E}} &= 18.8776 + 3.75275 i
 \end{aligned}$$

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain

Examples

Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

```

_____ computeManyKinematicsOneModel.xml _____
1 <scenario id="5" date="" description="Compute_Al_u_E1-DVCS_kinematics">
2   <task service="ObservableService" method="
computeManyKinematicOneModel" storeInDB="0">
3     <ObservableKinematic>
4       <param name="file"
5         value="/home/debian/workspace/PARTONS/data/E1DVCS.dat" />
6     </ObservableKinematic>
7     <Observable>
8       <param name="id" value="Alu" />
9     </Observable>
10    <DVCSModule>
11      <param name="id" value="BMJ2012Model" />
12      <param name="beam_energy" value="5.75" />
13    </DVCSModule>
14    <DVCSConvolCoeffFunctionModule>
15      <param name="id" value="DVCS_CFF_Model" />
16      <param name="qcd_order_type" value="LO" />
17    </DVCSConvolCoeffFunctionModule>
18    <GPDMModule>
19      <param name="id" value="GK11Model" />
20    </GPDMModule>
21  </task>

```

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples

Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

U. Paris-Saclay



Berthou



Chouika



Guidal



Lafitte



Moutarde



Sabatié



Sznajder

NCBJ



Wagner

ANL



Mezrag

ANL

U. Conn

U. Paris
Saclay

NCBJ

ECT*

U. Huelva

U. Conn



Colaneri



Joo

U. Huelva



Rodríguez-Quintero

ECT*/FBK



Binosi



Learning on the strong interaction from GPD models

Spin-0 Generalized Parton Distribution.

Definition and simple properties.

Nucleon
Reverse
Engineering

$$H_{\pi}^q(x, \xi, t) = \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \left\langle \pi, P + \frac{\Delta}{2} \left| \bar{q} \left(-\frac{z}{2} \right) \gamma^+ q \left(\frac{z}{2} \right) \right| \pi, P - \frac{\Delta}{2} \right\rangle_{\substack{z^+=0 \\ z_{\perp}=0}}$$

QCD


Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team



■ PDF forward limit

References

Müller *et al.*, Fortschr. Phys. **42**, 101 (1994)
Ji, Phys. Rev. Lett. **78**, 610 (1997)
Radyushkin, Phys. Lett. **B380**, 417 (1996)

$$H^q(x, 0, 0) = q(x)$$

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Spin-0 Generalized Parton Distribution.

Definition and simple properties.

Nucleon
Reverse
Engineering

$$H_{\pi}^q(x, \xi, t) = \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \left\langle \pi, P + \frac{\Delta}{2} \left| \bar{q} \left(-\frac{z}{2} \right) \gamma^+ q \left(\frac{z}{2} \right) \right| \pi, P - \frac{\Delta}{2} \right\rangle_{z_{\perp}=0}^{z_{\perp}=0}$$

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning


Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

with $t = \Delta^2$ and $\xi = -\Delta^+/(2P^+)$.



References

Müller *et al.*, Fortschr. Phys. **42**, 101 (1994)
Ji, Phys. Rev. Lett. **78**, 610 (1997)
Radyushkin, Phys. Lett. **B380**, 417 (1996)

- PDF forward limit
- Form factor sum rule

$$\int_{-1}^{+1} dx H^q(x, \xi, t) = F_1^q(t)$$

Spin-0 Generalized Parton Distribution.

Definition and simple properties.

Nucleon
Reverse
Engineering

$$H_{\pi}^q(x, \xi, t) = \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \left\langle \pi, P + \frac{\Delta}{2} \left| \bar{q} \left(-\frac{z}{2} \right) \gamma^+ q \left(\frac{z}{2} \right) \right| \pi, P - \frac{\Delta}{2} \right\rangle_{\substack{z^+=0 \\ z_{\perp}=0}}$$

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning


Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

with $t = \Delta^2$ and $\xi = -\Delta^+/(2P^+)$.



References

Müller *et al.*, Fortschr. Phys. **42**, 101 (1994)
Ji, Phys. Rev. Lett. **78**, 610 (1997)
Radyushkin, Phys. Lett. **B380**, 417 (1996)

- PDF forward limit
- Form factor sum rule
- H^q is an **even function** of ξ from time-reversal invariance.

Spin-0 Generalized Parton Distribution.

Definition and simple properties.

Nucleon
Reverse
Engineering

$$H_{\pi}^q(x, \xi, t) = \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \left\langle \pi, P + \frac{\Delta}{2} \left| \bar{q} \left(-\frac{z}{2} \right) \gamma^+ q \left(\frac{z}{2} \right) \right| \pi, P - \frac{\Delta}{2} \right\rangle_{z_{\perp}=0}^{z_{\perp}=0}$$

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning


Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

with $t = \Delta^2$ and $\xi = -\Delta^+/(2P^+)$.



References

Müller *et al.*, Fortschr. Phys. **42**, 101 (1994)
Ji, Phys. Rev. Lett. **78**, 610 (1997)
Radyushkin, Phys. Lett. **B380**, 417 (1996)

- PDF forward limit
- Form factor sum rule
- H^q is an **even function** of ξ from time-reversal invariance.
- H^q is **real** from hermiticity and time-reversal invariance.

Nucleon Reverse Engineering

■ Polynomiality

$$\int_{-1}^{+1} dx x^n H^q(x, \xi, t) = \text{polynomial in } \xi$$

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon
Reverse
Engineering

■ Polynomiality

Lorentz covariance

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon
Reverse
Engineering

■ Polynomiality

Lorentz covariance

■ Positivity

$$H^q(x, \xi, t) \leq \sqrt{q\left(\frac{x+\xi}{1+\xi}\right) q\left(\frac{x-\xi}{1-\xi}\right)}$$

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

■ H^q has support $x \in [-1, +1]$.

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

■ H^q has support $x \in [-1, +1]$.

Relativistic quantum mechanics

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon
Reverse
Engineering

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

■ H^q has support $x \in [-1, +1]$.

Relativistic quantum mechanics

■ Soft pion theorem (pion target)

$$H^q(x, \xi = 1, t = 0) = \frac{1}{2} \phi_\pi^q \left(\frac{1+x}{2} \right)$$

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

■ H^q has support $x \in [-1, +1]$.

Relativistic quantum mechanics

■ Soft pion theorem (pion target)

Dynamical chiral symmetry breaking

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

■ H^q has support $x \in [-1, +1]$.

Relativistic quantum mechanics

■ Soft pion theorem (pion target)

Dynamical chiral symmetry breaking

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

How can we implement *a priori* these theoretical constraints?

- There is no known GPD parameterization **relying only on first principles**.
- In the following, focus on **polynomiality** and **positivity**.

Nucleon Reverse Engineering

■ Representation of GPD:

$$H^q(x, \xi, t) = \int_{\Omega_{DD}} d\beta d\alpha \delta(x - \beta - \alpha\xi) (F^q(\beta, \alpha, t) + \xi G^q(\beta, \alpha, t))$$

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

- Support property: $x \in [-1, +1]$.
- Discrete symmetries: F^q is α -even and G^q is α -odd.
- **Gauge:** any representation (F^q, G^q) can be recast in one representation with a single DD f^q :

$$H^q(x, \xi, t) = x \int_{\Omega_{DD}} d\beta d\alpha f_{\text{BMKS}}^q(\beta, \alpha, t) \delta(x - \beta - \alpha\xi)$$

Belitsky *et al.*, Phys. Rev. **D64**, 116002 (2001)

$$H^q(x, \xi, t) = (1 - x) \int_{\Omega_{DD}} d\beta d\alpha f_{\text{P}}^q(\beta, \alpha, t) \delta(x - \beta - \alpha\xi)$$

Pobylitsa, Phys. Rev. **D67**, 034009 (2003)

Müller, Few Body Syst. **55**, 317 (2014)

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

- Choose $F^q(\beta, \alpha) = 3\beta\theta(\beta)$ ad $G^q(\beta, \alpha) = 3\alpha\theta(\beta)$:

$$H^q(x, \xi) = 3x \int_{\Omega} d\beta d\alpha \delta(x - \beta - \alpha\xi)$$

- Simple analytic expressions for the GPD:

$$H(x, \xi) = \frac{6x(1-x)}{1-\xi^2} \text{ if } 0 < |\xi| < x < 1,$$

$$H(x, \xi) = \frac{3x(x+|\xi|)}{|\xi|(1+|\xi|)} \text{ if } -|\xi| < x < |\xi| < 1.$$

■ Compute first Mellin moments.

n	$\int_{-\xi}^{+\xi} dx x^n H(x, \xi)$	$\int_{+\xi}^{+1} dx x^n H(x, \xi)$	$\int_{-\xi}^{+1} dx x^n H(x, \xi)$
0	$\frac{1+\xi-2\xi^2}{1+\xi}$	$\frac{2\xi^2}{1+\xi}$	1
1	$\frac{1+\xi+\xi^2-3\xi^3}{2(1+\xi)}$	$\frac{2\xi^3}{1+\xi}$	$\frac{1+\xi^2}{2}$
2	$\frac{3(1-\xi)(1+2\xi+3\xi^2+4\xi^3)}{10(1+\xi)}$	$\frac{6\xi^4}{5(1+\xi)}$	$\frac{3(1+\xi^2)}{10}$
3	$\frac{1+\xi+\xi^2+\xi^3+\xi^4-5\xi^5}{5(1+\xi)}$	$\frac{6\xi^5}{5(1+\xi)}$	$\frac{1+\xi^2+\xi^4}{5}$
4	$\frac{1+\xi+\xi^2+\xi^3+\xi^4+\xi^5-6\xi^6}{7(1+\xi)}$	$\frac{6\xi^6}{7(1+\xi)}$	$\frac{1+\xi^2+\xi^4}{7}$

■ Expressions get more complicated as n increases... But they always yield polynomials!

**Nucleon
Reverse
Engineering**

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

- Decompose an hadronic state $|H; P, \lambda\rangle$ in a Fock basis:

$$|H; P, \lambda\rangle = \sum_{N, \beta} \int [dx d\mathbf{k}_\perp]_N \psi_N^{(\beta, \lambda)}(x_1, \mathbf{k}_{\perp 1}, \dots, x_N, \mathbf{k}_{\perp N}) |\beta, k_1, \dots, k_N\rangle$$

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

- Derive an expression for the pion GPD in the DGLAP region $\xi \leq x \leq 1$:

$$H^q(x, \xi, t) \propto \sum_{\beta, j} \int [d\bar{x} d\bar{\mathbf{k}}_\perp]_N \delta_{j, q} \delta(x - \bar{x}_j) (\psi_N^{(\beta, \lambda)})^*(\hat{x}', \hat{\mathbf{k}}'_\perp) \psi_N^{(\beta, \lambda)}(\tilde{x}, \tilde{\mathbf{k}}_\perp)$$

Building

Computing chain
Examples
Team

with $\tilde{x}, \tilde{\mathbf{k}}_\perp$ (resp. $\hat{x}', \hat{\mathbf{k}}'_\perp$) generically denoting incoming (resp. outgoing) parton kinematics.

Learning

Definition

Overlap

Radon transform
Covariant extension

Diehl *et al.*, Nucl. Phys. **B596**, 33 (2001)

- Similar expression in the ERBL region $-\xi \leq x \leq \xi$, but with overlap of N - and $(N+2)$ -body LFWFs.

Conclusion

Appendix

Nucleon Reverse Engineering

- Physical picture.
- Positivity relations are fulfilled **by construction**.
- Implementation of **symmetries of N -body problems**.

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition

Overlap

Radon transform
Covariant extension

Conclusion

Appendix

What is not obvious anymore

What is *not* obvious to see from the wave function representation is however the **continuity of GPDs at $x = \pm\xi$** and the **polynomiality** condition. In these cases both the DGLAP and the ERBL regions must cooperate to lead to the required properties, and this implies **nontrivial relations between the wave functions** for the different Fock states relevant in the two regions. An *ad hoc* Ansatz for the wave functions would **almost certainly lead** to GPDs that **violate the above requirements**.

Diehl, Phys. Rept. **388**, 41 (2003)

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap

Radon transform
Covariant extension

Conclusion

Appendix



$$\mathcal{R}f(s, \phi) = \int_{-\infty}^{+\infty} d\beta d\alpha f(\beta, \alpha) \delta(s - \beta \cos \phi - \alpha \sin \phi)$$

For $s > 0$ and $\phi \in [0, 2\pi]$:

and:

$$\mathcal{R}f(-s, \phi) = \mathcal{R}f(s, \phi \pm \pi)$$

Relation to GPDs:

$$x = \frac{s}{\cos \phi} \text{ and } \xi = \tan \phi$$

Relation between GPD and DD in Belitsky *et al.* gauge

$$\frac{\sqrt{1 + \xi^2}}{x} H(x, \xi) = \mathcal{R}f_{\text{BMKS}}(s, \phi),$$

**Nucleon
Reverse
Engineering**

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning


Definition
Overlap

Radon transform

Covariant extension

Conclusion

Appendix



$$\mathcal{R}f(s, \phi) = \int_{-\infty}^{+\infty} d\beta d\alpha f(\beta, \alpha) \delta(s - \beta \cos \phi - \alpha \sin \phi)$$

For $s > 0$ and $\phi \in [0, 2\pi]$:

and:

$$\mathcal{R}f(-s, \phi) = \mathcal{R}f(s, \phi \pm \pi)$$

Relation to GPDs:

$$x = \frac{s}{\cos \phi} \text{ and } \xi = \tan \phi$$

Relation between GPD and DD in Pobylitsa gauge

$$\frac{\sqrt{1 + \xi^2}}{1 - x} H(x, \xi) = \mathcal{R}f_P(s, \phi),$$

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap

Radon transform

Covariant extension

Conclusion

Appendix

- The Mellin moments of a Radon transform are **homogeneous polynomials** in $\omega = (\sin \phi, \cos \phi)$.
- The converse is also true:

Theorem (Hertle, 1983)

*Let $g(s, \omega)$ an even compactly-supported distribution. Then g is itself the Radon transform of a compactly-supported distribution if and only if the **Ludwig-Helgason consistency condition** hold:*


- (i) g is C^∞ in ω ,
- (ii) $\int ds s^m g(s, \omega)$ is a homogeneous polynomial of degree m for all integer $m \geq 0$.

- Double Distributions and the Radon transform are the **natural solution** of the polynomiality condition.

DGLAP and ERBL regions

$$(x, \xi) \in \text{DGLAP} \Leftrightarrow |s| \geq |\sin \phi| ,$$

$$(x, \xi) \in \text{ERBL} \Leftrightarrow |s| \leq |\sin \phi| .$$



- Each point (β, α) with $\beta \neq 0$ contributes to **both** DGLAP and ERBL regions.
- Expressed in **support theorem**.

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Ill-posedness in the sense of Hadamard.

A first glimpse at the inverse Radon transform.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

- Numerical evaluation *almost unavoidable* (polar vs cartesian coordinates).
- Ill-posedness by **lack of continuity**.
- The **unlimited** Radon inverse problem is **mildly** ill-posed while the **limited** one is **severely** ill-posed.
- Careful selection of **algorithms** and **numerical methods**.



Mezrag
PhD dissertation

► See more on inverse Radon transform.

Conclusion

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

- What makes hadron structure studies so interesting:
 - Deep **physical questions** waiting for answers!
 - Well-defined **theoretical framework** and **observables**.
 - Active **experimental programs** worldwide.
- **Challenging constraints** expected from:
 - Jefferson Lab in the valence sector,
 - CERN in the sea sector,
 - EIC (later) in the gluon sector.
- **Good theoretical control** on the path between GPD models and experimental data: from theory to measurements, and conversely.
- Development of the PARTONS framework for **phenomenology** and **theory** purposes.
- **First release** of PARTONS in summer 2016!

Appendix

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Local fits

Take each kinematic bin independantly of the others.
Extraction of $Re\mathcal{H}$, $Im\mathcal{H}$, ...as independent parameters.

Global fit

Take all kinematic bins at the same time. Use a parametrization of GPDs or CFFs.

Hybrid : Local / global fit

Start from local fits and add smoothness assumption.

Neural networks

Exploratory stage for GPDs.

Nucleon Reverse Engineering

Local fits

Take each kinematic bin independantly of the others.
Extraction of $Re\mathcal{H}$, $Im\mathcal{H}$, ...as independent parameters.

M. Guidal, Eur. Phys. J. **A39**, 5 (2009)

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

- **Almost model-independent:** relies on twist-2 dominance assumption and assume bounds for the fitting domain.
- Interpretation of **uncertainties** on extracted quantities?
Contributions from measurements uncertainties, correlations between CFFs and fitting domain boundaries.
- Interpretation of **extracted quantities**? e.g.mixing of quark and gluon GPDs due to NLO effects.
- **Oscillations** between different (x_B, t, Q^2) bins may happen.
- **Extrapolation** problem left open.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Local fits: What can be achieved in principle?

- Structure of BSA at twist 2 :

$$\text{BSA}(\phi) = \frac{a \sin \phi + b \sin 2\phi}{1 + c \cos \phi + d \cos 2\phi + e \cos 3\phi}$$

where $a = \mathcal{O}(Q^{-1})$, $b = \mathcal{O}(Q^{-4})$, $c = \mathcal{O}(Q^{-1})$,
 $d = \mathcal{O}(Q^{-2})$, $e = \mathcal{O}(Q^{-5})$.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Local fits: What can be achieved in principle?

- Structure of BSA at twist 2 :

$$\text{BSA}(\phi) = \frac{a \sin \phi + b \sin 2\phi}{1 + c \cos \phi + d \cos 2\phi + e \cos 3\phi}$$

- **Underconstrained** problem (8 fit parameters : real and imaginary parts of 4 CFFs \mathcal{H} , \mathcal{E} , $\tilde{\mathcal{H}}$ and $\tilde{\mathcal{E}}$).

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Local fits: What can be achieved in principle?

- Structure of BSA at twist 2 :

$$\text{BSA}(\phi) = \frac{a \sin \phi + b \sin 2\phi}{1 + c \cos \phi + d \cos 2\phi + e \cos 3\phi}$$

- **Underconstrained** problem.
- Need other asymmetries on **same** kinematic bin to allow extraction of **all CFFs** (or **add** $\simeq 5\text{-}10\%$ **systematic uncertainty**).

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Local fits: What can be achieved in principle?

- Structure of BSA at twist 2 :

$$\text{BSA}(\phi) = \frac{a \sin \phi + b \sin 2\phi}{1 + c \cos \phi + d \cos 2\phi + e \cos 3\phi}$$

- **Underconstrained** problem.
- Need other asymmetries on **same** kinematic bin to allow extraction of **all CFFs**.
- Add physical input? **Dispersion relations**, etc.

Kumericki *et al.*, arXiv:1301.1230

Guidal *et al.*, Rept. Prog. Phys. **76**, 066202 (2013)

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Global fit

Take all kinematic bins at the same time. Use a parametrization of GPDs or CFFs.

Kumericki, Nucl. Phys. **B841**, 1 (2010)

- **Model-dependent** approach.
- Allows the **implementation of theoretical constraints** on GPDs or CFFs.
- Guideline for **extrapolation** outside the physical domain.
- Compromise between number of parameters and number of described GPDs (flavor dependence, higher-twists, ...)?
- Impact on the **choice of a fitting strategy**?

Nucleon Reverse Engineering

Hybrid : Local / global fit

Start from local fits and add smoothness assumption.

Moutarde, Phys. Rev. **D79**, 094021 (2009)

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

- Avoid unphysical oscillations between different (x_B, t, Q^2) bins by comparing to a **global fit by a smooth function**:

$$H^+ = 2 \sum_{n=0}^N \sum_{l=0}^{n+1} B_{nl}(t) \theta(|x| < \xi) \left(1 - \frac{x^2}{\xi^2}\right) C_{2n+1}^{(3/2)}\left(\frac{x}{\xi}\right) P_{2l}\left(\frac{x}{\xi}\right)$$

- Number of fit parameters describing the B_{nl} coefficients **increases with N^2** ...Extension to other GPDs seems difficult.
- **Extrapolation** problem left open.

**Nucleon
Reverse
Engineering**

Neural networks

Exploratory stage for GPDs.

Kumericki *et al.*, JHEP **1107**, 073 (2011)

QCD

- Mass without mass
- Nucleon structure
- Content of GPDs

Imaging

- Experimental access
- DVCS Kinematics
- Universality tests
- Towards 3D images

Building

- Computing chain
- Examples
- Team


Learning

- Definition
- Overlap
- Radon transform
- Covariant extension

Conclusion

Appendix

- Already used for PDF fits.
- **Almost model-independent:** neural network description, twist-2, H -dominance?
- Good agreement between model fit and neural network fit in the fitting domain.
- **More reliable uncertainties** in extrapolations?
- **Overtraining** as a generic feature of (too) flexible models.

$$\mathcal{H} \text{ at LO and NLO } (t = -0.1 \text{ GeV}^2, Q^2 = \mu_F^2 = 4. \text{ GeV}^2)$$


Moutarde *et al.*, Phys. Rev. **D87**, 054029 (2013)

- **Systematic** tests of perturbative QCD assumptions.
- **Wide kinematic range** (from JLab to EIC).
- **Accuracy** set by JLab 12 GeV expected statistical accuracy.
- **Model dependent** evaluations.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

What is large Q^2 ?



■ World data cover **complementary kinematic regions.**

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion


Appendix

What is large Q^2 ?




- World data cover **complementary kinematic regions**.
- Q^2 is **not so large** for most of the data.

What is large Q^2 ?



- World data cover **complementary kinematic regions**.
 - Q^2 is **not so large** for most of the data.
 - Higher twists?**
- ◀ Back to challenge

What is large Q^2 ?



- World data cover **complementary kinematic regions**.
 - Q^2 is **not so large** for most of the data.
 - Higher twists?**
- ◀ Back to challenge

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

- Write dispersion relation **at fixed t and Q^2** :

$$\text{Re}\mathcal{H}(\xi, t) = \Delta(t) + \frac{2}{\pi} \mathcal{P} \int_0^1 \frac{dx}{x} \frac{\text{Im}\mathcal{H}(x, t)}{\left(\frac{\xi^2}{x^2} - 1\right)}$$

- Use LO relation $\text{Im}\mathcal{H}(x, t) = \pi \left(H(x, x, t) - H(-x, x, t) \right)$.
- Up to the D-term form factor $\Delta(t)$, all the information accessible **at LO and fixed Q^2** is contained on the cross-over line.

Teryaev, hep-ph/0510031

Anikin and Teryaev, Phys. Rev. **D76**, 056007 (2007)

Diehl and Ivanov, Eur. Phys. J. **C52**, 919 (2007)

Dispersion relations and actual data.

Too few kinematic bins to provide model-independent constraints?

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Dispersion relations and actual data.

Too few kinematic bins to provide model-independent constraints?

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Dispersion relations and actual data.

Too few kinematic bins to provide model-independent constraints?

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension


Conclusion

Appendix



Nucleon
Reverse
Engineering

GPd H at $t = -0.23 \text{ GeV}^2$ and $Q^2 = 2.3 \text{ GeV}^2$.



GPd model: see Kroll *et al.*, Eur. Phys. J. **C73**, 2278 (2013)

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Need to know $H(x, \xi = 0, t)$ to do transverse plane imaging.



GPD model: see Kroll *et al.*, Eur. Phys. J. **C73**, 2278 (2013)

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

What is the physical region?



GPD model: see Kroll *et al.*, Eur. Phys. J. **C73**, 2278 (2013)

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

ξ_{\min} from finite beam energy.



GPD model: see Kroll *et al.*, Eur. Phys. J. **C73**, 2278 (2013)

Nucleon Reverse Engineering

ξ_{\max} from kinematic constraint on 4-momentum transfer.

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



GPD model: see Kroll *et al.*, Eur. Phys. J. **C73**, 2278 (2013)

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

The cross-over line $x = \xi$.



GPD model: see Kroll *et al.*, Eur. Phys. J. **C73**, 2278 (2013)

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

The black curve is what is needed for transverse plane imaging!



GPD model: see Kroll *et al.*, Eur. Phys. J. **C73**, 2278 (2013)

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team


Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Density plot of H at $t = -0.23 \text{ GeV}^2$ and $Q^2 = 2.3 \text{ GeV}^2$



GPD model: see Kroll *et al.*, Eur. Phys. J. **C73**, 2278 (2013)

◀ Back to challenges.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



- Steps of logic sequence in parent class.
- Model description and related mathematical methods in daughter class.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Modularity and layer structure.

Modifying one layer does not affect the other layers.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

For **any model of LFWF**, one has to address the following three questions:

- 1 Does the extension exist?
- 2 If it exists, is it unique?
- 3 How can we compute this extension?

Work in progress!

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Theorem

Let f be a compactly-supported locally summable function defined on \mathbb{R}^2 and $\mathcal{R}f$ its Radon transform.

Let $(s_0, \omega_0) \in \mathbb{R} \times S^1$ and U_0 an open neighborhood of ω_0 such that:

$$\text{for all } s > s_0 \text{ and } \omega \in U_0 \quad \mathcal{R}f(s, \omega) = 0.$$

Then $f(\mathbb{N}) = 0$ on the half-plane $\langle \mathbb{N} | \omega_0 \rangle > s_0$ of \mathbb{R}^2 .

Consider a GPD H being zero on the DGLAP region.

- Take ϕ_0 and s_0 s.t. $\cos \phi_0 \neq 0$ and $|s_0| > |\sin \phi_0|$.
- Neighborhood U_0 of ϕ_0 s.t. $\forall \phi \in U_0 \quad |\sin \phi| < |s_0|$.
- The underlying DD f has a zero Radon transform for all $\phi \in U_0$ and $s > s_0$ (DGLAP).
- Then $f(\beta, \alpha) = 0$ for all $(\beta, \alpha) \in \Omega_{\text{DD}}$ with $\beta \neq 0$.
- Extension **unique** up to adding a **D-term**: $\delta(\beta)D(\alpha)$.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

A discretized problem

Consider $N + 1$ Hilbert spaces H, H_1, \dots, H_N , and a family of continuous surjective operators $R_n : H \rightarrow H_n$ for $1 \leq n \leq N$. Being given $g_1 \in H_1, \dots, g_n \in H_n$, we search f solving the following system of equations:

$$R_n f = g_n \quad \text{for } 1 \leq n \leq N$$

Fully discrete case

Assume f piecewise-constant with values f_m for $1 \leq m \leq M$. For a collection of lines $(L_n)_{1 \leq n \leq N}$ crossing Ω_{DD} , the Radon transform writes:

$$g_n = \mathcal{R}f = \int_{L_n} f = \sum_{m=1}^M f_m \times \text{Measure}(L_n \cap C_m) \quad \text{for } 1 \leq n \leq N$$

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Kaczmarz algorithm

Denote P_n the orthogonal projection on the *affine* subspace $R_n f = g_n$. Starting from $f^0 \in H$, the sequence defined iteratively by:

$$f^{k+1} = P_N P_{N-1} \dots P_1 f^k$$

converges to the solution of the system.

The convergence is exponential if the projections are randomly ordered.

Strohmer and Vershynin, Jour. Four. Analysis and Appl. **15**,
437 (2009)

Computation of the extension.

Numerical evaluation of the inverse Radon transform (2/3).

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

And if the input data are inconsistent?

- Instead of solving $g = \mathcal{R}f$, find f such that $\|g - \mathcal{R}f\|_2$ is **minimum**.
- The solution **always exists**.
- The input data are **inconsistent** if $\|g - \mathcal{R}f\|_2 > 0$.

Nucleon Reverse Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Relaxed Kaczmarz algorithm

Let $\omega \in]0, 2[$ and:

$$P_n^\omega = (1 - \omega) \text{Id}_H + \omega P_n \quad \text{for } 1 \leq n \leq N$$

Write:

$$RR^\dagger = (R_i R_j^\dagger)_{1 \leq i, j \leq N} = D + L + L^\dagger$$

where D is diagonal, and L is lower-triangular with zeros on the diagonal.

Theorem

Let $0 < \omega < 2$. For $f^0 \in \text{Ran } R^\dagger$ (e.g. $f^0 = 0$), the Kaczmarz method with relaxation converges to the unique solution $f^\omega \in \text{Ran } R^\dagger$ of:

$$R^\dagger(D + \omega L)^{-1}(g - Rf^\omega) = 0 ,$$

where the matrix D and L appear in the decomposition of RR^\dagger . If $g = \mathcal{R}f$ has a solution, then f^ω is its solution of minimal norm. Otherwise:

$$f^\omega = f_{MP} + \mathcal{O}(\omega) ,$$

where f_{MP} is the minimizer in H of:

$$\langle g - \mathcal{R}f | g - \mathcal{R}f \rangle_D ,$$

the inner product being defined by:

$$\langle h | k \rangle_D = \langle D^{-1}h | k \rangle .$$

Nucleon
Reverse
Engineering

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building

Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix

Nucleon
Reverse
Engineering

A pion valence PDF-like example

Aim: reconstruct the PDF $q(x) = 30x^2(1 - x)^2$ from the knowledge of its first 30 Mellin moments.

QCD

Mass without mass
Nucleon structure
Content of GPDs

Imaging

Experimental access
DVCS Kinematics
Universality tests
Towards 3D images

Building


Computing chain
Examples
Team

Learning

Definition
Overlap
Radon transform
Covariant extension

Conclusion

Appendix



- Extensive testing *in progress*

- Various inputs: PDFs and LFWFs.
- Numerical noise.

- Piecewise-constant PDF: 20 values.
- Input: 30 Mellin moments.
- Unrelaxed method $\omega = 1$.
- 10000 iterations.

