

Cross section and polarization observables for the reaction

$$e^+ + e^- \rightarrow a_1(1260) + \pi$$

E. Tomasi-Gustafsson

*IRFU/Service de Physique Nucléaire,
CEA/Saclay, 91191 Gif-sur-Yvette, France*

G. I. Gakh

*National Science Centre "Kharkov Institute of Physics and Technology",
61108 Akademicheskaya 1, Kharkov, Ukraine*

C. Adamušćin

Institute of Physics, Slovak Academy of Sciences, Bratislava, Slovakia

(Dated: March 5, 2008)

Abstract

A model independent formalism for the electron positron annihilation reaction $e^+ + e^- \rightarrow a_1(1260) + \pi$ has been derived. The differential and total cross sections and the elements of the spin-density matrix of the a_1 -meson were calculated in terms of the electromagnetic form factors of the corresponding $\gamma^* a_1 \pi$ current. Simple models of a_1 form factors have been fitted to the available cross section data and they allow to give numerical predictions for the different observables.

PACS numbers: 12.20.-m, 13.40.-f, 13.60.-Hb, 13.88.+e

I. INTRODUCTION

The electron positron annihilation into hadrons constitutes an important source of information on the internal structure of the mesons: the light quarks and their interactions as well as the spectroscopy of their bound states. The experimental data about these reactions in the low-energy region are also relevant to the determination of the strong interaction contribution to the anomalous magnetic moment of the muon, to the test of standard model predictions for the hadronic tau-lepton decay, which is related by the conservation of vector currents.

Recently, the construction of the new detectors with a large solid angle, which can operate at new colliders with high luminosity, opened new possibilities for the investigation of the reactions $e^+ + e^- \rightarrow$ multihadrons [1]. Not only the statistic is highly increased, but also the possibility to detect charged as well as neutral pions allows to draw conclusions on the nature of the intermediate states.

In the energy region $1 \leq W \leq 2.5$ GeV (W is the total energy of the colliding beams) the process of four pion production is one of the dominant processes of the reaction $e^+ + e^- \rightarrow$ hadrons. Its cross section is larger than 2π production and comparable to $e^+ + e^- \rightarrow \mu^+ + \mu^-$.

The process of e^+e^- annihilation into four pions was firstly detected in Frascati [2] and later on in Novosibirsk [3]. Through a simultaneous analysis of the differential distributions in two final channels: $2\pi^+2\pi^-$ and $\pi^+\pi^-2\pi^0$, it was shown in [1] that the reaction predominantly occurs through the $a_1(1260)\pi$ and $\omega\pi^0$ intermediate states in the energy range 1.05–1.38 GeV. It was also found that the relative fraction of the $a_1(1260)\pi$ state increases with the beam energy. The measurement of the $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$ cross section was extended to lower energies. Data obtained with larger statistical and systematic precision [4] confirmed that the dominant production mechanism is consistent with the $a_1(1260)\pi$ intermediate state.

The process of the multihadron production at large energies was also investigated with the BABAR detector at the PEP-II asymmetric electron-positron storage ring using the initial-state radiation [5]. In particular, the cross section for the process $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$ was measured for center-of-mass (CMS) energies from 0.6 to 4.5 GeV, providing evidence of a resonant structure, with preferred quasi-two-body production of $a_1(1260)\pi$. A detailed

understanding of the four-pion final state requires also information from final states such as $\pi^+\pi^-\pi^0\pi^0$, to which the $\rho^+\rho^-$ intermediate state can contribute. A summary of the hadronic cross section measurements performed with BABAR via radiative return is given in Ref. [6].

In this paper we consider the reaction

$$e^-(k_1) + e^+(k_2) \rightarrow a_1(p_1) + \pi(p_2), \quad (1)$$

where a_1 is the axial-vector meson $a_1(1260)$ with the following quantum numbers $I^G(J^{PC}) = 1^-(1^{++})$. The notation for the particle four-momenta is given in brackets.

The determination of the pseudoscalar-meson FFs as pions and kaons requires only cross section measurements. They have been extensively studied both in the space-like and time-like regions (see, for example, [7]). Light vector mesons are less known, because their experimental determination is more difficult, due to their short lifetimes. However, the t dependence of the cross section for diffractive vector-meson electroproduction gives (model-dependent) information on the charge radius, and radiative decays, such as $\rho^+ \rightarrow \pi^+\pi^0\gamma$, allow to obtain their magnetic moment.

From the theoretical point of view, the processes of the vector- and axial-mesons production in the electron-positron annihilation were considered in a number of papers. The predictions for the differential and total cross sections of the reaction $e^+e^- \rightarrow a_1^\pm + \pi^\mp$ was given in Ref. [8] in the framework of the hard-pion current algebra models.

It was shown that the reaction cross section alone could be, in principle, discriminative toward models. Using VMD model, the authors of Ref. [9] investigated the reaction $e^+e^- \rightarrow mesons$ assuming two-body (or quasi-two-body) final states, as $a_1(1260)\pi$ and $\rho^+\rho^-$. All FFs were taken equal to unity. Estimations of the cross sections of the processes $e^+e^- \rightarrow 3\pi$, 4π were also obtained, using the VMD model, in Ref. [10].

It appears that the magnitude and energy dependence of the cross section alone can not constitute a decisive test on the validity of VMD models. A recent discussion can be found in Ref. [11], where existing models [12–14] have been phenomenologically modified, including parameters to be fitted on the data. A good description of the cross section is obtained assuming $a_1\pi$ intermediate state, in addition to ρ and π , and including higher ρ resonances for energies over 1 GeV.

Due to the conservation of vector current the cross section of the $e^+e^- \rightarrow 4\pi$ process can be related to the probability of the $\tau \rightarrow 4\pi\nu_\tau$ decay. Therefore, all realistic models describing

the first process, should also be applicable to the description of the latter one. It was found [15] that the assumption of the $a_1(1260)\pi$ dominance is in qualitative agreement with all available data. The free-parameter investigation of the branching ratios and distribution functions of the four particle decay of $\tau \rightarrow \rho\pi\pi\nu$, in terms of the effective chiral theory of mesons, is consistent with the data [16]. The theory predicted the a_1 dominance in these four particle decay of the tau-lepton.

The purpose of this paper is to calculate the differential (and total) cross sections and the elements of the spin-density matrix of the a_1 -meson in terms of the electromagnetic form factors (FFs) of the corresponding $\gamma^*a_1\pi$ current. A model independent formalism, derived in [17] for spin one particles, and applied to the process $e^+ + e^- \rightarrow \rho^+ + \rho^-$ in [18], allows to express the experimental observables (differential cross section, polarization observables, elements of the density matrix..) in terms of hadron FFs. In annihilation reactions, these FFs should be known, or extrapolated from the space-like region into the time-like (TL) region, on the basis of analytical arguments.

II. FORMALISM

The following derivation is based on the one-photon exchange mechanism. In principle, at large q^2 , one should take into account the two-photon-exchange contribution, as it was suggested a few decades ago [19]. In case of spin one particles, model independent properties of the two-photon-exchange contribution in elastic electron-deuteron scattering have been discussed in Refs. [20, 21]. However no experimental evidence has been found, up to now, on the presence of two-photon-exchange in the scattering [22] as well as in the annihilation channels [23], and we do not include such contribution which makes the formalism very complicated.

A. The spin structure of the matrix element

In the one-photon approximation, the differential cross section of the reaction (1) in terms of the hadronic, $W_{\mu\nu}$, and leptonic, $L_{\mu\nu}$, tensors, neglecting the electron mass, is written as

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{q^6} \frac{p}{2W} L_{\mu\nu} W_{\mu\nu}, \quad (2)$$

where $\alpha = 1/137$ is the electromagnetic constant, $p = \sqrt{(q^2 + m^2 - M^2)^2 - 4m^2q^2}/2W$ is the final-particle momentum in the reaction CMS, m and M are the masses of the pion and of the a_1 meson, respectively. The four momentum of the virtual photon is $q = k_1 + k_2 = p_1 + p_2$, with $q^2 = W^2$, and W is the total energy of the initial beams (note that the cross section is not averaged over the spins of the initial beams).

The leptonic tensor (for the case of longitudinally polarized electron beam) is

$$L_{\mu\nu} = -q^2 g_{\mu\nu} + 2(k_{1\mu}k_{2\nu} + k_{2\mu}k_{1\nu}) + 2i\lambda\varepsilon_{\mu\nu\sigma\rho}k_{1\sigma}k_{2\rho}, \quad (3)$$

where λ is the degree of the electron beam polarization (further we assume that the electron beam is completely polarized and consequently $\lambda = 1$).

The hadronic tensor can be expressed via the electromagnetic current J_μ , describing the transitions $\gamma^* \rightarrow \pi a_1$, as follows

$$W_{\mu\nu} = J_\mu J_\nu^*. \quad (4)$$

The hadron tensor $W_{\mu\nu}$ can be expressed in terms of FFs of the $\gamma^* \rightarrow \pi a_1$ transition, using the explicit form of the electromagnetic current J_μ . The spin-density matrix of the a_1 -meson is composed of three terms, corresponding to unpolarized, vector and tensor polarized meson:

$$\rho_{\mu\nu} = -\left(g_{\mu\nu} - \frac{p_{1\mu}p_{1\nu}}{M^2}\right) + \frac{i}{2M}\varepsilon_{\mu\nu\rho\sigma}s_\rho p_{1\sigma} + 3Q_{\mu\nu}. \quad (5)$$

Here s_μ and $Q_{\mu\nu}$ are the a_1 -meson polarization four vector and quadrupole tensor, respectively. The four vector of the a_1 -meson vector polarization s_μ and the a_1 -meson quadrupole-polarization tensor $Q_{\mu\nu}$ satisfy the following conditions:

$$s^2 = -1, \quad sp_1 = 0, \quad Q_{\mu\nu} = Q_{\nu\mu}, \quad Q_{\mu\mu} = 0, \quad p_{1\mu}Q_{\mu\nu} = 0.$$

Taking into account Eqs. (4) and (5), the hadronic tensor in the general case can be written as the sum of three terms

$$W_{\mu\nu} = W_{\mu\nu}(0) + W_{\mu\nu}(V) + W_{\mu\nu}(T), \quad (6)$$

where $W_{\mu\nu}(0)$ corresponds to the case of unpolarized particles in the final state and $W_{\mu\nu}(V)(W_{\mu\nu}(T))$ corresponds to the case of the vector (tensor) polarized a_1 -meson.

These expressions are general, for any spin one particle in the final state. Let us consider more particularly, the reaction $e^+ + e^- \rightarrow \pi + a_1$ which has been shown to be the main contribution to the 4π final state.

The electromagnetic current of the $\gamma^* \rightarrow \pi a_1$ transition is described by two FFs. Assuming the P- and C-invariance of the hadron electromagnetic interaction this current can be written as [10]

$$J_\mu = f_1(q^2)(q^2 U_\mu^* - q \cdot U^* q_\mu) + f_2(q^2)(q \cdot p_2 U_\mu^* - q \cdot U^* p_{2\mu}), \quad (7)$$

where U_μ is the polarization four-vector describing the spin one a_1 -meson, and $f_i(q^2)$, ($i = 1, 2$) are the electromagnetic FFs describing the $\gamma^* \rightarrow \pi a_1$ transition (note that we singled out explicitly the electron charge e from the expression for the electromagnetic current). The FFs $f_i(q^2)$ are complex functions of the variable q^2 in the region of the TL momentum transfer ($q^2 > 0$).

In case of real photon, f_1 does not contribute, and the value $f_2(0)$ can be obtained from the experimental data on the decay width $\Gamma(a_1 \rightarrow \pi\gamma)$. The expression of the radiative decay width Γ in the axial-vector meson rest frame is

$$\Gamma = \frac{1}{\pi} \frac{\omega}{24M^2} |\mathcal{M}|^2, \quad (8)$$

where ω is the photon momentum in the axial-vector meson rest frame, $\omega = (M^2 - m^2)/2M$, and the matrix element of the radiative decay is written as: $\mathcal{M} = ee_\mu J_\mu$, where e_μ is the photon polarization four-vector. So, the value of FF f_2 at point $q^2 = 0$ can be obtained from the following expression

$$f_2^2(0) = 12 \frac{\Gamma M^3}{\alpha(M^2 - m^2)^3}. \quad (9)$$

Another estimation of these FFs was obtained in Ref. [10] where the processes $e^+e^- \rightarrow 3\pi$, 4π were studied. The spin structure of the matrix element of the $a_1\rho\pi$ transition is similar to the $\gamma^*a_1\pi$ transition. The two coupling constants describing the $a_1\rho\pi$ transition were estimated by using the method of hard pions and knowing the width of the a_1 -meson (taken about 130 MeV) [10]. The corresponding values: $g^2/4\pi \approx 1.3$ and $\beta = 1.6$ were obtained. The following relations hold:

$$ef_1(q^2 = m_\rho^2) = \frac{g}{m_\rho}, \quad ef_2(q^2 = m_\rho^2) = \frac{g}{m_\rho}(\beta - 1).$$

The explicit expressions for the contributions to the hadronic tensor are:

- Unpolarized term $W_{\mu\nu}(0)$:

$$W_{\mu\nu}(0) = W_1(q^2)\tilde{g}_{\mu\nu} + \frac{W_2(q^2)}{M^2}\tilde{p}_{1\mu}\tilde{p}_{1\nu}, \quad \tilde{g}_{\mu\nu} = g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}, \quad \tilde{p}_{1\mu} = p_{1\mu} - \frac{p_1 q}{q^2}q_\mu,$$

where

$$W_1(q^2) = -|q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2)f_2|^2, \quad W_2(q^2) = q^2 (q^2 |f_1 + f_2|^2 - M^2 |f_2|^2), \quad (10)$$

- Term for vector polarization $W_{\mu\nu}(V)$:

$$\begin{aligned} W_{\mu\nu}(V) &= \frac{i}{M}V_1(q^2)\varepsilon_{\mu\nu\sigma\rho}s_\sigma q_\rho + \frac{i}{M^3}V_2(q^2)[\tilde{p}_{1\mu}\varepsilon_{\nu\alpha\sigma\rho}s_\alpha q_\sigma p_{1\rho} - \tilde{p}_{1\nu}\varepsilon_{\mu\alpha\sigma\rho}s_\alpha q_\sigma p_{1\rho}] + \\ &\quad + \frac{1}{M^3}V_3(q^2)[\tilde{p}_{1\mu}\varepsilon_{\nu\alpha\sigma\rho}s_\alpha q_\sigma p_{1\rho} + \tilde{p}_{1\nu}\varepsilon_{\mu\alpha\sigma\rho}s_\alpha q_\sigma p_{1\rho}], \\ V_1(q^2) &= -M^2(q^2 + M^2 - m^2)^{-1}|q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2)f_2|^2, \\ V_2(q^2) &= -M^2 q^2 (q^2 + M^2 - m^2)^{-1} \left[q^2 |f_1|^2 + \frac{1}{2}(q^2 - M^2 + m^2)|f_2|^2 + \right. \\ &\quad \left. \frac{1}{2}(3q^2 - M^2 + m^2)Re f_1 f_2^* \right], \\ V_3(q^2) &= -\frac{1}{2}M^2 q^2 Im f_1 f_2^*, \end{aligned} \quad (11)$$

- Term for tensor polarization $W_{\mu\nu}(T)$:

$$\begin{aligned} W_{\mu\nu}(T) &= T_1(q^2)\bar{Q}\tilde{g}_{\mu\nu} + T_2(q^2)\frac{\bar{Q}}{M^2}\tilde{p}_{1\mu}\tilde{p}_{1\nu} + T_3(q^2)(\tilde{p}_{1\mu}\tilde{Q}_\nu + \tilde{p}_{1\nu}\tilde{Q}_\mu) + \\ &\quad T_4(q^2)\tilde{Q}_{\mu\nu} + iT_5(q^2)(\tilde{p}_{1\mu}\tilde{Q}_\nu - \tilde{p}_{1\nu}\tilde{Q}_\mu), \end{aligned} \quad (12)$$

where

$$\begin{aligned} \tilde{Q}_\mu &= Q_{\mu\nu}q_\nu - \frac{q_\mu}{q^2}\bar{Q}, \quad \tilde{Q}_\mu q_\mu = 0, \quad \tilde{Q}_{\mu\nu} = Q_{\mu\nu} + \frac{q_\mu q_\nu}{q^4}\bar{Q} - \frac{q_\nu q_\alpha}{q^2}Q_{\mu\alpha} - \frac{q_\mu q_\alpha}{q^2}Q_{\nu\alpha}, \\ \tilde{Q}_{\mu\nu}q_\nu &= 0, \quad \bar{Q} = Q_{\alpha\beta}q_\alpha q_\beta, \\ T_1(q^2) &= 0, \quad T_2(q^2) = 3M^2 |f_2|^2, \quad T_3(q^2) = \frac{3}{2}(q^2 - M^2 + m^2)|f_2|^2 + 3q^2 Re f_1 f_2^*, \\ T_4(q^2) &= 3|q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2)f_2|^2, \quad T_5(q^2) = -3q^2 Im f_1 f_2^*. \end{aligned} \quad (13)$$

B. Expressions for the observables

Using the definitions of the cross section (2), of the leptonic (3) and hadronic (6) tensors, one can derive the expression for the unpolarized differential cross section in terms of the

structure functions $W_{1,2}$ (after averaging over the spins of the initial particles)

$$\frac{d\sigma^{un}}{d\Omega} = \frac{\alpha^2}{2q^4} \frac{p}{W} \left\{ -W_1(q^2) + \frac{1}{2}W_2(q^2) \left[\tau - 1 - \frac{(u-t)^2}{4M^2q^2} + \frac{M^2 - m^2}{4M^2q^2}(2q^2 + M^2 - m^2) \right] \right\}, \quad (14)$$

where $\tau = q^2/(4M^2)$, $t = (k_1 - p_1)^2$ and $u = (k_1 - p_2)^2$. In the reaction CMS this expression can be written as

$$\frac{d\sigma^{un}}{d\Omega} = \frac{\alpha^2}{2q^4} \frac{p}{W} (A + B \sin^2 \theta),$$

$$A = |q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2)f_2|^2, B = 2\tau p^2 [q^2 |f_1 + f_2|^2 - M^2 |f_2|^2], \quad (15)$$

where θ is the angle between the momenta of the axial-meson (\vec{p}) and of the electron beam (\vec{k}). Integrating this expression with respect to the axial-meson angular variables one obtains the following formula for the total cross section:

$$\sigma_{tot}(e^+ e^- \rightarrow \pi a_1) = \frac{2\pi\alpha^2}{3q^4} \frac{p}{W} \left[3|q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2)f_2|^2 + 4\tau p^2 [q^2 |f_1 + f_2|^2 - M^2 |f_2|^2] \right]. \quad (16)$$

Let us define an angular asymmetry, R , with respect to the differential cross section, $\sigma_{\pi/2}$, measured at $\theta = \pi/2$,

$$\frac{d\sigma^{un}}{d\Omega} = \sigma_{\pi/2} (1 + R \cos^2 \theta), R = -B/(A + B). \quad (17)$$

As it was previously shown in the case of $e^+ + e^- \rightarrow d + \bar{d}$ [17], this observable is very sensitive to the different underlying assumptions on the axial-meson FFs and does not require polarization measurements.

The differential cross section in terms of FFs $f_i(q^2)$ contains not only the moduli of these FFs but also their interference (15). One can choose a linear combinations of these FFs, in such a way that the unpolarized differential cross section will contain only moduli. Let us introduce new FFs $g_i(q^2)$ which related to the old ones as follows

$$f_1 = g_1 + g_2, \quad f_2 = c g_1 + d g_2, \quad (18)$$

where $c = -2q^2/(q^2 - M^2 + m^2)$, $d = (q^2 + M^2 - m^2)/(M^2 - q^2 + m^2)$. Then the structure functions W_i describing the unpolarized part of the hadronic tensor can be written as

$$W_1(q^2) = -\frac{4p^4 q^4}{(q^2 - M^2 - m^2)^2} |g_2|^2, \quad (19)$$

$$W_2(q^2) = 4p^2 q^4 \left[\frac{q^2}{(q^2 - M^2 + m^2)^2} |g_1|^2 - \frac{M^2}{(q^2 - M^2 - m^2)^2} |g_2|^2 \right]. \quad (20)$$

The cross section can be written, in the general case, as the sum of unpolarized and polarized terms, corresponding to the different polarization states and polarization directions of the incident and scattered particles:

$$\frac{d\sigma}{d\Omega} = \frac{d\sigma^{un}}{d\Omega} [1 + P_y + \lambda P_x + \lambda P_z + P_{zz} R_{zz} + P_{xz} R_{xz} + P_{xx} (R_{xx} - R_{yy}) + \lambda P_{yz} R_{yz}], \quad (21)$$

where P_i , P_{ij} , and R_{ij} , $i, j = x, y, z$ are, respectively, the components of the vector, tensor polarization and of the quadrupole polarization tensor of the outgoing a_1 -meson $Q_{\mu\nu}$, in its rest system and $d\sigma^{un}/d\Omega$ is the unpolarized differential cross section. λ is the degree of longitudinal polarization of the electron beam. It is explicitly indicated, in order to stress that these specific polarization observables are induced by the beam polarization.

Let us consider the different polarization observables and give their expression in terms of the $\gamma^* \rightarrow a_1\pi$ transition FFs.

- The vector polarization of the outgoing axial-meson, P_y , which does not require polarization in the initial state is

$$P_y = \frac{1}{8} \frac{\sqrt{7}}{\sigma_0} [(q^2 + M^2 - m^2)^2 - 4M^2 q^2] \sin(2\theta) \text{Im} f_1 f_2^*, \quad (22)$$

where $\sigma_0 = A + B \sin^2 \theta$. One can see that this polarization is determined by non-zero phase difference of the complex FFs f_1 and f_2 .

- The axial-vector meson can be tensor polarized also in case of unpolarized initial beams.

The components of the tensor polarization are

$$\begin{aligned} P_{xx} &= -\frac{3}{4} \frac{1}{\sigma_0} \sin^2 \theta \left| q^2 f_1 + \frac{1}{2} (q^2 - M^2 + m^2) f_2 \right|^2, \\ P_{xz} &= \frac{3}{4} \frac{\sqrt{7}}{\sigma_0} \frac{\sin(2\theta)}{q^2} \left\{ 2(q^2 + M^2 - m^2) \left| q^2 f_1 + \frac{1}{2} (q^2 - M^2 + m^2) f_2 \right|^2 + \right. \\ &\quad \left. + [(q^2 + M^2 - m^2)^2 - 4M^2 q^2] \left[\frac{1}{2} (q^2 - M^2 + m^2) |f_2|^2 + q^2 \text{Re} f_1 f_2^* \right] \right\}, \\ P_{zz} &= \frac{3}{8} \frac{1}{\sigma_0} \frac{1}{M^2 q^2} [(q^2 + M^2 - m^2)^2 - 4M^2 q^2] \left\{ \left| q^2 f_1 + \frac{1}{2} (q^2 - M^2 + m^2) f_2 \right|^2 + \right. \\ &\quad \left. + \frac{1}{2} \sin^2 \theta \left[-q^4 |f_1|^2 + q^2 (q^2 + 3M^2 - 3m^2) \text{Re} f_1 f_2^* + \right. \right. \\ &\quad \left. \left. + \left(2q^2 (q^2 - M^2) - \frac{3}{4} (q^2 - M^2 + m^2)^2 \right) |f_2|^2 \right] \right\}. \end{aligned} \quad (23)$$

A possible non-zero phase difference between the axial-meson FFs leads to another T-odd polarization observable proportional to the R_{yz} component of the axial-meson tensor polarization. It takes the form

$$P_{yz} = -\frac{3}{2} \frac{\sqrt{7}}{\sigma_0} \left[(q^2 + M^2 - m^2)^2 - 4M^2 q^2 \right] \sin \theta \operatorname{Im} f_1 f_2^*. \quad (24)$$

- Let us consider now the case of a longitudinally polarized electron beam. The other two components of the axial-meson vector polarization (P_x , P_z) require the initial particle polarization and are

$$\begin{aligned} P_x &= -\frac{1}{4} \frac{\sqrt{7}}{\sigma_0} \sin \theta \left\{ 2q^2 (q^2 + M^2 - m^2) |f_1|^2 + [(q^2 - M^2)^2 - m^4] |f_2|^2 + \right. \\ &\quad \left. + [q^2 (3q^2 - 2M^2 - 2m^2) - (M^2 - m^2)^2] \operatorname{Re} f_1 f_2^* \right\}, \\ P_z &= \frac{1}{2} \frac{1}{\sigma_0} \cos \theta \left| q^2 f_1 + \frac{1}{2} (q^2 - M^2 + m^2) f_2 \right|^2. \end{aligned} \quad (25)$$

III. SPIN-DENSITY MATRIX OF AXIAL-MESON

For unstable particles, the vector and tensor polarizations are directly related to the angular distribution of their decay products; one can show that the angular distribution can be expressed in terms of the spin-density matrix. Let us calculate the elements of the spin-density matrix of the axial-meson which is produced in the reaction $e^+ + e^- \rightarrow \pi + a_1$. The calculation is done in CMS of this reaction.

In case of unpolarized initial lepton beams, the convolution of the lepton $L_{\mu\nu}$ and hadron $W_{\mu\nu}$ tensors can be written as

$$S^{un} = S_{\mu\nu} U_\mu U_\nu^*, \quad (26)$$

where U_μ is the polarization four-vector of the detected axial-meson and the $S_{\mu\nu}$ tensor can be represented in the following general form

$$S_{\mu\nu} = S_1 g_{\mu\nu} + S_2 q_\mu q_\nu + S_3 k_{1\mu} k_{1\nu} + S_4 (k_{1\mu} q_\nu + q_\mu k_{1\nu}) + i S_5 (k_{1\mu} q_\nu - q_\mu k_{1\nu}). \quad (27)$$

The functions S_i ($i = 1 - 5$) can be written in terms of the two transition FFs of the axial-meson. Their explicit form is

$$S_1 = -q^2 \left| q^2 f_1 + \frac{1}{2} (q^2 - M^2 + m^2) f_2 \right|^2,$$

$$\begin{aligned}
S_2 &= \left[q^2(q^2 - M^2) - \frac{1}{4}(q^2 - M^2 + m^2 + 2Wp \cos \theta)^2 \right] |f_2|^2 + \\
&\quad + q^2(q^2 + M^2 - m^2 - 2Wp \cos \theta) \text{Re} f_1 f_2^*, \\
S_3 &= -4 \left| q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2) f_2 \right|^2, \\
S_4 &= 2q^4 |f_1|^2 + 2q^2(q^2 - M^2 + m^2 + Wp \cos \theta)^2 \text{Re} f_1 f_2^* + \\
&\quad + \frac{1}{2}(q^2 - M^2 + m^2)(q^2 - M^2 + m^2 + 2Wp \cos \theta) |f_2|^2, \\
S_5 &= -2q^2 Wp \cos \theta \text{Im} f_1 f_2^*. \tag{28}
\end{aligned}$$

The T-odd structure function S_5 is not zero here since the transition FFs of the axial-meson are complex functions.

The elements of the spin-density matrix of the axial-meson are defined as

$$S\rho_{mm'} = S_{\mu\nu} U_\mu^{(m)} U_\nu^{(m')*}, \quad S = S_{\mu\nu} \left(-g_{\mu\nu} + \frac{p_{1\mu} p_{1\nu}}{M^2} \right), \tag{29}$$

where $S = 2q^2(A + B \sin^2 \theta)$, and $U_\mu^{(m)}$ is the polarization four-vector of the axial-meson with definite ($m = 0, \pm 1$) projection on the z axis. In our case it is directed along the axial-meson momentum and thus $U_\mu^{(m)}$ are the polarization vectors with definite helicity.

The elements of the spin-density matrix of the axial-meson are

$$\begin{aligned}
\rho_{++} &= \rho_{--} = \frac{q^2}{2S} (1 + \cos^2 \theta) \left| q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2) f_2 \right|^2, \\
\rho_{00} &= \frac{q^4}{M^2 S} \sin^2 \theta \left\{ \frac{1}{4}(q^2 + M^2 - m^2)^2 |f_1|^2 + [2p^2 q^2 + M^2(q^2 - M^2 + m^2)] \text{Re} f_1 f_2^* + \right. \\
&\quad \left. + (m^2 M^2 + p^2 q^2) |f_2|^2 \right\}, \\
\rho_{+-} &= \rho_{-+} = \frac{q^2}{2S} \sin^2 \theta \left| q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2) f_2 \right|^2, \\
\rho_{+0} &= -\frac{q^4}{S} \sqrt{\frac{\tau}{2}} \sin \theta \cos \theta \left\{ (q^2 + M^2 - m^2) |f_1|^2 + \left(1 - \frac{M^2 - m^2}{q^2} \right) [(q^2 + M^2 - m^2) \text{Re} f_1 f_2^* + \right. \\
&\quad \left. + \frac{1}{2}(q^2 - M^2 - m^2) |f_2|^2] + 2p^2 f_2 f_1^* \right\}, \\
\rho_{-0} &= -\rho_{+0}, \quad \rho_{0+} = \rho_{+0}^*, \quad \rho_{0-} = \rho_{-0}^*. \tag{30}
\end{aligned}$$

The spin-density matrix is normalized as $\text{Tr} \rho = 1$ or $\rho_{++} + \rho_{--} + \rho_{00} = 1$. The element ρ_{+0} is complex and the real and imaginary parts are written as:

$$\begin{aligned}
\text{Re} \rho_{+0} &= -\frac{q^4}{S} \sqrt{\frac{\tau}{2}} \sin \theta \cos \theta \left\{ (q^2 + M^2 - m^2) |f_1|^2 + \frac{1}{2} \left(1 - \frac{M^2 - m^2}{q^2} \right) \right. \\
&\quad \left. (q^2 - M^2 - m^2) |f_2|^2 + \frac{1}{2} \left[3q^2 - 2M^2 - 2m^2 - \frac{(M^2 - m^2)^2}{q^2} \right] \text{Re} f_1 f_2^* \right\}, \\
\text{Im} \rho_{+0} &= \frac{p^2 q^4}{S} \sqrt{\frac{\tau}{2}} \sin 2\theta \text{Im} f_1 f_2^*. \tag{31}
\end{aligned}$$

Let us consider the case when the electron beam is longitudinally polarized. Then the convolution of the spin-dependent part of the lepton and hadron tensors can be written as

$$S(\lambda) = S_{\mu\nu}(\lambda)U_\mu U_\nu^*, \quad (32)$$

where λ is the degree of the electron beam polarization and the $S_{\mu\nu}(\lambda)$ tensor can be written as

$$S_{\mu\nu}(\lambda) = Q_1 \epsilon_{\mu\nu\alpha\beta} k_{1\alpha} k_{2\beta} + Q_2 (q_\mu a_\nu - q_\nu a_\mu) + Q_3 (q_\mu a_\nu + q_\nu a_\mu), \quad (33)$$

where $a_\mu = \epsilon_{\mu\alpha\beta\gamma} p_\alpha k_{1\beta} k_{2\gamma}$, $p = p_1 - p_2$. The structure functions Q_i ($i = 1 - 3$) can be written in terms of the two transition FFs of the axial-meson as

$$\begin{aligned} Q_1 &= -2i\lambda \left| q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2) f_2 \right|^2, \\ Q_2 &= -2i\lambda \text{Re} \left[q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2) f_2 \right] f_2^*, \\ Q_3 &= 2\lambda q^2 \text{Im} f_1 f_2^*. \end{aligned} \quad (34)$$

The T-odd structure function Q_3 is not zero since FFs are complex functions in the TL region.

The elements of the spin-density matrix of the a_1 -meson that depend on the longitudinal polarization of the electron beam can be defined as

$$S\rho_{mm'}(\lambda) = S_{\mu\nu}(\lambda)U_\mu^{(m)}U_\nu^{(m')*}, \quad (35)$$

and they are expressed in terms of FFs as

$$\begin{aligned} \rho_{++}(\lambda) &= -\rho_{--}(\lambda) = \frac{\lambda}{S} q^2 \cos \theta \left| q^2 f_1 + \frac{1}{2}(q^2 - M^2 + m^2) f_2 \right|^2, \\ \rho_{00}(\lambda) &= \rho_{+-}(\lambda) = \rho_{-+}(\lambda) = 0, \\ \rho_{+0}(\lambda) &= -\frac{\lambda}{S} \sqrt{\frac{\tau}{2}} q^4 \sin \theta \left\{ (q^2 + M^2 - m^2) |f_1|^2 + \right. \\ &\quad \left. + \left(1 - \frac{M^2 - m^2}{q^2} \right) \left[(q^2 + M^2 - m^2) \text{Re} f_1 f_2^* + \frac{1}{2}(q^2 - M^2 - m^2) |f_2|^2 \right] + 2p^2 f_2 f_1^* \right\}, \\ \rho_{0+}(\lambda) &= \rho_{+0}^*(\lambda), \quad \rho_{-0}(\lambda) = \rho_{+0}(\lambda), \quad \rho_{0-}(\lambda) = \rho_{-0}^*(\lambda). \end{aligned} \quad (36)$$

The spin-density matrix element $\rho_{+0}(\lambda)$ is a complex quantity and its real and imaginary parts are

$$\begin{aligned} \text{Re} \rho_{+0}(\lambda) &= -\frac{\lambda}{S} \sqrt{\frac{\tau}{2}} q^4 \sin \theta \left\{ (q^2 + M^2 - m^2) |f_1|^2 + \frac{1}{2} \left(1 - \frac{M^2 - m^2}{q^2} \right) (q^2 - M^2 - m^2) |f_2|^2 + \right. \\ &\quad \left. + \frac{1}{2} \left[(3q^2 - 2M^2 - 2m^2 - \frac{(M^2 - m^2)^2}{q^2}) \text{Re} f_1 f_2^* \right] \right\}, \\ \text{Im} \rho_{+0}(\lambda) &= \frac{\lambda}{S} \sqrt{2\tau} p^2 q^4 \sin \theta \text{Im} f_1 f_2^*. \end{aligned} \quad (37)$$

The axial-meson FFs are complex functions in the TL region. So, for the complete determination of FFs it is necessary to measure three quantities: two moduli of FFs and their phase difference. Therefore, the measurement of the unpolarized differential cross section does not allow to determine completely FFs. It is necessary to determine the spin-density matrix elements of the produced axial meson measuring the angular distribution of its decay products.

The measurement of the angular dependence of the unpolarized differential cross section allows to determine the structure functions A and B . So, one can determine the following ratio

$$R_1 = \frac{B}{A} = \frac{p^2}{2M^2} \left[1 + 2r \cos \alpha + \left(1 - \frac{M^2}{q^2} \right) r^2 \right] \left[1 + \left(1 + \frac{m^2 - M^2}{q^2} \right) r \cos \alpha + \frac{1}{4} \left(1 + \frac{m^2 - M^2}{q^2} \right)^2 r^2 \right]^{-1}, \quad (38)$$

where $r = |f_2|/|f_1|$ and α is the relative phase of two complex FFs f_1 and f_2 defined as follows: $\alpha = \alpha_1 - \alpha_2$, where $\alpha_1 = \text{Arg}f_1$ and $\alpha_2 = \text{Arg}f_2$. Thus, Eq. 38 contains two unknown quantities: α and r . Another equation for the determination of these quantities can be obtained, for example, from the following ratio of the spin-density matrix elements of the produced axial meson

$$R_2 = \frac{\rho_{00}}{\rho_{+-}} = 8\tau \left\{ \frac{1}{4}(q^2 + M^2 - m^2)^2 + [2p^2q^2 + M^2(q^2 - M^2 + m^2)] r \cos \alpha + (m^2M^2 + p^2q^2)r^2 \right\} \left[q^4 + q^2(q^2 - M^2 + m^2)r \cos \alpha + \frac{1}{4}(q^2 - M^2 + m^2)^2 r^2 \right]^{-1}. \quad (39)$$

And, finally, the phase difference α can be determined by fixing the sign of $\sin \alpha$ from the measured spin-density matrix element $\text{Im}\rho_{0+}$. Note that the complete determination of two complex FFs f_1 and f_2 does not require the polarization of the initial beams. The measurement of the angular distribution of the unpolarized differential cross section allows to determine the structure functions A and B separately. Knowing the ratio r , one can determine the moduli $|f_1|$ and $|f_2|$.

IV. NUMERICAL RESULTS

VMD inspired models have proved to be very successful in describing the structure of hadrons. Such models contain a small number of parameters, with transparent physical

meaning, and can be analytically extended to the full region of momentum transfer squared. A monopole-like behavior reproduce quite well the existing experimental data on pion FFs, and satisfies pQCD asymptotic [24]. In order to predict the behavior of polarization observables, we suggest a simple model for the a_1 transition FFs, in TL region.

We used a simple VMD-based parametrization saturated by vector mesons. The contribution of one vector meson is given by the Breit-Wigner form

$$\Delta f_i = \frac{C_{v,i} M_v}{M_v^2 - q^2 + i M_v \Gamma_v}, \quad i = 1, 2, \quad (40)$$

where M_v and Γ_v are the mass and the width of a vector meson carrying the interaction. In general one should introduce all allowed vector mesons, but, as shown in Refs. [11, 25], the largest contribution to the cross section is given by the $\rho(770)$, $\rho'(1450)$, and, at higher energies, $\rho''(1700)$. We consider only data for the total cross section at energies above the $a_1\pi$ kinematical threshold, which have been compiled from Refs. [5, 26–28]. The experimental data [5] show a clear contribution from J/ψ around $\sqrt{s} = 3.01$ GeV. We excluded the corresponding (four) data points from the fit. The final form of our parametrization of the two $\gamma^*\pi a_1$ transition form factors

$$f_i = \frac{C_{\rho,i} M_\rho}{M_\rho^2 - q^2 + i M_\rho \Gamma_\rho} + \frac{C_{\rho',i} M_{\rho'}}{M_{\rho'}^2 - q^2 + i M_{\rho'} \Gamma_{\rho'}} + \frac{C_{\rho'',i} M_{\rho''}}{M_{\rho''}^2 - q^2 + i M_{\rho''} \Gamma_{\rho''}}, \quad i = 1, 2. \quad (41)$$

Such parametrization has, therefore, in total six parameters: three normalization constants, $C_{\rho,i}$, $C_{\rho',i}$, $C_{\rho'',i}$ for each FF.

The result of the fit for the available data on the total cross section of the $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$ process from [5, 26–28] is shown in Fig. 1, and corresponds to $\chi^2/ndf = 455/154 = 2.9$. The different sets of data points are quite dispersed, especially in the region of the maximum. The resulting normalization constants are given in Table I. The $\gamma^*\pi a_1$ transition form factors (moduli) are presented in the Fig.2. Peaks can be seen in correspondence with the masses of the chosen vector mesons. For $|f_2|$, one can see a bump around $q^2 = 4$ GeV², which results from the interference of the different terms.

Figs. 3, 4 show the vector and tensor polarization observables. One should note here that the vector polarization of vector mesons can not be measured through their decays which are driven by strong and electromagnetic interaction with conservation of P-parity [29]. Fig. 5 shows the predictions for the density matrix elements. These quantities can be quite large and show a particular behavior, which can be experimentally verified. Finally, Fig. 6 shows the predictions for the ratios R_1 , R_2 Eqs. (38,39).

i	$C_{\rho,i}$	$C_{\rho',i}$	$C_{\rho'',i}$
1	2.35 ± 0.01	-0.089 ± 0.008	-0.131 ± 0.009
2	-4.45 ± 0.11	0.56 ± 0.05	0.38 ± 0.03

TABLE I: Optimal values of the normalization constants $C_{\rho,i}$, $C_{\rho',i}$, $C_{\rho'',i}$ obtained by the fitting procedure.




FIG. 1: Total cross section data for the annihilation $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$ from [5, 26–28]. The line represents the fit of the total cross section (16) for the reaction $e^+ + e^- \rightarrow a_1 + \pi$ with VMD-based model of transition form factors (41).




FIG. 2: q^2 dependence of the $\gamma^*\pi a_1$ transition form factors f_1 , f_2 (41).




FIG. 3: Angular dependence of the vector polarization observables P_x , P_y , P_z at $q^2 = 2 \text{ GeV}^2$, Eqs. (22,25).




FIG. 4: Angular dependence of the tensor polarization observables P_{xx} , P_{xz} , P_{zz} , P_{yz} at $q^2 = 2 \text{ GeV}^2$, Eqs. (23,24).

V. CONCLUSION

Using the parametrization of the electromagnetic transition $\gamma^* \rightarrow a_1(1260)\pi$ in terms of two complex FFs, we investigated the polarization phenomena in the annihilation reaction




FIG. 5: Angular dependence of the elements of the spin density matrix $\rho_{++}, \rho_{00}, \rho_{+-}, \rho_{+0}$ at $q^2 = 2$ GeV^2 , in case of unpolarized collision (solid lines) from Eqs. (30,31). In case of longitudinally polarized electrons the corresponding elements are shown also (dashed lines, (d),(e)) from Eqs. (37) and (f) from Eq. (36).




FIG. 6: q^2 behavior of the ratios R_1, R_2 , Eqs. (38,39).

(1). We calculated the differential (and total) cross section and various polarization observables as functions of the FFs. The spin-density matrix elements of the produced axial meson have been also calculated.

The q^2 dependence of the FFs shows, as expected, peaks in correspondence of the masses of the considered vector meson. An interesting feature appears for f_2 at $Q^2 \sim 4 \text{ GeV}^2$, which is due to an interference of the different terms. Note that none of the considered vector mesons corresponds to a mass of such value. An interference of such origin can be the reason for the plateau seen in the cross section data, which, however does not appear with large intensity in our fit.

The reaction (1) has been clearly detected in the experiments. The present results can be useful for the analysis of the experimental data and for the determination of the $\gamma^* \rightarrow a_1(1260)\pi$ transition FFs.

VI. ACKNOWLEDGMENT

This work was inspired by enlightening discussions with Prof. M. P. Rekaló. Two of us (G.I.G. and C.A.) acknowledge the hospitality of CEA, Saclay. This work was partly supported by the Slovak Grant Agency for Sciences VEGA under Grant N. 2/4099/26 (CA) and by grant INTAS Ref. Nr 05-1000008-8328 (G.I.G.). We acknowledge the french Groupement de Recherche Nucleon, for useful meetings and continuous support.

-
- [1] R. R. Akhmetshin *et al.* [CMD2 Collaboration], Phys. Lett. B **466**, 392 (1999).
 - [2] B. Bartoli *et al.*, Nuovo Cim. **A70**, 615 (1970).
 - [3] L. M. Kurdadze *et al.*, Phys. Lett. **B42**, 515 (1972).
 - [4] R. R. Akhmetshin *et al.* [CMD-2 Collaboration], Phys. Lett. B **595**, 101 (2004).
 - [5] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **71**, 052001 (2005).
 - [6] A. Denig [BaBar Collaboration], arXiv:hep-ex/0611029.
 - [7] R. Baldini, S. Dubnicka, P. Gauzzi, S. Pacetti, E. Pasqualucci and Y. Srivastava, Eur. Phys. J. C **11**, 709 (1999); Nucl. Phys. **A666**, 38 (2000); J. Volmer *et al.*, Phys. Rev. Lett. **86**, 1713 (2001).

- [8] M. T. Vaughn, P. J. Polito, *Lett. Nuovo Cim.* **1**, 74 (1971).
- [9] G. Kramer and J. L. Uretsky and T. F. Walsh, *Phys. Rev.* **D3**, 719 (1991).
- [10] A. M. Altukhov and I. B. Khriplovitch, *Sov. J. Nucl. Phys.* **14**, 440 (1972).
- [11] P. Lichard and J. Juran, *Phys. Rev. D* **76**, 094030 (2007).
- [12] S. I. Eidelman, Z. K. Silagadze and E. A. Kuraev, *Phys. Lett. B* **346**, 186 (1995).
- [13] R. S. Plant and M. C. Birse, *Phys. Lett. B* **365**, 292 (1996).
- [14] N. N. Achasov and A. A. Kozhevnikov, *Phys. Rev. D* **71**, 034015 (2005).
- [15] A. E. Bondar, S. I. Eidelman, A. I. Milstein, N. I. Root, *Phys. Lett.* **B466**, 403 (1999).
- [16] B. A. Li, *Phys. Rev. D* **58**, 097302 (1998).
- [17] G. I. Gakh, E. Tomasi-Gustafsson, C. Adamuscin, S. Dubnicka, and A. Z. Dubnickova, *Phys. Phys.* **C74**, 025202 (2006).
- [18] C. Adamuscin, G. I. Gakh and E. Tomasi-Gustafsson, *Phys. Rev. C* **75**, 065202 (2007).
- [19] J. Gunion and L. Stodolsky, *Phys. Rev. Lett.* **30**, 345 (1973); V. Franco, *Phys. Rev. D* **8**, 826 (1973); V. N. Boitsov, L. A. Kondratyuk and V. B. Kopeliovich, *Sov. J. Nucl. Phys.* **16**, 287 (1973) [*Yad. Fiz.* **16**, 515 (1972)]. F. M. Lev, *Sov. J. Nucl. Phys.* **21**, 45 (1973);
- [20] M. P. Rekalov, E. Tomasi-Gustafsson and D. Prout, *Phys. Rev. C* **60**, 042202R (1999).
- [21] G. I. Gakh and E. Tomasi-Gustafsson, *Nucl. Phys. A* **799**, 127 (2008).
- [22] E. Tomasi-Gustafsson and G. I. Gakh, *Phys. Rev. C* **72**, 015209 (2005).
- [23] E. Tomasi-Gustafsson, E. A. Kuraev, S. Bakmaev and S. Pacetti, *Phys. Lett. B* **659**, 197 (2008).
- [24] S. J. Brodsky, J. R. Hiller, *Phys. Rev.* **D46**, 2141 (1992).
- [25] R. R. Akhmetshin *et al.* [CMD-2 Collaboration], *Phys. Lett. B* **475**, 190 (2000).
- [26] C. Bacci *et al.*, *Phys. Lett. B* **95**, 139 (1980).
- [27] B. Esposito *et al.*, *Lett. Nuovo Cim.* **28**, 195 (1980).
- [28] A. Cordier, D. Bisello, J. C. Bizot, J. Buon, B. Delcourt, L. Fayard and F. Mane, *Phys. Lett. B* **109**, 129 (1982).
- [29] E. Tomasi-Gustafsson and M. P. Rekalov, *Phys. Part. Nucl.* **33**, 220 (2002) [*Fiz. Elem. Chast. Atom. Yadra* **33**, 436 (2002)].