

The $g_{V\sigma\gamma}$ -coupling constants in hadron electrodynamics

Michail P. Rekalo ¹

Middle East Technical University, Physics Department, Ankara 06531, Turkey

Egle Tomasi-Gustafsson ²

DAPNIA/SPhN, CEA/Saclay, 91191 Gif-sur-Yvette Cedex, France

Abstract

Recent measurements of the branching ratios for the decays $\omega \rightarrow \pi^0\pi^0\gamma$ and $\rho^0 \rightarrow \pi^0\pi^0\gamma$ lead to coupling constants of the $V\sigma\gamma$ -interaction (V is a vector meson) one order of magnitude smaller than previously assumed to describe the threshold cross section of $\gamma + p \rightarrow p + \rho^0$ and the HERMES effect. The new $g_{V\sigma\gamma}^2$ couplings are in contradiction with the predictions of the QCD sum rules, but are in good agreement with VDM-estimation of the $\sigma \rightarrow \gamma\gamma$ -width.

1 Introduction

The coupling constants $g_{V\sigma\gamma}$, $V = \rho$ or ω are important ingredients for the theoretical analysis of many different hadronic electromagnetic processes. If $m_\sigma < m_V$, the decays $V \rightarrow \sigma + \gamma$ can occur directly-with radiation of electric dipole photons. In this respect, these decays are essentially different from the decays $V \rightarrow P + \gamma$, $P = \pi, \eta, \eta'$ -with radiation of magnetic dipole photons. In the quark models, these last decays are induced by the quark magnetic moment, with transition $S = 1 \rightarrow S = 0$, where S is the total spin of the $q\bar{q}$ -system (in the corresponding meson), whereas the decays $V \rightarrow \sigma + \gamma$ are induced by the internal motion of quarks, with transition $S = 1 \rightarrow S = 1$ [1].

Let us mention the main applications of the $g_{V\sigma\gamma}$ -coupling constants:

¹ Permanent address: *National Science Center KFTI, 310108 Kharkov, Ukraine*

² Corresponding author: etomasi@cea.fr

- The σ -exchange for the vector meson production [2–6] $\gamma + p \rightarrow p + \rho^0$, near threshold.
- The electromagnetic transition $\gamma^* \rightarrow \sigma + \omega$, in the space-like region of the virtual photon γ^* four-momenta, enters in the calculation of meson exchange currents, in particular in the analysis of the deuteron electromagnetic form factors at large momentum transfer [7–9].
- The branching ratio of rare radiative decays of vector mesons as $\rho^0 \rightarrow \pi^0 \pi^0 \gamma$ and $\omega \rightarrow \pi^0 \pi^0 \gamma$ are controlled by the $g_{V\sigma\gamma}$ -coupling constants [10,11].
- The HERMES effect [12], concerning the inclusive electroproduction cross section on light nuclei can be explained under specific assumptions [13] about the absolute value of the $g_{\rho\sigma\gamma}$ and $g_{\omega\sigma\gamma}$ coupling constants and the corresponding electromagnetic form factors.
- The coupling constant of the $\sigma \rightarrow \gamma\gamma$ -vertex, which can be calculated on the basis of the $g_{V\sigma\gamma}$ couplings, is important for the analysis of real and virtual Compton scattering on nucleons [14–16].
- The exact value of the $g_{V\sigma\gamma}$ coupling constant, in principle, allows one to constrain the $g_{\sigma NN}$ -coupling, which describes the scalar exchange in the NN -potential [17].
- Finally, the $g_{V\sigma\gamma}$ coupling constants have an implicit theoretical interest, being fundamental coupling constants of hadron electrodynamics.

Attempts to estimate these coupling constants from QCD -sum rules, give relatively large absolute values [18]. Phenomenological methods based on the existing experimental data about different electromagnetic processes need serious additional assumptions. For example, in the framework of σ -exchange for $\gamma + p \rightarrow p + \rho^0$, it is possible to estimate the product $g_{\rho\sigma\gamma}g_{\sigma NN}$, assuming a definite form of the phenomenological form factors, which insure the correct behavior of the differential cross section $d\sigma/dt$, in the near-threshold region, and depend on ad-hoc cut-off parameters. Taking $g_{\sigma NN}^2/4\pi = 8$, one obtains $g_{\rho\sigma\gamma} = 2.71$ [2]. Considering only σ -exchange in $\gamma + p \rightarrow p + \rho^0$, the sign of this constant can not be determined, but some polarization observables are sensitive to this sign [19]. Other possible contributions to the matrix element for the process $\gamma + p \rightarrow p + \rho^0$, such as N^* -excitation, N -exchange in s - and u -channel are neglected in such oversimplified consideration [2].

Also the estimation of the discussed coupling constants done in framework of the HERMES effect [13] can not be considered as direct and model independent.

Another source of information about the $\rho\sigma\gamma$ -vertex is the radiative decay $\rho^0 \rightarrow \pi^+\pi^-\gamma$ -with relatively large branching ratio [20]. Namely this decay has been considered to favor large values of $g_{\rho\sigma\gamma}$ in comparison with $g_{\omega\sigma\gamma}$, because of the corresponding widths $\Gamma(\omega \rightarrow \pi^+\pi^-\gamma) \ll \Gamma(\rho \rightarrow \pi^+\pi^-\gamma)$ [2]. However it is necessary to point out that for the decay $\rho^0 \rightarrow \pi^+\pi^-\gamma$ the bremsstrahlung mechanism dominates, therefore the σ -contribution, being of the order of the

error bars, can not give the $g_{\rho\sigma\gamma}$ with good accuracy. Note also that the precise determination of $g_{\rho\sigma\gamma}$ depends essentially on the mass and the width of the σ -meson. In [11] different values of $g_{\rho\sigma\gamma}$ could be found, with the general conclusion that the $g_{\rho\sigma\gamma}$ value from $\rho^0 \rightarrow \pi^+\pi^-\gamma$ is not in contradiction with the estimation done in [2].

Finally, having a definite value of $g_{\rho\sigma\gamma}$, it is possible to estimate the branching ratio for $\rho \rightarrow \pi^0\pi^0\gamma$, in framework of the effective Lagrangian approach. In this case, the absence of the bremsstrahlung mechanism results in the fact that the σ -exchange mechanism is important. But the predicted $BR(\rho^0 \rightarrow \pi^0\pi^0\gamma) \simeq (45 - 200) \cdot 10^{-5}$, with $g_{\rho\sigma\gamma} = 3 \div 5$, were too large in comparison with other theoretical expectations for this decay [21].

Moreover, and this will be the important background of this paper, large values of $g_{\rho\sigma\gamma}$ contradict essentially the recent experimental results from Novosibirsk, concerning the direct measurement of the decay $\rho^0 \rightarrow \pi^0\pi^0\gamma$ [22]:

$$Br(\rho \rightarrow \pi^0\pi^0\gamma) = (4.1_{-0.9}^{+1.0} \pm 0.3) \cdot 10^{-5},$$

$$Br(\rho \rightarrow \sigma\gamma \rightarrow \pi^0\pi^0\gamma) = (1.9_{-0.8}^{+0.9} \pm 0.4) \cdot 10^{-5}.$$

The main aim of this paper is to estimate the coupling constant $g_{\rho\sigma\gamma}$ on the basis of these new important experimental data, which is, in our opinion, the most straightforward way.

2 The σ -contribution to the $\rho \rightarrow \pi^0 + \pi^0 + \gamma$ -decay

A crude estimation of the $g_{\rho\sigma\gamma}$ coupling constant can be easily obtained from the experimental data about $\rho^0 \rightarrow \sigma\gamma \rightarrow \pi^0\pi^0\gamma$ [22], under the assumption that the σ -mass is smaller than the ρ -meson mass, so that the decay $\rho \rightarrow \sigma + \gamma$ is allowed.

Neglecting for a moment the σ -width (which is an evident oversimplification of reality, as $\Gamma_\sigma = (0.6 \div 1)$ GeV [20]), we can find:

$$\Gamma(\rho^0 \rightarrow \sigma\gamma \rightarrow \pi^0\pi^0\gamma) = \frac{1}{3}\Gamma(\rho^0 \rightarrow \sigma\gamma), \quad (1)$$

taking into account the identity of the neutral π^0 -mesons, produced in $\rho^0 \rightarrow \pi^0\pi^0\gamma$, and the isotopic relation: $g(\sigma \rightarrow \pi^0\pi^0) = g(\sigma \rightarrow \pi^+\pi^-)$.

The matrix element of the decay $V \rightarrow \sigma + \gamma$ can be written in the following

form:

$$\mathcal{M} = \frac{e g_{V\sigma\gamma}}{M} (\epsilon^* \cdot U \ k \cdot p - \epsilon^* \cdot p \ U \cdot k), \quad (2)$$

where $\epsilon_\mu(k)$ and $U_\mu(p)$ are the four-vectors describing the vector polarization (four-momenta) of the photon and of the V -meson.

Averaging over the V -meson polarizations and summing over the photon polarizations, one can find for the width of the decay $V \rightarrow \sigma + \gamma$ (still considering the σ -meson as a stable particle):

$$\Gamma(V \rightarrow \sigma\gamma) = \alpha \frac{g_{V\sigma\gamma}^2}{24} M \left(1 - \frac{m_\sigma^2}{M^2}\right)^3 \simeq 231 g_{V\sigma\gamma}^2 \left(1 - \frac{m_\sigma^2}{M^2}\right)^3 \text{ keV}, \quad (3)$$

where $M(m_\sigma)$ is the mass of the $V(\sigma)$ -meson. The numerical estimation (3) is done for $M=0.77$ GeV (ρ -meson). Evidently, at a given value of the $g_{V\sigma\gamma}$ -coupling constant, the value of $\Gamma(V \rightarrow \sigma\gamma)$ depends essentially on the value taken for the σ -meson mass, which can be in a wide range: $m_\sigma = (0.4 \div 1.2)$ GeV [20]. In Table 1 we report the values of $\Gamma(\rho^0 \rightarrow \sigma\gamma)$ and $\Gamma(\rho^0 \rightarrow \sigma\gamma \rightarrow \pi^0\pi^0\gamma)$, calculated for the values of the $g_{\rho\sigma\gamma}$ -constant and m_σ from Refs. [2] and [13]. Such estimations for $\Gamma(\rho^0 \rightarrow \sigma\gamma \rightarrow \pi^0\pi^0\gamma)$ are almost two orders of magnitude larger than the value given by the experiment [22]:

$$\Gamma(\rho^0 \rightarrow \sigma\gamma \rightarrow \pi^0\pi^0\gamma) = (2.85_{-1.2}^{+1.4} \pm 0.6) \text{ keV}. \quad (4)$$

Note that the predictions of QCD sum rules [18] are also in serious disagreement with the direct experimental estimation (4).

m_σ [GeV]	$g_{\rho\sigma\gamma}$	$\Gamma(\rho^0 \rightarrow \sigma\gamma)$ [keV]	$\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)$ [keV]	Ref.
0.5	3	379	126	[2]
0.6	5-6	308-444	103-148	[13]

Table 1

Radiative widths for different masses of σ -meson, m_σ and coupling constant $g_{\rho\sigma\gamma}$.

This crude preliminary estimation leads to a conclusion which is inconsistent with the suggested interpretation of the HERMES effect [13] and the description of threshold behavior of the cross section for $\gamma + p \rightarrow p + \rho^0$ [2]. Therefore let us estimate more precisely the $g_{\rho\sigma\gamma}$ coupling constant on the basis of the decay $\rho^0 \rightarrow \sigma\gamma \rightarrow \pi^0\pi^0\gamma$, removing the hypothesis taken above, i.e. taking

into account the finite value of the σ -width, and the possibility that in some cases it is possible to have $m_\sigma > M$.

Considering the sequence of decays $\rho^0 \rightarrow \sigma + \gamma \rightarrow \pi^0 + \pi^0 + \gamma$, one can find, for the effective $\pi^0 + \pi^0$ -mass distribution, the following formula (see Appendix):

$$\frac{d\Gamma}{dw^2} = \frac{e^2 g_{\rho\sigma\gamma}^2}{288\pi^2} M \left(1 - \frac{w^2}{M^2}\right)^3 \frac{\beta_w}{\beta_\sigma} \frac{m_\sigma \Gamma_\sigma}{(w^2 - m_\sigma^2)^2 + \Gamma_\sigma^2 m_\sigma^2}, \quad (5)$$

where Γ_σ is the total width for the σ -meson, and:

$$\beta_\sigma = \sqrt{1 - \frac{4m_\pi^2}{m_\sigma^2}}, \quad \beta_w = \sqrt{1 - \frac{4m_\pi^2}{w^2}}.$$

For the estimation of the coupling constant $g_{\rho\sigma\gamma}^2$ on the basis of $\Gamma(\rho^0 \rightarrow \sigma\gamma \rightarrow \pi^0\pi^0\gamma)$ the following formula can be used:

$$g_{\rho\sigma\gamma}^2 = \frac{\Gamma(\rho^0 \rightarrow \sigma\gamma \rightarrow \pi^0\pi^0\gamma) 72\pi}{M\alpha} \frac{1}{r_1 r_2 f(r_1, r_2)} \quad (6)$$

with

$$f(r_1, r_2) = \int_a^1 dx \frac{(1-x)^3}{(x-r_1^2)^2 + r_1^2 r_2^2} \left(\frac{1-a/x}{1-a/r_1^2}\right)^{1/2}, \quad r_1 = \frac{m_\sigma}{M}, \quad r_2 = \frac{\Gamma_\sigma}{M}, \quad a = \frac{4m_\pi^2}{M^2}.$$

Using the experimental information for the branching ratio for the radiative decay $\rho^0 \rightarrow \sigma\gamma \rightarrow \pi^0 + \pi^0 + \gamma$ [22], with the help of Eq. (6), one can deduce the constant $g_{\rho\sigma\gamma}$ as a function of the σ -meson parameters, its mass and its width (Fig. 1). One can see a strong dependence of the value of $g_{\rho\sigma\gamma}$ on the two coordinates r_1 and r_2 , which vary in the wide interval suggested by [20]. Moreover, in the whole region of parameters r_1 and r_2 one can see that $g_{\rho\sigma\gamma}^2 \leq 2.5$; for example, for $m_\sigma = 0.5$ GeV and $\Gamma_\sigma = 0.6$ GeV, we find $g_{\rho\sigma\gamma}^2 = 0.196$.

3 Estimation of $g_{V\sigma\gamma}$ couplings from $\sigma \rightarrow 2\gamma$

Another source for an independent estimation of the $g_{\rho\sigma\gamma}$ coupling constant, which confirms our previous finding, is the radiative decay $\sigma \rightarrow 2\gamma$. In principle it is possible to estimate the width of this decay analyzing the σ -contribution in the amplitude of the elastic scattering of photons by nucleons, in the framework of the t -channel exchange, Fig. 2a, or in the two-photon production of a $\pi^+\pi^-$ or $\pi^0\pi^0$ -pair in $\gamma\gamma$ -collisions, Fig. 2b, in the corresponding region of the $\pi\pi$ -effective mass [24].




Fig. 1. Two-dimensional plot of the coupling constant $g_{\rho\sigma\gamma}^2$ as a function of $r_1 = \frac{m_\sigma}{M}$ and $r_2 = \frac{\Gamma_\sigma}{M}$.




Fig. 2. Feynman diagrams for σ -meson contribution to nucleon Compton scattering (a) and to $\pi^+\pi^-$ -production in $\gamma\gamma$ -collisions (b).

Following the VDM approach, (Fig. 3), the matrix element for $\sigma \rightarrow 2\gamma$ can be written in terms of the $g_{V\sigma\gamma}$ coupling constants, as follows:

$$\mathcal{M}(\sigma \rightarrow 2\gamma) = e^2(g_\rho g_{\rho\sigma\gamma} + g_\omega g_{\omega\sigma\gamma})\vec{e}_1 \cdot \vec{e}_2 m_\sigma^2 / M, \quad (7)$$

where \vec{e}_1 and \vec{e}_2 are the three-vectors of polarization for the two produced photons. The coupling constant g_V determines the width of the V -meson leptonic




Fig. 3. Feynman diagrams for $\sigma \rightarrow 2\gamma$ -decay in the VDM.

decay $V \rightarrow e^+e^-$ (through one-photon exchange): $\Gamma(V \rightarrow e^+e^-) = \alpha^2 g_V^2 M \frac{4\pi}{3}$. Note that this formula holds for zero lepton mass. The existing data about $\rho(\omega) \rightarrow e^+e^-$ decays [20] allow to estimate g_ρ and g_ω with good accuracy: $g_\rho^2 \simeq 0.04$, $g_\omega^2 \simeq 0.09g_\rho^2$.

As both couplings $g_{\rho\sigma\gamma}$ and $g_{\omega\sigma\gamma}$ enter in Eq. (7), we assume, for simplicity, the validity of the $SU(3)$ relation: $g_{\rho\sigma\gamma}/g_{\omega\sigma\gamma} \simeq 3$. The width $\Gamma(\sigma \rightarrow 2\gamma)$ can be written as:

$$\Gamma(\sigma \rightarrow 2\gamma) = \pi\alpha^2 \frac{m_\sigma^3}{M^2} g_\rho^2 g_{\rho\sigma\gamma}^2 \left(1 + \frac{g_\omega}{3g_\rho}\right)^2 \simeq 13.7 g_{\rho\sigma\gamma}^2 \left(\frac{m_\sigma}{1 \text{ GeV}}\right)^3 \text{ keV}. \quad (8)$$

Taking $g_{\rho\sigma\gamma} \simeq 3$ [2], one finds

$$\Gamma(\sigma \rightarrow 2\gamma) = 123 \text{ keV}, \text{ for } m_\sigma = 1 \text{ GeV}. \quad (9)$$

The same or even larger numbers for $\Gamma(\sigma \rightarrow 2\gamma)$ follow from the QCD-sum rule estimation for $g_{\rho\sigma\gamma}$ [18] or from the value deduced from the analysis of $\rho^0 \rightarrow \pi^+\pi^-\gamma$ -decays [11]. These numbers contradict the corresponding experimental estimations:

$$\Gamma(\sigma \rightarrow 2\gamma) = 10.6 \text{ keV} [20], \quad \Gamma(\sigma \rightarrow 2\gamma) = (3.8 \pm 1.5) \text{ keV} [24]. \quad (10)$$

And, again, the discrepancy is large, up to one order magnitude. The situation is even worse, taking the value of $g_{\rho\sigma\gamma}$ from Ref. [13], which leads to

$$\Gamma(\sigma \rightarrow 2\gamma) = (340 \div 490) \text{ keV}. \quad (11)$$

On the other hand, taking the experimental data about the $\sigma \rightarrow 2\gamma$ -decay,

(10), one can deduce directly a value

$$0.28 \leq g_{\rho\sigma\gamma}^2 \left(\frac{m_\sigma}{1 \text{ GeV}} \right)^3 \leq 0.78, \quad (12)$$

in evident contradiction with the previous large $g_{\rho\sigma\gamma}$ values [2,11,13,18], but in good agreement with the present estimation, based on the new data about the decay $\rho \rightarrow \pi^0 \pi^0 \gamma$ [22].

4 Discussion and conclusions

Let us discuss the possible consequences of a smaller value of the $g_{\rho\sigma\gamma}$ coupling constant; more precisely, let us consider the case $g_{\rho\sigma\gamma}^2 \leq 1$. Here this range is derived from two independent sources: the radiative decay of the ρ^0 -meson, $\rho^0 \rightarrow \sigma\gamma \rightarrow \pi^0 + \pi^0 + \gamma$, on one side, and the radiative decay of the σ -meson: $\sigma \rightarrow 2\gamma$, from another side. Both these decays can be described in a relative simple and transparent theoretical framework: the σ -dominance for the first one and the VDM approach for the second one. Note that the VDM model gives a good description of the different numerous radiative decays involving vector and pseudoscalar mesons. The coupling constants g_ρ and g_ω (for the $\gamma \rightarrow V^0$ -transition), which enter in our consideration of the decay $\sigma \rightarrow 2\gamma$, are well known from the experimental data about the decays $V \rightarrow \ell^+ \ell^-$, which have quite good accuracy. Therefore, the main uncertainty in the determination of $g_{\rho\sigma\gamma}$ on the basis of the existing data on $\sigma \rightarrow 2\gamma$, derives from the relatively large interval for $\Gamma(\sigma \rightarrow 2\gamma)$. The parameters of the σ -meson also do not affect very much this estimation: there is no dependence on the σ -width, and only a cubic dependence on the σ -meson mass, so that:

$$g_{\rho\sigma\gamma}^2 = (g_{\rho\sigma\gamma}^2)_0 \left(\frac{m_\sigma}{1 \text{ GeV}} \right)^{-3}, \quad (13)$$

where $(g_{\rho\sigma\gamma}^2)_0$ is the value of the considered coupling constant for $m_\sigma=1 \text{ GeV}$. From this scaling law it follows the $g_{\rho\sigma\gamma}$ constant gets smaller for higher σ -masses.

The estimation of $g_{\rho\sigma\gamma}^2$ on the basis of the decay $\rho^0 \rightarrow \sigma + \gamma \rightarrow \pi^0 + \pi^0 + \gamma$ depends essentially on the mass and width of the σ -meson, only, without additional unknown coupling constants. We can build a two-dimensional representation on $g_{\rho\sigma\gamma}^2$, with non trivial dependence on m_σ and Γ_σ . It turns out that, for the 'standard' value $m_\sigma = 0.5 \text{ GeV}$, in the interval $0.6 \leq \Gamma_\sigma \leq 1 \text{ GeV}$, one has the following limits:

$$0.19 \leq g_{\rho\sigma\gamma}^2 \leq 0.30, \quad (14)$$

definitely smaller than the values used in [2] and [13,18]. On the other hand, the interval (14) is in agreement with the interval (12). We must stress that the value $g_{\rho\sigma\gamma}^2 = 9$ in the consideration of the process $\gamma + p \rightarrow p + \rho^0$ [2], has been found for $m_\sigma=0.5$ GeV, which has been chosen quite arbitrarily. The σ -exchange amplitude and the corresponding estimation of $g_{\rho\sigma\gamma}$ depend essentially on m_σ . For example, in the near-threshold conditions for $\gamma + p \rightarrow p + \rho^0$, another scaling law holds for $g_{\rho\sigma\gamma}$:

$$\frac{g_{\rho\sigma\gamma}^2}{(m_\sigma^2 + a)^2} = \text{const}, \quad (15)$$

where $a = M^2 m_N / (m_N + M) \simeq 0.32$ GeV² (m_N is the nucleon mass). This relation shows a large correlation between $g_{\rho\sigma\gamma}$ deduced from the data about $\gamma + p \rightarrow p + \rho^0$ and the σ -mass, in such a way that larger value of m_σ correspond to larger value of the coupling constant.

Moreover, the differential cross section for $\gamma + p \rightarrow p + \rho^0$, calculated in framework of σ -exchange, contains a strong dependence on the σ -mass, through the phenomenological form factor, which has to be introduced here to improve the t -behavior of the differential cross section $d\sigma(\gamma p \rightarrow p\rho^0)/dt$ at large $|t|$. To deduce a definite value of the $g_{\rho\sigma\gamma}$ coupling constant from a fit of the differential cross section data, it is also necessary to assume a definite value for the $g_{\sigma NN}$ -coupling. Typically this value is determined from the NN -potential.

The estimation of $g_{\rho\sigma\gamma}$ from the data about $\rho^0 \rightarrow \pi^+ + \pi^- + \gamma$, which suggests the large value: $g_{\rho\sigma\gamma}^2 > 10$, can not be considered very precise, as the possible σ -exchange is hidden by the large contribution of bremsstrahlung (photon radiation by charged pions). Therefore the coupling $g_{\rho\sigma\gamma}$ derived by this decay, results in a very large width for $\Gamma(\rho^0 \rightarrow \pi^0 + \pi^0 + \gamma)$, in strong contradiction with the experiment and with the other theoretical estimations of this width.

The HERMES-effect can not be considered a reliable source of information about the $g_{\rho\sigma\gamma}$ -coupling constant. It is more correct to say, that, in order to explain this effect in the framework of the model [13], one needs large values, which are difficult to justify on the basis of what is known from the radiative decays $V \rightarrow \pi\pi\gamma$. For example, the large value of $g_{\omega\sigma\gamma}$ was justified in [13] by taking the lower limit for $\omega \rightarrow \pi^+\pi^-\gamma$ from [20], instead of a much smaller and more precise value from $\omega \rightarrow \pi^0\pi^0\gamma$. Following selection rules, the relation $\Gamma(\omega \rightarrow \pi^+\pi^-\gamma) = 2\Gamma(\omega \rightarrow \pi^0\pi^0\gamma)$ [23] is correct independently on the decay mechanism. One deduces, for $g_{\omega\sigma\gamma}$, a value which is two orders of magnitude lower than in Ref. [13].

If the small values for the $g_{\rho\sigma\gamma}$ -coupling constant, as we suggest in the present paper are correct, the interpretation of the following different electromagnetic

processes has to be revised:

- The explanation of the near-threshold cross section for $\gamma + p \rightarrow p + \rho^0$ in frame of the σ -model can not work, because the σ -contribution has to be small. The σ -contribution, taking $g_{\rho\sigma\gamma}^2 \leq 1$, turns out to be one order of magnitude smaller than necessary for the explanation of the existing data. In principle, one can recover by increasing correspondingly the $g_{\sigma NN}$ -coupling constant, but this would seriously modify the NN -potential. Moreover, such increase would change the scale of the σ -contribution to the differential cross section of $\gamma + p \rightarrow p + \omega$, in the near threshold region, making the σ and π -exchanges of the same order. So other contributions, such as Pomeron and (or) f_2 -exchanges, which have been extrapolated up to threshold, have to be taken into account. This allows to decrease $g_{\rho\sigma\gamma}$ up to unity [5].
- The explanation of the HERMES effect, in terms of coherent contribution of mesons, does not hold anymore.
- The predictions of these constants in the framework of the QCD -sum rules do not hold.

5 Acknowledgment

We thank J. M. Laget for useful discussions on the problem of the $g_{\rho\sigma\gamma}$ -coupling constant in different electromagnetic processes. One of us (M.P.R.) acknowledges the kind hospitality of Saclay, where this work was completed.

6 Appendix

In this Appendix we discuss the effect of the σ -width on the decay $\rho \rightarrow \pi + \pi + \gamma$.

The matrix element, corresponding to this diagram (Fig. 4) can be written as:

$$\mathcal{M} = \mathcal{M}(V\sigma^*\gamma) \frac{1}{w^2 - m_\sigma^2} m_\sigma g_{\sigma\pi\pi}, \quad (16)$$

where $\mathcal{M}(V\sigma^*\gamma)$ is the matrix element for the decay $V \rightarrow \sigma^* + \gamma$, with production of a virtual σ -meson, which mass w (different from the mass of the σ -meson, m_σ), coincides here with the effective mass of the produced $\pi^+\pi^-$ -system, $m_\sigma \rightarrow m_\sigma - i\Gamma_\sigma/2$, Γ_σ is the total width of the σ -meson.

Taking the expression for $\mathcal{M}(V\sigma^*\gamma)$, Eq.(2), (where the $g_{V\sigma^*\gamma}$ coupling constant does not depend, following our assumption, on the virtuality of the




Fig. 4. Feynman diagrams for $V \rightarrow \pi^+ + \pi^- + \gamma$ decay through σ -meson production. (σ -meson), one can find, after summing over the photon polarizations and summing over the polarization of the V -meson:

$$|\overline{\mathcal{M}}(V\sigma^*\gamma)|^2 = e^2 \frac{g_{V\sigma^*\gamma}^2}{6} \frac{(M^2 - w^2)^2}{M^2}. \quad (17)$$

We are interested here in the $d\Gamma/dw^2$ -distribution (integrated over the effective $\pi^+\pi^-$ -mass) for the decay $V^0 \rightarrow \pi^+ + \pi^- + \gamma$ (we are considering here $\pi^+\pi^-$ -production to have different final pions):

$$d\Gamma = \frac{|\overline{\mathcal{M}}|^2}{2M} (2\pi)^4 \frac{d^3k}{(2\pi)^3 2E_\gamma} \int \frac{d^3k_1}{(2\pi)^3 2E_1} \frac{d^3k_2}{(2\pi)^3 2E_2} \delta(q - k - k_1 - k_2).$$

As $|\overline{\mathcal{M}}|^2$ depends only on the variable w^2 , it can be taken outside the integration, so that:

$$d\Gamma = \frac{|\overline{\mathcal{M}}|^2}{2^9 \pi^5 M} \frac{d^3k}{E_\gamma} \mathcal{I}, \quad (18)$$

where \mathcal{I} is the result of an invariant integration (which has to be done in the CMS of the produced $\pi^+\pi^-$ -system):

$$\mathcal{I} = \int \frac{d^3k_1}{E_1} \frac{d^3k_2}{E_2} \delta(q - k_1 - k_2) = 2\pi \frac{p^*}{E^*}, \quad E^* = \frac{w}{2}, \quad p^* = \sqrt{\frac{w^2}{4} - m_\pi^2} \quad (19)$$

So $\mathcal{I} = 2\pi\beta_w$, where $\beta_w = \frac{p^*}{E^*} = \sqrt{1 - 4m_\pi^2/w^2}$ is the velocity of the pion produced in the decay of the virtual σ -meson, $\sigma^* \rightarrow \pi^+ + \pi^-$, with mass w . It is the energy of the photon, produced in the decay $V \rightarrow \pi^+ + \pi^- + \gamma$, which determines the effective mass of the produced $\pi^+ + \pi^-$ -system, through the following relation:

$$M^2 = w^2 + 2ME_\gamma \quad (20)$$

which holds in the rest system of the decaying V -meson. Therefore we can write:

$$\frac{d^3k}{E_\gamma} \rightarrow 4\pi E_\gamma dE_\gamma = -\pi \left(1 - \frac{w^2}{M^2}\right) dw^2. \quad (21)$$

Substituting Eqs. (19) and (21) in (17) one can find:

$$\begin{aligned} \frac{d\Gamma}{dw} &= \frac{|\overline{\mathcal{M}}|^2}{2^8 \pi^3 M} \left(1 - \frac{w^2}{M^2}\right) \sqrt{1 - 4\frac{m_\pi^2}{w^2}} \\ &= \frac{e^2 g_{V\sigma^*\gamma}}{3\pi^3 2^9} M \left(1 - \frac{w^2}{M^2}\right)^3 \sqrt{1 - 4\frac{m_\pi^2}{w^2}} g_{\sigma^*\pi^+\pi^-} \frac{m_\sigma^2}{(w^2 - m_\sigma^2)^2} \end{aligned} \quad (22)$$

The coupling constant $g_{\sigma\pi^+\pi^-}$ can be related to the total width Γ_σ of the σ -meson, assuming that $\sigma \rightarrow 2\pi$ is its main decay: $\Gamma_\sigma = \Gamma(\sigma \rightarrow \pi^+\pi^-) + \Gamma(\sigma \rightarrow \pi^0\pi^0)$. In terms of the coupling constant $g = g_{\sigma\pi^+\pi^-} = g_{\sigma\pi^0\pi^0}$ (on the basis of isotopic invariance), one can write:

$$\Gamma(\sigma \rightarrow \pi^+\pi^-) = \frac{|\overline{\mathcal{M}}(\sigma \rightarrow \pi^+\pi^-)|^2}{8\pi m_\sigma} q = \frac{g^2}{16\pi} m_\sigma \beta_\sigma, \quad (23)$$

where $\beta_\sigma = \sqrt{1 - 4m_\pi^2/m_\sigma^2}$ is the velocity of the pion produced in the decay of the real σ -meson, with mass m_σ . Taking into account the relation: $\Gamma(\sigma \rightarrow \pi^+\pi^-) = 2\Gamma(\sigma \rightarrow \pi^0\pi^0)$, due to the identity of the two neutral pions in the decay $\sigma \rightarrow \pi^0\pi^0$, one can write:

$$\Gamma_\sigma = \frac{3}{2}\Gamma(\sigma \rightarrow \pi^+\pi^-) = \frac{3}{32} \frac{g^2}{\pi} m_\sigma \beta_\sigma, \quad (24)$$

Substituting Eq. (24) in (22) one obtains:

$$\frac{d\Gamma}{dw^2} = \frac{e^2 g_{V\sigma^*\gamma}}{144\pi^2} M \left(1 - \frac{w^2}{M^2}\right)^3 \frac{\beta_w}{\beta_\sigma} \frac{m_\pi \Gamma_\sigma}{(w^2 - m_\sigma^2)^2 + \Gamma_\sigma^2 m_\sigma^2} \quad (25)$$

So, for the full width of the decay $V \rightarrow \pi^+ + \pi^- + \gamma$, the following formula holds:

$$\Gamma(V \rightarrow \pi^+\pi^-\gamma) = \int_{4m_\pi^2}^{M^2} \frac{d\Gamma}{dw^2} dw^2. \quad (26)$$

For the decay $V \rightarrow \pi^0 + \pi^0 + \gamma$, Eq. (26) holds, divided by a coefficient 2 - due to the identity of pions in the final state.

Let us test the validity of these formulas, considering the limit of zero width, $\Gamma_\sigma \rightarrow 0$, i.e. considering the σ -meson as a stable particle. We use the following symbolic relation:

$$\begin{aligned} \left| \frac{1}{(w^2 - m_\sigma^2) - i\Gamma_\sigma m_\sigma} \right|^2 &= \frac{1}{(w^2 - m_\sigma^2) + i\Gamma_\sigma m_\sigma} \frac{1}{(w^2 - m_\sigma^2) - i\Gamma_\sigma m_\sigma} \xrightarrow{\Gamma_\sigma \rightarrow 0} \\ &= \frac{1}{(w^2 - m_\sigma^2) + i\Gamma_\sigma m_\sigma} \left[\frac{1}{w^2 - m_\sigma^2} + i\pi\delta(w^2 - m_\sigma^2) \right] \rightarrow \frac{\pi}{m_\sigma\Gamma_\sigma} \delta(w^2 - m_\sigma^2) \end{aligned}$$

This δ -function allows one to integrate easily Eq. (26), getting the following result:

$$\Gamma(V \rightarrow \sigma\gamma \rightarrow \pi^+\pi^-\gamma) = \frac{\alpha}{36} g_{V\sigma\gamma}^2 M \left(1 - \frac{m_\sigma^2}{M^2} \right)^3. \quad (27)$$

To find the radiative width for the radiative decay $V \rightarrow \sigma\gamma$, taking into account both possibilities, $\sigma \rightarrow \pi^+ + \pi^-$ and $\sigma \rightarrow \pi^0 + \pi^0$, we have to introduce a factor 3/2, so that:

$$\Gamma(V \rightarrow \sigma\gamma) = \frac{\alpha}{24} g_{V\sigma\gamma}^2 M \left(1 - \frac{m_\sigma^2}{M^2} \right)^3$$

in agreement with the direct calculation, see Eq. (3).

References

- [1] F. Iachello and D. Kusnetzov, Phys. Rev. D **45**, 4156 (1992).
- [2] B. Friman and M. Soyeur, Nucl. Phys. A **100**, 477 (1996).
- [3] Q. Zhao, Z. Li, and C. Bennhold, Phys. Rev. C **58**, 2393 (1998); Phys. Lett. B **436**, 42 (1998).
- [4] A. I. Titov, T. S. H. Lee, H. Toki and O. Streltsova, Phys. Rev. C **60**, 035205 (1999).
- [5] J. M. Laget, Phys. Lett. B **489**, 313 (2000).
- [6] H. Babacan, T. Babacan, A. Gokalp and O. Yilmaz, Eur. Phys. J. A **13**, 355 (2002).
- [7] M. Chemtob, E. J. Moniz and M. Rho, Phys. Rev. C **10**, 344 (1974).

- [8] E. Hummel and J. A. Tjon, Phys. Rev. Lett. **63** , 1788 (1989); Phys. Rev. C **42**, 423 (1990).
- [9] J. W. Van Orden, N. Devine and F. Gross, Phys. Rev. Lett. **75**, 4369 (1995).
- [10] S. M. Renard, Nuovo Cimento **62A**, 475 (1969).
- [11] A. Gokalp and O. Yilmaz, Phys. Rev. D **62**, 093018 (2000); Phys. Lett. B **508**, 25 (2001).
- [12] K. Ackerstaff *et al.* [HERMES Collaboration], Phys. Lett. B **475**, 386 (2000).
- [13] G. A. Miller, S. J. Brodsky and M. Karliner, Phys. Lett. B **481**, 245 (2000).
- [14] A. Yu. Korchin, O. Scholten, R. G. E. Timmermans, Phys. Lett. B **438**, 1 (1998).
- [15] V. Pascalatsa, O. Sholten, Nucl. Phys. A **591**, 658 (1995).
- [16] I. Guiasa, C. Pomponia, E. E. Radesca, Ann. Phys. (NY), 114, 296 (1979).
- [17] R. Machleidt, Adv. Nucl. Phys. **19**, 189 (1989).
- [18] A. Gokalp and O. Yilmaz, Phys. Rev. D **64**, 034012 (2001); T. M. Aliev, A. Ozpineci and M. Savci, Phys. Rev. D **65**, 076004 (2002).
- [19] M. P. Rekalo and E. Tomasi-Gustafsson, to appear in Nucl. Phys. A
- [20] D. E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).
- [21] S. Fajfer and R. J. Oakes, Phys. Rev. D **42**, 2392 (1990);
A. Bramon, A. Grau and G. Pancheri, Phys. Lett. B **283**, 416 (1992);
A. Bramon, A. Grau and G. Pancheri, Phys. Lett. B **289**, 97 (1992);
E. Marco, S. Hirenzaki, E. Oset and H. Toki, Phys. Lett. B **470**, 20 (1999);
A. Bramon, R. Escribano, J. L. Lucio Martinez and M. Napsuciale, Phys. Lett. B **517**, 345 (2001).
- [22] M. N. Achasov *et al.*, Phys. Lett. B **537**, 201 (2002).
- [23] O. Singer, Phys. Rev. 128, 2789 (1962), Phys Rev 130, 2441 (1963); Phys. Rev. 161, 1694 (1967).
- [24] M. Boglione and M. R. Pennington, Eur. Phys. J. C **9**, 11 (1999).