

NLTE Solar Irradiance Modeling with the COSI code

A. I. Shapiro¹, W. Schmutz¹, M. Schoell^{1,2}, M. Haberreiter³, and E. Rozanov^{1,4}

¹ Physikalisch-Meteorologisches Observatorium Davos, World Radiation Center, 7260 Davos Dorf, Switzerland
e-mail: alexander.shapiro@pmodwrc.ch

² Institute for Astronomy ETH, Zurich, Switzerland

³ Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO 80303, USA

⁴ Institute for Atmospheric and Climate science ETH, Zurich, Switzerland

Received ; accepted

ABSTRACT

Context. The solar irradiance is known to change over time scales from minutes to decades, and it is suspected that its substantial fluctuations are partially responsible for climate variations in the past.

Aims. We are developing a solar atmosphere code which allows the physical modeling of the entire solar spectrum composed of quiet Sun and active regions. This code is a tool for modeling the temporal variability of the solar irradiance and understanding its influence on Earth.

Methods. We exploit a further development of the radiative transfer code COSI that now incorporates the calculation of molecular lines. We validate COSI under the conditions of local thermodynamic equilibrium (LTE) against the synthetic spectra calculated with the ATLAS code. The synthetic solar spectra are also calculated in non-local thermodynamic equilibrium (NLTE) and compared to the available measured spectra. Doing so we define the main problems of the modeling, e.g. the lack of opacity in the UV part of the spectrum and the inconsistency in the calculations of the visible continuum level, and we describe a solution to these problems.

Results. The improved version of COSI allows us to reach a good agreement between the calculated and observed solar spectrum as measured by SOLSTICE and SIM onboard the SORCE satellite and ATLAS 3 mission operated from the Space Shuttle. We find that NLTE effects are very important for the modeling of the solar spectrum even in the visual part of the spectrum and for its variability over the entire solar spectrum. In addition to the strong effect on the UV part of the spectrum, NLTE effects influence the concentration of the negative ion of hydrogen which results in a significant change of the visible continuum level and the irradiance variability.

Key words. Line: formation – atomic data – Molecular processes – Sun: atmosphere – Sun: UV radiation: Radiative transfer

1. Introduction

The solar radiation is the main source of the input of energy to the terrestrial atmosphere and therefore, it determines Earth's thermal balance and climate. Although it is known since 1978 that the solar irradiance is not constant but varies on scales from several minutes to decades (cf. Fröhlich 2005; Krivova & Solanki 2008), the influence of this variability to the climate is not yet fully understood. Nowadays several datasets for the past spectral solar irradiance (SSI) based on the different reconstruction approaches (e.g. Lean et al. 2005; Krivova et al. 2009) and satellite measurements are available. However the remaining disagreements between these data lead to different atmospheric responses when they are used in the climate models (Shapiro et al. 2009). The task of constructing a self-consistent physical model in order to reconstruct the past Solar Spectral Irradiance (SSI) remains of high importance.

Modern reconstructions of the SSI are based on the assumption that the irradiance changes are determined by the evolution of the solar surface magnetic field (Foukal & Lean 1988; Krivova et al. 2003; Domingo et al. 2009). Areas of the solar disk are associated to several components (e.g. quiet Sun, bright network, plage and sunspot) according to the measured surface magnetic field and the contrast of the features. These components are represented by corresponding atmosphere structures

(cf. Kurucz 1991; Fontenla et al. 1999) and therefore different emergent radiation. The SSI is calculated by weighting the irradiance from each model with the corresponding filling factor.

The calculation of the emergent solar radiation, even from the atmosphere with known thermal structure, is a very sophisticated problem because consistent models have to account for the NLTE effects in the solar atmosphere. As the importance of these effects became clear over the last several decades, several numerical codes were developed. One of the first NLTE-codes, LINEAR, was published by Auer et al. (1972) who used a complete linearization method developed by Auer & Mihalas (1969, 1970). Later, the MULTI code published by Carlsson (1986). This code is based on the linearization technique developed by Scharmer (1981) and Scharmer & Carlsson (1985a,b). More recently, the RH code, which is based on the MALI (Multi-level Approximate Lambda Iteration) formalism of Rybicki & Hummer (1991, 1992) was developed by Uitenbroek (2001).

Combining the radiative transfer code by (Hamann & Schmutz 1987; Schmutz et al. 1989) and SYNSPEC by Hubeny (1981), Haberreiter et al. (2008) developed the 1D spherical symmetrical COde for Solar Irradiance (COSI). The NLTE opacity distribution function (ODF) concept, which was implemented in this code, allows to indirectly account for the NLTE effects in several millions lines. This made COSI especially suitable for calculating the overall energy distribution in the solar spectrum. In this paper we introduce a new version of COSI (COSI version 2) describe the main modifications, and present first results.

In Sect. 2 we describe the datasets of the measured spectral irradiance which were used for comparison with our calculated spectrum. In Sect. 3 we introduce the integration of molecular lines into COSI and we show that it can solve the discrepancies to observations and to other codes which has been described in Haberreiter et al. (2008). In Sect. 4 we present the resulting solar spectrum from NLTE calculations. Due to missing opacity in the UV the flux appears to be significantly higher than measured (Busá et al. 2001; Short & Hauschildt 2009). We solve this problem by introducing additional opacity to the ODF for selected spectral ranges (Sect. 4.2), while the problem of the NLTE visible continuum due to the deviations in the concentration of the hydrogen negative ion is addressed in Sect. 4.3. In Sect. 5 we present the synthetic spectra of the active regions and its implications to the solar variability study. Finally, we summarize the main results in Sect. 6.

2. Measured solar spectra irradiance

We have compare our calculated solar spectrum with several available datasets. We use the observations taken with the Solar-Stellar Irradiance Comparison Experiment (SOLSTICE) (McClintock et al. 2005) up to 320 nm and the Solar Irradiance Monitor (SIM) (Harder et al. 2005) from 320 nm onward instruments onboard the SORCE satellite (Rottman 2005) obtained during the 23th solar cycle minimum (hereafter SORCE measurements). For the comparisons shown in this paper we use the average of the observations from 21 April 2008 to 28 April 2008.

Our calculated spectrum was converted with 1 nm box-car profile for comparison with the SOLSTICE measurements and trapezoidal profile for comparison with the SIM measurements. The SIM resolution strongly depends on wavelength (it is about 1.5 nm for the 310 nm and about 22 nm for the 800 nm) so the parameters of this trapezoidal profile are also wavelength dependent and were provided by by Harder (2009).

We also used the SOLar SPECTral Irradiance Measurements (SOLSPEC) (Thuillier et al. 2004) during the ATLAS 3 mission in November 1994. For this comparison we convolved the calculated spectrum with a Gaussian with FWHM = 0.6 nm.

3. Molecules in the COSI code

COSI simultaneously solves the equations of statistical equilibrium and radiative transfer in 1D spherical geometry. All spectral lines in COSI can be divided into two groups. The first group comprises about one thousand lines that are most prominent in the solar spectrum. These lines are explicitly treated in NLTE. The second group contains several millions background lines provided by Kurucz (2006) and calculated under the assumption that their upper and lower levels are populated in LTE relative to the explicit NLTE levels. These background lines are taken into account in the spectral synthesis part of the code but also affect NLTE calculations via the ODF. The ODF is iteratively recalculated until it becomes self-consistent with populations of the NLTE levels (Haberreiter 2006; Haberreiter et al. 2008). This allows us to indirectly account for the NLTE effects in the background lines.

In the previous version of COSI (Haberreiter et al. 2008) molecular lines were not included. However, molecular lines play an important role in the formation of the solar spectrum. In some spectral regions they are even the dominant opacity source. Therefore, a considerable amount of opacity was missing. This led to a discrepancy with observations. In the present version




Fig. 1. The ratio R of the number concentration of carbon and oxygen which are not attached to the molecules to their total element amount (upper panels) and the CN and CO concentrations as a function of height (lower panels) for three atmospheric models: FALA99 (dashed curves), FALC99 (continuous curves), and FALP99 (dotted curves). The zero-height depth point in all models is defined as the radius at which the continuum optical depth at 5000Å is equal to one.

of COSI molecular lines are taken into account. This requires the computation of molecular concentrations, preparation of the molecular line lists, and incorporation of molecular opacities and emissivities into the code.

3.1. Chemical equilibrium

Under the assumption of the instantaneous chemical equilibrium the concentration of molecule AB is connected with the concentrations of consisting atoms A and B by the Guldberg-Waage law (cf. Tatum 1966):

$$\frac{n_A n_B}{n_{AB}} = K_{AB}(T) = \left(\frac{2\pi m k T}{h^2} \right)^{3/2} \frac{Q_A Q_B}{Q_{AB}} kT \exp(-D_0/kT), \quad (1)$$

where K_{AB} is the temperature dependent equilibrium constant, Q is internal partition function for the atoms A and B, and molecule AB, m is the reduced molecular mass, and D_0 is the dissociation energy. The contribution to the opacity from the transitions between lower level i and upper level j of molecule AB is given by:

$$\kappa_{AB}^{ij}(\nu) = \frac{n_{AB} g_i \exp(-E_i/kT) h\nu}{Q_{AB}} \frac{1}{4\pi} (n_i B_{ij} - n_j B_{ji}) \phi(\nu - \nu_{ij}), \quad (2)$$

where g_i is statistical weight of the lower level, B 's are the Einstein coefficients and $\phi(\nu)$ is the line profile normalized to one.

There are several available temperature polynomial approximations for the atomic and molecular partition functions as well as for the equilibrium constants K_{AB} (cf. Tatum 1966; Tsuji 1973; Irwin 1981; Sauval & Tatum 1984; Rossi et al. 1985). Although molecular partition functions can contain significant errors due to the inaccurate values of the energy levels and even due to unknown electronic states (Irwin 1981), this does not affect significantly the accuracy of molecular opacity calculations. This insensitivity is due to the fact that the overall molecular

concentration is proportional to the partition function (see Eq. 1), while the opacity in a given line is inversely proportional to it (see Eq. 2). Therefore it is crucial to either calculate equilibrium constants directly from partition functions or use the same source for both approximations.

Tsuji (1973) shows that abundances of hydrogen, carbon, nitrogen, and oxygen can be determined quite accurately ignoring their dependence on other elements. As we are interested only in molecular bands which can significantly contribute to the opacity over a broad spectral range, the system of equations like Eq. (1) were solved only for these four main elements and the diatomic molecules built from them.

As a significant fraction of atoms can be associated with molecules the chemical equilibrium calculation also affects the atomic lines. In Fig. 1 we present the changes in the carbon and oxygen concentrations due to the association with molecules together with CN and CO concentrations for three atmospheric models by Fontenla et al. (1999): the relatively cold supergranular cell center model FALA (hereafter FALA99), the averaged quiet Sun model FALC (hereafter FALC99) and the relatively warm bright network model FALP (hereafter FALP99). Both molecular concentrations and deviations in atomic concentration show strong temperature sensitivities which is important for assessing of the solar spectral variability.

3.2. Main molecular bands in the solar spectrum

The line list for the spectrum synthesis was compiled using the molecular databases of Kurucz (1993) and the SRPM database, which is based on Gray & Corbally (1994), and HITRAN. Wavelengths and line strengths of the most significant lines of the CN violet system and CH G band were calculated based on the molecular constants by Krupp (1974), Knowles et al. (1988), and Wallace et al. (1999).

In Fig. 2 we present the part of the solar spectrum where the LTE calculations with the previous version of COSI, which did not account for molecular lines, showed significant deviations from the calculations with the LTE radiative transfer code ATLAS12 carried out by Kurucz (2005). Both calculations used the same atmosphere structure and abundances by Kurucz (1991) (hereafter K91). One can see that introducing the molecular lines into COSI quite strongly affects the spectrum and significantly diminishes disagreements with the ATLAS 12 code. The most prominent features are the CH G band around 430 nm, CN violet G band around 380 nm, CN, NH, and OH bands between 300 and 350 nm. The remaining differences can be possibly attributed to the different photoionization cross sections implemented in COSI and ATLAS12.

In Fig. 3 we compare the spectra calculated by both codes with the SORCE measurements. One can see that both codes predict the correct value of the continuum in the red part of the visible spectrum (600 nm) but fail to reproduce the molecular bands and UV part of the spectrum. The LTE spectra calculated with the FALC99 and two sets of abundances by Grevesse & Anders (1991) (hereafter G91) and by Asplund et al. (2005) (hereafter A05) are presented in Fig. 4. The calculations with the G91 abundances predict the correct value of the visible and IR continuum. Furthermore, they give quite a good fit of molecular bands (see right panel of the picture where the calculation of CN violet system and CN G band are shown in more detail). However the calculated irradiance in the UV part is again much higher than observed for both abundance sets.

The metal abundances by A05 are significantly lower (up to about 2 times) than in G91. The lower metal abundance leads




Fig. 2. LTE calculation of molecular lines with COSI. Upper panel: Synthetic solar spectrum calculated with (solid line) and without (dashed line) molecular lines. Lower panel: Corresponding ratios to the irradiance calculated by the ATLAS12 code. All spectra are averaged with a 1-nm boxcar.




Fig. 3. The irradiance calculated by COSI in LTE (thin solid line) and the ATLAS 12 (dashed line), using K91 atmosphere structure and abundances. The spectrum observed by SORCE is given by the thick solid line. All spectra are convolved with the instrument profile of SORCE.

to a significantly lower electron density and consequently to a lower concentration of negative hydrogen ion. The latter is the main source of the continuum opacity in the solar atmosphere (cf. Mihalas 1978, p. 102). Therefore the application of the A05 abundances results in a lower continuum opacity for the visible and IR wavelengths and as a consequence in a higher irradiance. This effect is especially prominent at longer wavelengths (from about 400 nm) where the number of strong lines is relatively low and the opacity of negative hydrogen dominates over all other sources of opacity. In turn, the lower metal abundance given by

A05 leads to lower line opacities and photoionization opacities in the UV. Followingly, the A05 abundances also lead to a higher irradiance in the UV than the G91 abundances.

4. NLTE calculations

4.1. Introduction

Most of the solar radiation emerges from regions of the atmosphere with a steep temperature gradient. Therefore a significant part of the escaping photons is not in the thermodynamic equilibrium with the surrounding medium. This implies that a self-consistent radiative transfer model has to account for NLTE effects. It is well-known that these effects are of extreme importance for the strong lines and the UV radiation. In this section we present NLTE calculations of the solar spectrum with COSI and show that the NLTE effects are also important for the overall energy distribution in the solar spectrum, including the visible and infrared wavelength ranges.

In the left panel of Fig. 5 the NLTE calculations of the solar spectrum with the atmosphere structure FALC99 and the abundance sets A05 and G91 are compared to the observed solar spectrum. The right panel of Fig. 5 shows the ratios of the NLTE to the LTE calculations. The most prominent overall NLTE effects are the strong increase of the irradiance in the UV and mild decrease of the irradiance in the red part of the spectrum. The first effect is discussed in details in Sect. 4.2, while the latter in Sect. 4.3.

4.2. Calculation of the UV radiation

4.2.1. NLTE opacity effects

Independent of the solar atmosphere model and abundances, the LTE calculations overestimate the UV irradiance up to about 300 nm (see Figs. 3 and 4). This problem is even more severe in the NLTE calculations as they predict a UV irradiance that is higher than in the case of the LTE calculations (see left panel of Fig. 5).

The effect of the increase of the far UV flux in the NLTE calculations is actually well-known (cf. Short & Hauschildt 2009). The UV radiation incident from the higher and hotter parts of the solar atmosphere onto the photosphere causes a stronger ionization of iron and other metals relative to the LTE case. This decreases the populations of the neutral atoms and consequently weakens the strength of the corresponding line and photoionization cross section, which are the main sources of the opacity in the far-UV. Therefore it leads to a decrease of the opacity and consequently an increase of the flux in the wavelengths up to about 250 nm (red threshold of the photoionization from the Mg I 2 level). In the longer wavelengths the line and the photoionization opacities of the ionized metals become more important so the increase of their populations results in lower irradiance.

The effect of the increased UV flux in the NLTE calculations can be decreased significantly by the use of the opacity distribution function (see Fig. 6), which allows to incorporate the opacity from the background lines into the solution of statistical equilibrium equations (Haberreiter et al. 2008). This background opacity diminishes the amount of the penetrating UV radiation and decreases the degree of metal ionization. It leads to the relative decrease of the spectral irradiance up to 250 nm and an increase at higher wavelengths (in agreement with the above discussion). The complicated wavelength dependence of the spectrum alterations caused by the influence of the ODF can be explained by the large number of the strong metal lines and

photoionization thresholds. Overall the introduction of the ODF causes the redistribution of the solar irradiance over the entire spectrum as it decreases the far-UV part and increases near-UV and visible.

Although the concept of the ODF allows to improve the agreement between calculations and observations, the ODF based on the Kurucz linelist (Haberreiter et al. 2008) can not solve the discrepancy from measurements.

The most probable source of the problem is inaccuracy of the atomic and molecular line list and possible missing continuum opacity. The production of a complete line list is an extremely laborious task. Considerable progress was reached during last decades thanks to the effort of the OPACITY project (Seaton 2005) in calculating atomic data and in particular of Kurucz (1993). Concerning the problem of the UV blanketing it is important to keep in mind that about 99%(!) of the atomic and molecular lines are predicted only theoretically and only 1% of all lines are observed in the laboratory (Kurucz 2005). This leads to potential errors in the total opacity as lines could still be missing as well as wavelengths and oscillator strengths of the predicted lines could be inaccurate. An incompleteness of the atomic data is especially significant in the UV where the immense number of lines form the UV line haze.

The consistent account for the line blanketing effect is very important for the overall absolute flux distribution, especially for the UV (cf. Collet et al. 2005; Avrett & Loeser 2008; Fontenla et al. 2009). Busá et al. (2001) used version 2.2 of the MULTI code, which UV opacity were contributed only from several hundred calculated in the NLTE lines, bound-free and free-free processes. They showed that NLTE calculations without proper account for the line blanketing effect can lead to the overestimation of the UV flux by six orders of magnitude in the case of cold metal-rich stars. In order to compensate this, they multiplied the continuum opacity by a wavelength-dependent factor and gave explicit expression for the factor parameterization. The similar approach was used by Bruls (1993) to calculate the NLTE populations of Ni I. Short & Hauschildt (2009) showed that the use of different line lists for the line blanketing calculations can lead to a significantly different spectral irradiance distributions. They also showed that the problem of the excess of the UV flux in the 300-420 nm spectral region can be solved by increasing of the continuum absorption coefficient.

4.2.2. Additional opacities in COSI

COSI takes into account the opacity from the several million lines in the UV (Haberreiter et al. 2008). However the significant disagreement between the calculated and measured solar UV flux as discussed above indicates that the computations still miss an essential part of the opacity. In order to compensate for this missing opacity Haberreiter et al. (2008) calculated the UV part of the solar spectrum with artificially increased Doppler broadening. This approach proves itself successful. However it cannot be used to fit high-resolution spectra. Therefore, here we have extended the approach by Busá et al. (2001) and Short & Hauschildt (2009) and included additional opacity in our NLTE-ODF scheme, as described below.

COSI overestimates the irradiance i.e. shows an opacity deficiency in the spectral region between 160 nm and 320 nm. At shorter wavelengths the opacity is dominated by continuum photoionization, while at longer wavelengths tabulated atomic and molecular lines are sufficient to reproduce the observations. In this wavelength region we grouped the solar spectrum into 1 nm intervals (corresponding to the resolution of the




Fig. 4. Spectra calculated by COSI in LTE using the FALC99 atmosphere structure and A05 and G91 abundances, vs. SORCE measurements. All spectra are convolved with the instrument profile of SORCE.




Fig. 5. Left panel: as the left panel in Fig. 4 but calculated in NLTE. Right panel: The ratio between NLTE and LTE emergent flux.




Fig. 6. Left panel: NLTE calculations with and without iterated ODF, compared to the solar irradiance as observed by SORCE. Right panel: Ratio between the emergent flux from NLTE calculated with and without iterated ODF. All calculations are done with the FALC atmosphere model and G91 abundances.

SOLSTICE/SORCE spectrum). For each interval we empirically determined the coefficient $f_c(\lambda)$ by which the continuum opacity has to be multiplied to reproduce the SOLSTICE measurements. As the correction of the ODF affects the statistical equilibrium of the populations we iteratively solve the statistical equilibrium equations and the factors $f_c(\lambda)$ using newly updated ODFs until we reached a self-consistent solution. Thus, the $f_c(\lambda)$ factors

were found iteratively. We stopped the iteration when the deviations to the observed spectrum was smaller than 10%. In the considered region there were very few wavelength points where the irradiance was underestimated. We did not apply any correction to these points. This underestimation can be explained by the inaccuracy of the line list as it can be caused by a small wavelength inaccuracy of a few strong lines.




Fig. 7. The factor $f_c(\lambda)$ (upper panel) and the ratio of the additional opacity to and total opacity at the height 200 km (lower panel) for the G91 and A05 abundances.

The wavelength dependence of $f_c(\lambda)$ is shown in Fig. 7 for the calculations with FALC99 atmosphere model and G91 and A05 sets of abundances. One can see that $f_c(\lambda)$ is very close to 1 (which means little additional opacity) for wavelengths longer than 280 nm. There are also several prominent peaks which indicate strong lack of opacity at specific wavelengths. Although different abundances lead to a change of the peak amplitudes, the overall profile of wavelength dependence stays the same for both sets of abundances, G91 and A05, respectively. The averaged line opacity is larger by several order of magnitude than the continuum opacity. So the ratio of the additional opacity, which was needed to achieve agreement with the observed solar spectrum, to the total opacity which was already included into COSI is for most wavelength below 5 % (see lower panel of Fig. 7). Therefore, the correction of the overestimation of the emergent UV flux can be achieved by a moderate modification of the input line list.

Let us also note that calculations with the G91 abundances lead to a lower continuum than calculations with A05 abundances. Therefore the flux overestimation is stronger in the case of calculations with A05 abundances. Consequently one has to use larger correction factors $f_c(\lambda)$ for A05 abundances.

The correction factor strongly depends on the chosen line list. We tested our calculations using the Vienna Atomic Line Database (VALD) (Kupka et al. 1999, 2000) and retrieving the lines with known radiative damping parameter. As this database contains a smaller number of lines than provided by Kurucz (2006) a larger factor $f_c(\lambda)$ is necessary to achieve an agreement with the observations

4.3. Calculation of the visible and IR radiation

NLTE effects decrease the visible and near IR irradiance by about 10 % (see Fig. 5). The reason is that NLTE effects influence the concentration of electrons and the negative hydrogen ions which determines the continuum opacity as well as the continuum source function.

4.3.1. NLTE effects in electron and negative hydrogen concentrations

The deviations in the electron and hydrogen negative ion concentrations are presented in Fig. 8 for the calculations with FALC99 atmosphere structure and A05 abundances. The upper panel illustrates the main electron sources in the solar atmosphere. In the lower part of the photosphere and in the chromosphere hydrogen is ionized and provides most of the electrons. However in the photosphere, which is responsible for the formation of the visible and near-IR irradiance, the temperature is too low for ionizing hydrogen. Therefore, here the electrons are mainly provided by the metals with low ionization potential, such as iron, silicon and magnesium.

As one can see from the second upper panel of Fig. 8 the ionization of silicon and iron are strongly affected by the NLTE effects. This is basically the same effect of NLTE “overionization” which has already been discussed in Sect. 4.2, where the main emphasis was on the deviations of the concentration of the neutral atoms. Let us notice that applying of the ODF significantly decreases the NLTE effects on the electron concentration as it strongly increases the UV opacity. From the last part of Fig. 8, where the contribution functions for the continuum radiation are plotted for different μ -positions, one can see that although the continuum is mainly formed in the lower part of the photosphere where the NLTE effects in the electron concentration are not so prominent, they still can be important especially for the radiation emitted close to the solar limb.

The photoionization cross section of negative hydrogen has its threshold at about 16 500 Å and it reaches its maximum at about 8500 Å (cf. Mihalas 1978, p. 102). Although the photosphere is optically thick for the UV and radiation in the strong atomic or molecular lines, the visible and IR photons can easily escape even from the lower part of the photosphere (the optical depth of the zero point in the continuum radiation is about one, depending on the wavelengths). This means that all over the photosphere there is a lack of photons which are able to photoionize the negative hydrogen. Thus, in the NLTE case, the radiative destruction rate of H^- is decreased, which in turn, increases the concentration of the negative hydrogen ion compared to the LTE equilibrium. This is illustrated in the third panel of Fig. 8. We note that the effect is especially strong in the layers where the continuum radiation is formed (see the contribution functions in the lower panel of Fig. 8). Moreover, in contrast with effects on the electron concentration, the applying of the ODF does not decrease the NLTE deviations of the negative hydrogen concentration as UV radiation does not have significant contribution to its ionizations.

Compared to the LTE case, NLTE calculations result in higher electron and negative hydrogen concentrations. This increases the continuum opacity so that continuum radiation comes from higher and cooler layers of the photosphere. In addition, there is also a decrease of the NLTE continuum source function which contributes to the decrease of the continuum emission.

The effects described above significantly decrease the level of the emergent flux in visible and near IR. Although the NLTE treatment is physically more consistent than LTE, it leads to a severe deviation from the observed solar spectrum (see Fig. 5).

4.3.2. Collisions with negative hydrogen

One possible cause for the inability of the NLTE calculations to reproduce the measurements is the inadequate treatment of col-




Fig. 8. NLTE effects in electron and H^- concentrations. Upper panel: Depth dependence of the total electron concentration and the electron concentrations resulting from the ionization of elements as indicated. Second panel: Ratios of the change of the electron concentrations due to the NLTE effects and total electron concentration calculated in LTE. Plotted are total change without and with applying ODF and resulting from the ionization of particular elements change without ODF. Third panel: depth dependence of the ratio of negative hydrogen and electron concentration calculated in NLTE and LTE. Lower panel: Contribution functions for the 5000 Å (dotted lines) continuum radiation and temperature on depth dependence (solid curve) for the FALC99 atmosphere model. The contribution functions are plotted for eight values of the cosine of the angle between the propagation direction of radiation and the local solar radius μ . From right to left: $\mu = 0.05$ (almost solar limb), 0.1, 0.2, 0.4, 0.6, 0.8, 0.9, 1.0 (solar disk center). All calculations are carried out for the FALC99 atmosphere model and A05 abundances

lisional rates for the negative hydrogen ion. The present version of COSI accounts for the electron detachment through collisions with electrons and neutral hydrogen as well as charge neutralization with protons as given by Lambert & Pagel (1968). We do not treat collisions of H^- with heavy elements, which in principle could lead to a strong underestimation of the collisional rates.

We found that increasing the collisional rates involving H^- by a factor of ten solves the disagreement between the measured and calculated continuum flux for the models using A05 abundances. However, when using increased collisional rates, the

molecular bands (especially CH G band) appear to be weaker than the observed ones. To make the molecular lines consistent with observations we changed the A05 carbon and nitrogen abundances and used them as given by G91. This allows us to reach a very good agreement with measurements (see Sect. 4.4).

Let us emphasize that an increase of the collisional rates for H^- cannot help to adjust the NLTE calculations performed with G91 abundances. The reason is that the considered increase of collisions can only enforce a LTE population ratio between negative hydrogen and H I 1 level. The latter is however in a NLTE regime mainly due to the NLTE “overionization”. As the calculations with G91 abundances give a good agreement in a pure LTE regime (see Fig. 4) the considered increase of the collisional rates for H^- is not enough to enforce full LTE agreement. A NLTE treatment of the solar atmosphere yields population numbers of negative hydrogen that strongly deviate from LTE.

4.3.3. Change of abundances

The effect of the use of different abundances on the emergent synthetic spectrum is illustrated in Fig. 9. One can see that the abundance change of elements like iron, magnesium and silicon strongly affects the UV radiation (mainly due to the photoionization opacities) but also the visible and infrared continuum because these elements are the main donors of the photospheric electrons. In contrast, a change of nitrogen and oxygen abundances has no effect on the continuum and only affects the spectrum via the molecular bands. The carbon abundance slightly influences the continuum but the main effect of carbon in the visible spectral range is due to molecular systems: The CN violet system at about 380 nm and CH G band at about 430 nm are the most prominent features. Thus, the carbon and nitrogen abundance changes from the A05 to G91 values almost do not affect the continuum level (see Fig. 9) and lead to a good agreement with the measurements (see Fig. 10). Therefore, we can conclude that A05 abundances should be given preference for the calculation of the NLTE continuum, except for carbon and nitrogen, for which the G91 abundances lead to a better reproduction of the molecular bands. The favored combination of abundances leads to a good overall agreement with the observed solar spectrum.

4.4. Comparison with measurements and discussion

In Fig. 10 we present the solar spectrum calculated with the FALC99 atmosphere model compared with the SORCE measurements. The agreement in the 160–320 nm spectral region is automatically good (see Sect. 4.2.2). At shorter wavelengths the observed flux is determined by several strong emission lines which are treated in LTE in the current version of COSI. This leads to a strong underestimation of the irradiance. In a future version of COSI these lines will be treated in NLTE. The Ly α line is already computed in NLTE and we obtain an overestimation of its flux. In a forthcoming paper we will investigate it in more detail (Schoell et al. 2010).

The detailed comparison of our synthetic spectrum with SOLSPEC measurements is given in Figs. 13 and 14 in the Online Material. One can see that starting from 160 nm two spectra are in remarkable agreement with each other. Although several strong emission lines are currently not properly calculated in shorter wavelengths, the continuum level there is also consistent with SOLSPEC measurements. This can be achieved only with NLTE calculations.




Fig. 9. Ratios of the emergent fluxes calculated with A05 abundances to the fluxes calculated with the abundances of the indicated element increased by a factor of two.

Thus the assumption of enhanced negative hydrogen collisions and the use of the combined G91 and A05 abundances set provide us with the possibility to successfully model the entire solar spectrum. Another possible cause of the problem that synthetic NLTE models have difficulty to fit the observed solar spectrum is that the FALC99 atmosphere model was developed to fit the visible and IR continuum with the radiative transfer code PANDORA (see Fontenla et al. 1999), who used a different approximate approach for the treatment of the NLTE effects, different sets of the atomic and molecular data as well as different sets of abundances. The Fontenla et al. (1999) atmosphere structures were successfully used in various applications (cf. Penza et al. 2004a,b; Vitas & Vince 2005) in which however the continuum flux was calculated in LTE. The quest of constructing a self-consistent atmosphere model which fully accounts for all NLTE effects will be addressed in a future investigation.

5. Irradiance variations

It is generally accepted that most of the solar variability is introduced by the competition between the solar flux decrease due to the dark sunspots and a flux increase due to the bright magnetic network and plage (e.g. Willson & Hudson 1991; Krivova et al. 2003). Therefore it is important to consistently calculate the solar irradiance from these active components of the solar atmosphere.

For our calculations we use the Fontenla et al. (1999) atmosphere model FALC99 for the quiet Sun, model FALF99 for the bright network, model FALP99 for plage, and model FALS99 for sunspot. Both NLTE and LTE spectra are presented in Fig. 11. The NLTE spectra produce a lower visible and UV flux and ac-

ordingly, lower total solar irradiance (TSI). This effect for the quiet Sun irradiance was already discussed in Sect. 4.3.

An further interesting detail is the decrease of the contrast between the quiet Sun, plage and bright network in the NLTE calculations. The main reason for this is that the NLTE effects which decrease the continuum source function is stronger in the hotter FALP model as it corresponds to the lower particle density (and consequently lower collisional rates).

Fig. 12 shows the flux differences between the active components of the solar atmosphere and the quiet Sun calculated in LTE with A05 abundances and NLTE with enhanced collisions with negative hydrogen and combined A05 and G91 abundances (see Sect. 4). The most prominent peak at about 3890 Å corresponds to the CN violet system. Although the differences in the CH G band around 4300 Å are also clearly visible, they are less pronounced than in the CN violet system. This is mainly due to the differences in the CH and CN dissociation energies (3.465 eV and 7.76 eV), so the CN concentration is more sensitive to temperature variations (see Sect. 3.1). Furthermore, the contrast between the active components and the quiet Sun is significantly decreased in the NLTE calculations. As a consequence the NLTE calculations reduce the solar variability in the visible and IR, shifting it to the UV. As the corrected NLTE calculations do not change the visible and IR continuum they are free from this effect.

The differences between the spectral fluxes for active components and for the quiet Sun integrated from 900 Å to 40 000 Å are presented in Table 1 for the LTE (A05 and G91 abundances), NLTE with G91 abundances and NLTE with enhanced collisions with negative hydrogen and combined A05 and G91 abundances (NLTE comb.). We conclude that the NLTE calculations (both




Fig. 10. Upper panel: SORCE observations vs. calculations with FALC atmosphere model and described in the text abundances. Lower panel: ratio between the calculations and measurements.




Fig. 11. NLTE (left panel) and LTE (right panel) calculations of the solar irradiance from the quiet Sun, bright network, plage and sunspot.

Table 1. TSI differences with FALC model per 1 % area (in W/m^2).

	FALF	FALP	FALS
LTE G91	0.28	1.48	-10.7
LTE A05	0.25	1.27	-10.82
NLTE G91	0.18	0.76	-11.02
NLTE comb.	0.15	0.99	-11.05

with and without ODF) significantly reduce the variations of the TSI.

6. Conclusions

We present a further development of the radiative transfer code COSI. The code accounts for the NLTE effects in several hundred lines, while the NLTE effects in the other several millions lines are indirectly included via iterated opacity distribution function. The radiative transfer is solved in spherical symmetry. The main conclusions can be summarized as follows:

- The inclusion of the molecular lines into COSI allows us to reach a good agreement with the SORCE observations in the main molecular bands (especially in the CN violet system and CH G band). It also solves the previous discrepancies be-




Fig. 12. The flux differences between plage and quiet Sun (upper panel) and bright network and quiet Sun (lower panel) calculated in NLTE and LTE.

tween the LTE calculations with the COSI code and ATLAS 12 calculations. We show that due to their strong temperature sensitivity, molecular lines significantly contribute to the solar irradiance variability.

- We introduce additional opacities to the opacity distribution function. It allows us to solve well-known problem of the overestimation of the UV flux in the synthetic spectrum.
- We show that the concentration of negative hydrogen is strongly affected by NLTE effects. It decreases the level of the visible and infrared continuum and leads to a discrepancy with the measured level if the calculations are done with the current models of the quiet Sun atmosphere.
- We present calculations of the total and spectral solar irradiance changes due to the presence of the active regions. We show that NLTE effects can strongly affect both of these values.

Acknowledgements. The investigations have benefited from stimulating discussions at meetings organized at ISSI in Bern, Switzerland. The research leading to this paper has received funding from the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement N 218816 (SOTERIA). We thank J. Fontenla for supplying the SRPM database.

References

- Asplund, M., Grevesse, N., Sauval, A. J., Allende Prieto, C., & Blomme, R. 2005, *A&A*, 431, 693
- Auer, L. H., Heasley, J. N., & Milkey, R. W. 1972, A computational program for the solution of non-LTE transfer problems by the complete linearization method, 55–+
- Auer, L. H. & Mihalas, D. 1969, *ApJ*, 158, 641
- Auer, L. H. & Mihalas, D. 1970, *MNRAS*, 149, 65
- Avrett, E. H. & Loeser, R. 2008, *ApJS*, 175, 229
- Bruls, J. H. M. J. 1993, *A&A*, 269, 509
- Busá, I., Andretta, V., Gomez, M. T., & Terranegra, L. 2001, *A&A*, 373, 993
- Carlsson, M. 1986, *Upps. Astron. Obs. Rep.*, 33
- Collet, R., Asplund, M., & Thévenin, F. 2005, *A&A*, 442, 643
- Domingo, V., Ermolli, I., Fox, P., et al. 2009, *Space Science Reviews*, 145, 337
- Fontenla, J., White, O. R., Fox, P. A., Avrett, E. H., & Kurucz, R. L. 1999, *ApJ*, 518, 480
- Fontenla, J. M., Curdt, W., Haberreiter, M., Harder, J., & Tian, H. 2009, *ApJ*, 707, 482
- Foukal, P. & Lean, J. 1988, *ApJ*, 328, 347
- Fröhlich, C. 2005, *Memorie della Societa Astronomica Italiana*, 76, 731
- Gray, R. O. & Corbally, C. J. 1994, *AJ*, 107, 742
- Grevesse, N. & Anders, E. 1991, *Solar element abundances*, 1227–1234
- Haberreiter, M. 2006, PhD thesis, Physikalisch-Meteorologisches Observatorium Davos, World Radiation Center, Dorfstrasse 33, 7260 Davos, Switzerland
- Haberreiter, M., Schmutz, W., & Hubeny, I. 2008, *A&A*, 492, 833
- Hamann, W.-R. & Schmutz, W. 1987, *A&A*, 174, 173
- Harder, J. 2009, personal communication
- Harder, J. W., Fontenla, J., Lawrence, G., Woods, T., & Rottman, G. 2005, *Sol. Phys.*, 230, 169
- Hubeny, I. 1981, *A&A*, 98, 96
- Irwin, A. W. 1981, *ApJS*, 45, 621
- Knowles, P. J., Werner, H.-J., Hay, P. J., & Cartwright, D. C. 1988, *J. Chem. Phys.*, 89, 7334
- Krivova, N., Solanki, S., Wenzler, T., & Podlipnik, B. 2009, *Journal of Geophysical Research*, in press
- Krivova, N. A. & Solanki, S. K. 2008, *Journal of Astrophysics and Astronomy*, 29, 151
- Krivova, N. A., Solanki, S. K., Fligge, M., & Unruh, Y. C. 2003, *A&A*, 399, L1
- Krupp, B. M. 1974, *ApJ*, 189, 389
- Kupka, F., Piskunov, N., Ryabchikova, T. A., Stempels, H. C., & Weiss, W. W. 1999, *A&AS*, 138, 119
- Kupka, F. G., Ryabchikova, T. A., Piskunov, N. E., Stempels, H. C., & Weiss, W. W. 2000, *Baltic Astronomy*, 9, 590
- Kurucz, R. L. 1991, in *NATO ASIC Proc. 341: Stellar Atmospheres - Beyond Classical Models*, 441–+
- Kurucz, R. L. 1993, CD 13
- Kurucz, R. L. 2005, *Memorie della Societa Astronomica Italiana Supplement*, 8, 86
- Kurucz, R. L. 2006, personal communication

- Lambert, D. L. & Pagel, B. E. J. 1968, *MNRAS*, 141, 299
- Lean, J., Rottman, G., Harder, J., & Kopp, G. 2005, *Sol. Phys.*, 230, 27
- McClintock, W. E., Snow, M., & Woods, T. N. 2005, *Sol. Phys.*, 230, 259
- Mihalas, D. 1978, *Stellar Atmospheres*, 2nd edn. (San Francisco: Freeman and Company)
- Penza, V., Caccin, B., & Del Moro, D. 2004a, *A&A*, 427, 345
- Penza, V., Caccin, B., Ermolli, I., & Centrone, M. 2004b, *A&A*, 413, 1115
- Rossi, S. C. F., Maciel, W. J., & Benevides-Soares, P. 1985, *A&A*, 148, 93
- Rottman, G. 2005, *Sol. Phys.*, 230, 7
- Rybicki, G. B. & Hummer, D. G. 1991, *A&A*, 245, 171
- Rybicki, G. B. & Hummer, D. G. 1992, *A&A*, 262, 209
- Sauval, A. J. & Tatum, J. B. 1984, *ApJS*, 56, 193
- Scharmer, G. B. 1981, *ApJ*, 249, 720
- Scharmer, G. B. & Carlsson, M. 1985a, *Journal of Computational Physics*, 59, 56
- Scharmer, G. B. & Carlsson, M. 1985b, in *NATO ASIC Proc. 152: Progress in Stellar Spectral Line Formation Theory*, ed. J. E. Beckman & L. Crivellari, 189–198
- Schmutz, W., Hamann, W.-R., & Wessolowski, U. 1989, *A&A*, 210, 236
- Schoell, M., Schmutz, W., Haberreiter, M., & Shapiro, A. I. 2010, *A&A*, in preparation
- Seaton, M. J. 2005, *MNRAS*, 362, L1
- Shapiro, A. V., Rozanov, E., Egorova, T., et al. 2009, *J. Atmos. Solar-Terrest. Phys.*, submitted
- Short, C. I. & Hauschildt, P. H. 2009, *ApJ*, 691, 1634
- Tatum, J. B. 1966, *Publications of the Dominion Astrophysical Observatory Victoria*, 13, 1
- Thuillier, G., Floyd, L., Woods, T. N., et al. 2004, in *Washington DC American Geophysical Union Geophysical Monograph Series*, Vol. 141, *Solar Variability and its Effects on Climate*. Geophysical Monograph 141, ed. J. M. Pap, P. Fox, C. Frohlich, H. S. Hudson, J. Kuhn, J. McCormack, G. North, W. Sprigg, & S. T. Wu, 171–+
- Tsuji, T. 1973, *A&A*, 23, 411
- Uitenbroek, H. 2001, *ApJ*, 557, 389
- Vitas, N. & Vince, I. 2005, *Memorie della Societa Astronomica Italiana*, 76, 1064
- Wallace, L., Hinkle, K., Li, G., & Bernath, P. 1999, *ApJ*, 524, 454
- Willson, R. C. & Hudson, H. S. 1991, *Nature*, 351, 42

Online Material




Fig. 13. The spectra calculated with COSI (solid line) vs. SOLSPEC measurements (dotted line).




Fig. 14. As Fig. 13. Continuation.