
Surface rotation period extraction plots KOIs


Plots marked a, b, c,... correspond to light curves obtained with the KADACS corrections using a filter of 30 days applied to the whole light curves.


Plots marked a', b', c',... correspond to light curves obtained with the KADACS corrections using a filter of 55 days applied quarter by quarter before stitching the quarters together.


For both corrections, the plots are as follows:


- a: Original light curve (light blue) and rebinned light curve (black). The vertical dotted lines mark the limits of the *Kepler* quarters.
- **b**: Power density spectrum of the rebinned light curve, normalised to its maximum.
- c: Wavelet Decomposition of the rebinned light curve. The red/black areas mark the most relevant periods while the blue/white areas mark the less relevant ones. The hatched region corresponds to the unreliable zone due to border effects.
- d: Global Wavelet Power Spectrum (i.e. the Wavelet Decomposition projected onto the periods axis) of the rebinned light curve, normalised to its maximum (black) and multiple gaussian functions fit (green). The horizontal dotted line marks the period retruned by this method.
- e: Autocorrelation Function (ACF) of the rebinned light curve (black) and smoothed ACF (blue). The vertical dotted line marks the period retruned by this method.
- **f**: Composite Spectrum of the rebinned light curve (black) and multiple gaussian functions fit (green). The vertical dotted line marks the period retruned by this method.





















KIC 9955598









KIC 11807274