

The cooling of atomic and molecular gas in DR21

H. Jakob¹, C. Kramer¹, R. Simon¹, N. Schneider^{4,2,1}, V. Ossenkopf^{1,3}, S. Bontemps², U. U. Graf¹, and J. Stutzki¹

¹ KOSMA, I. Physikalisches Institut, Universität zu Köln, Zùlpicher StraÙe 77, 50937 Köln, Germany

² Observatoire de Bordeaux, Université de Bordeaux 1, BP 89, 33270 Floirac, France

³ SRON, National Institute for Space Research, PO Box 800, 9700 AV Groningen, the Netherlands

⁴ SAp/CEA Saclay, 91191 Gif-sur-Yvette, France

Received 19 June 2006 / Accepted 11 September 2006

ABSTRACT

Aims. We present an overview of a high-mass star formation region through the major (sub-)mm, and far-infrared cooling lines to gain insight into the physical conditions and the energy budget of the molecular cloud.

Methods. We used the KOSMA 3m telescope to map the core ($10' \times 14'$) of the Galactic star forming region DR 21/DR 21 (OH) in the Cygnus X region in the two fine structure lines of atomic carbon ($C\text{ I }^3P_1-^3P_0$ and $^3P_2-^3P_1$) and four mid- J transitions of CO and ^{13}CO , and CS $J = 7-6$. These observations have been combined with FCRAO $J = 1-0$ observations of ^{13}CO and C^{18}O . Five positions, including DR21, DR21 (OH), and DR21 FIR1, were observed with the ISO/LWS grating spectrometer in the [O I] 63 and 145 μm lines, the [C II] 158 μm line, and four high- J CO lines. We discuss the intensities and line ratios at these positions and apply Local Thermal Equilibrium (LTE) and non-LTE analysis methods in order to derive physical parameters such as masses, densities and temperatures. The CO line emission has been modeled up to $J = 20$.

Results. From non-LTE modeling of the low- to high- J CO lines we identify two gas components, a cold one at temperatures of $T_{\text{kin}} \sim 30 - 40\text{ K}$, and one with $T_{\text{kin}} \sim 80 - 150\text{ K}$ at a local clump density of about $n(\text{H}_2) \sim 10^4 - 10^6\text{ cm}^{-3}$. While the cold quiescent component is massive containing typically more than 94 % of the mass, the warm, dense, and turbulent gas is dominated by mid- and high- J CO line emission and its large line widths. The medium must be clumpy with a volume-filling of a few percent. The CO lines are found to be important for the cooling of the cold molecular gas, e.g. at DR21 (OH). Near the outflow of the UV-heated source DR21, the gas cooling is dominated by line emission of atomic oxygen and of CO. Atomic and ionised carbon play a minor role.

Key words. ISM: clouds – ISM: abundances – Radio lines: ISM – Line: profiles – Stars: formation – ISM: individual objects: DR 21, DR21 (OH)

1. Introduction

Emission lines of C II, C I, and CO in molecular clouds are among the most important diagnostic probes of star formation and particularly important for studying the influence of massive stars and their UV radiation on the environment. Obtaining the physical parameters of such regions is often done with simplified radiative transfer assumptions. Local Thermal Equilibrium (LTE) and the Large Velocity Gradient (LVG) analysis have proven to yield good results for the bulk of the cold gas (see, e.g., Black (2000)). However, the assumption of a homogeneous medium does not represent the typical physical conditions of a high mass star formation region. Instead, the observing beam will contain several gas components with differ-

ent excitation conditions and with unknown filling factors. The observed line transitions are sensitive to gas at different density, temperature and opacity, corresponding to different regions. By selecting designated lines as tracers of the different parts of the gas, one can obtain a rather complete coverage of excitation conditions accounting for different physical and also chemical conditions within the beam. A more differentiated analysis using non-LTE radiative transfer algorithms (see van Zadelhoff et al. (2002) for an overview) is needed for this purpose. It can provide a more detailed description of the structure of the cloud including clumpiness and temperature or density gradients (e.g., Ossenkopf et al. (2001), Williams et al. (2004)).

Using the results of this analysis, we can try to retrieve a self-consistent picture including all major processes driving and driven by high mass star formation (e.g. with respect to turbulence or chemistry). Thus, we obtain a

Send offprint requests to: H. Jakob

Correspondence to: jakob@ph1.uni-koeln.de




Fig. 1. View of the DR21/DR21(OH) high mass star forming region. *Left:* Overlay of the KOSMA ^{13}CO 3–2 map (cf. Schneider et al. (2006)/SBS2006) in contours on a Spitzer IRAC $8\ \mu\text{m}$ greyscale map (Marston et al. 2004). Contours are on a T_{mb} scale and range from 16.8 to 100.8 in steps of $16.8\ \text{K km s}^{-1}$ ($\sigma = 5.6\ \text{K km s}^{-1}$). The integration interval is -10 to $20\ \text{km s}^{-1}$. The left-top circle indicates the HPBW of KOSMA. The map also shows 6 positions of radio continuum sources identified as OB stars by Roelfsema et al. (1989) (white crosses), 2 infrared sources and maser positions (Garden et al. 1986) (small white circles), and 8 compact 1.3-mm dust emission sources (Chandler et al. 1993) (open rectangles). *Right:* Overlay of the Spitzer map at $4.5\ \mu\text{m}$ with contours of the KOSMA ^{12}CO 3–2 map (integration interval of -20 to $30\ \text{km s}^{-1}$; discussed in SBS2006) ranging from 18 to 450 in steps of $48\ \text{K km s}^{-1}$ ($\sigma = 6\ \text{K km s}^{-1}$). The labeled circles mark the positions and beam sizes of the ISO/LWS observations that we use to constrain the physical parameters as discussed in the text. The LWS beam is comparable to the resolution of the KOSMA maps.

detailed view of the Photon Dominated Regions (PDRs) in the environment of young stars. PDRs are transition regions from ionized and atomic to dense molecular gas in which physics and chemistry of the gas are dominated by the UV radiation from young massive stars. Processes heating and cooling the clouds are qualitatively well understood on a theoretical level (e.g., Tielens & Hollenbach 1985; Sternberg & Dalgarno 1989), and in general we find a reasonable agreement between the theoretical predictions and observations of the brightest cooling lines. Still, there is an ongoing debate about the details of the energy input contributing to the heating. We find a combination of mechanisms such as stellar far-ultraviolet radiation (FUV), dynamical energy through shocks or supernovae, or an enhanced cosmic ray rate. There are also many open questions with respect to the chemical composition of these regions. The role of nonequilibrium chemistry is still not well understood. For instance Polycyclic Aromatic Hydrocarbons (PAHs) as a reservoir of fresh gas phase carbon (e.g., Oka et al. (2004), Habart et al. (2005)) may change the overall picture.

Emission from warm carbon in the form of atoms, ions or in carbonated molecules (mainly CO) provides a significant fraction of the gas cooling at sub-millimeter and far-infrared (FIR) wavelengths (e.g., Giannini et al. (2000, 2001), Schneider et al. (2003), Kramer et al. (2004), or Bradford et al. (2003, 2005)). Finding the prevailing cooling channels among these and other species (e.g., [OI], H_2O , OH) may provide significant insight into the details of the chemical composition and the energy balance of PDRs. Since systematic observations of sub-mm- and FIR-line cooling lines with a good spatial resolution are still sparse, data to test the model predictions are still largely missing and many questions concerning the details of the chemistry and energy balance remain unanswered. By studying a well-known, massive Galactic star forming region, we aim at gaining new insight into these problems.

The DR21 region

The DR21 H II-region/molecular cloud is part of the Cygnus X complex of molecular clouds. Cygnus X is known to host a large number of H II-regions

Table 1. List of observational parameters for KOSMA. The columns give line frequency, receiver used, main beam efficiency η_{mb} , half power beam width, observing mode, number of mapped positions, velocity channel width Δv (the resolution is a factor of ~ 1.4 larger), mean atmospheric zenith opacity $\langle \tau_0^{\text{atm}} \rangle$, and observing period.

Species	Transition	Frequency [GHz]	Receiver	η_{mb}^a	HPBW ^a [']	Mode	No.	Δv [m s ⁻¹]	$\langle \tau_0^{\text{atm}} \rangle$	Observing period
¹³ CO ^b	$J = 3-2$	330.5880	DualSIS	0.68	80	OTF ^c	693	308	0.26	01/2006
¹² C ³² S	$J = 7-6$	342.8829		0.68	80	OTF ^c	525	294	0.08	12/2005
¹² CO	$J = 3-2$	345.7960		0.68	80	OTF ^c	525	294	0.08	12/2005
¹² CO	$J = 4-3$	461.0408	SMART	0.5	57	OTF ^c	3458	677	0.85 – 1.4	1/2005
[¹² C I]	³ P ₁ – ³ P ₀	492.1607		0.5	55	DBS ^d	1380	630	0.5 – 1.6	1/2003 – 3/2004
¹³ CO	$J = 6-5$	661.067		0.4	~ 40	PSw ^e	33	307	1.0 – 1.3	1/1998
¹² CO	$J = 6-5$	691.473		0.4	~ 40	PSw ^e	58	295	1.0 – 1.3	1/1998
¹² CO	$J = 7-6$	806.6518		0.31	42	DBS ^d	964	775	0.4 – 1.6	1/2003 – 3/2004
[¹² C I]	³ P ₂ – ³ P ₁	809.3420		0.31	42	DBS ^d	964	775	0.4 – 1.6	1/2003 – 3/2004

^a The main beam efficiency η_{mb} and the HPBW were determined from cross scans of Jupiter.

^b See Schneider et al. (2006) for a ¹³CO 3–2 map covering a larger part of Cygnus X.

^c OTF – On-The-Fly mapping mode with periodic switching to a emission free reference position at $\alpha(J2000) = 20^{\text{h}}37^{\text{m}}10^{\text{s}}$ $\delta(J2000) = 42^{\circ}30'00''$.

^d DBS – Dual Beam Switch observing mode with a chopping secondary mirror in azimuth.

^e The CO and ¹³CO 6–5 observations were carried out on a 35'' grid (Köster 1998) in a Position-Switched observing mode (PSw). The OFF position was at $\alpha(J2000)=20^{\text{h}}26^{\text{m}}43^{\text{s}}.56$, $\delta(J2000)=42^{\circ}09'39''.5$.

(Downes & Rinehart 1966; Wendker 1984) associated with molecular clouds as seen, e.g., in the CfA ¹²CO 1–0 survey (Leung & Thaddeus (1992)). An extended (11 square degrees) map of the Cygnus X region in the ¹³CO 2–1 line and smaller maps in the CO and ¹³CO 3–2 lines were obtained with the KOSMA telescope and are presented in Schneider et al. (2006) (from now on SBS2006). The distance to DR 21 was sometimes estimated to be ~ 3 kpc, based on visual interstellar extinction and radial velocity (e.g., Campbell et al. 1982). For this paper, we follow the argumentation from SBS2006 based on a morphological comparison of the molecular line data with mid-IR emission and the OB associations, and favour a value of 1.7 kpc as distance toward DR21 and DR21 (OH). However, for easier comparison with the literature, we explicitly give distance depended physical properties as a function of distance.

The DR 21 region has been subject to numerous studies at different wavelengths. Several active star formation sites and cometary shaped H II-regions were identified in maps of radio continuum emission (e.g., Dickel et al. 1983; Cyganowski et al. 2003). Ammonia observations by Mauersberger et al. (1985) were interpreted as evidence for high densities ($\sim 10^{5.5} \text{ cm}^{-3}$) paired with a high degree of clumpiness. H₂O masers (Genzel & Downes 1977) and a recently obtained map of 1.3-mm continuum emission (Motte et al. 2005) show that DR 21 belongs to a north-south orientated chain of *massive star forming complexes*. Additional indication of active star formation is also given by observations of vibrationally excited H₂, tracing the hot, shocked gas in the vicinity of DR 21 and DR21 (OH) (e.g., Garden et al. (1986), and more recent results by Davis et al. (2006)). H₂ images reveal an ex-

tended ($5' = 2.5 \text{ pc} (D/1.7 \text{ kpc})$) north-east to south-west orientated outflow centered on DR21. To the west, the outflow is highly collimated, whereas toward the east side, gas is expanding in a blister style (Lane et al. 1990). Jaffe et al. (1989) found very intense CO 7–6 emission peaking on the H II-region and following the H₂-emission.

In this paper, we present the results of sub-mm mapping observations with KOSMA and discuss far-IR line intensities observed with ISO. This comprehensive data set allows to study the spatial and partly also the kinematic structure of all major cooling lines of the ISM in the vicinity of DR 21 and DR 21(OH). In Sect. 3, we analyse the line emission and continuum data toward five positions, by comparing the spatial distribution, line profiles, and the physical properties through selected line ratios and the far-IR dust continuum. In Sect. 4, we model the integrated line intensity ratios and absolute intensities with radiative transfer models. The impact of the line emission on the total cloud cooling is discussed in Sect. 5. A summary is given in Sect. 6.

2. Observations

2.1. KOSMA

The region centered on DR21/DR21 (OH) was mapped in the two atomic carbon fine structure transitions at 492 GHz (609 μm , ³P₁–³P₀; hereafter 1–0) and 809 GHz (370 μm , ³P₂–³P₁; hereafter 2–1) and the mid- J rotational transitions of CO ($J=3-2$, $J=4-3$, $J=6-5$, and $J=7-6$), ¹³CO ($J=3-2$, and $J=6-5$) using the dual channel 345 & 230/660 GHz SIS receiver, and the SubMillimeter Array Receiver for Two frequencies




Fig. 2. Integrated intensity map of CO 4–3 (*left*) and CO 7–6 (*right*) of DR 21 & DR 21 (OH). The integration interval is -20 to 30 km s^{-1} . The contours for CO 4–3 run from $66(3\sigma)$ to 462 K km s^{-1} in steps of 3σ , and from $68(2\sigma)$ to 671 K km s^{-1} in steps of 2σ for CO 7–6. The HPBW is indicated in the upper left corner. The positions of ISO/LWS observations (cf. Fig. 1) are indicated by circles. The dashed white lines correspond to the cuts shown in Fig. 6.

Table 2. Positions of ISO observations in absolute coordinates. DR 21 C is centered on the H II-region. Projected distance of DR 21 E (resp. DR 21 W) from position DR 21 C is 0.77 pc (resp. 0.89 pc) at a distance of 1.7 kpc. The TDT-number is given in the second column.

Name	TDT-No.	R.A. [J2000]	Dec. [J2000]
DR 21 E/East	15200785	$20^{\text{h}}39^{\text{m}}09^{\text{s}}.14$	$42^{\circ}20'05''.0$
DR 21 C/Center	15200786	$20^{\text{h}}39^{\text{m}}00^{\text{s}}.93$	$42^{\circ}19'42''.0$
DR 21 W/West	15200787	$20^{\text{h}}38^{\text{m}}52^{\text{s}}.23$	$42^{\circ}19'00''.5$
DR 21 (OH)	34700439	$20^{\text{h}}39^{\text{m}}00^{\text{s}}.95$	$42^{\circ}22'37''.6$
DR 21 FIR1	35500317	$20^{\text{h}}39^{\text{m}}00^{\text{s}}.2$	$42^{\circ}24'34''.6$

(SMART) at the KOSMA-3m telescope. SMART is a dual-frequency, eight-pixel SIS-heterodyne receiver that simultaneously observes in the 650 and 350 μm atmospheric windows (Graf et al. 2002). All observations were taken in the period between 2003 and 2006.

An area of $\sim 10' \times 14'$ was mapped in the two [C I] lines and the CO 7–6 line on a $27.5'' \times 27.5''$ grid with a total integration time of 160 s per position. The observations before 2005 were performed in Dual-Beam-Switch mode (DBS), with a secondary mirror chop throw of $6'$ fixed in azimuth. Due to the extent of the DR 21 cloud complex, self-chopping is present in some of the DBS-mode observations. In Appendix B, we describe how we processed these data in order to reduce the impact of the self-chopping. CO and ^{13}CO 3–2, and CO 4–3 were ob-

served in an On-The-Fly (OTF) mapping mode on a fully-sampled grid. A zeroth order baseline was subtracted from the calibrated spectra. In addition to CO 3–2, we simultaneously obtained the CS 7–6 line in the image sideband of the 345 GHz channel. Pointing was frequently checked on Jupiter, the sun, and then confirmed on DR 21 itself. The resulting accuracy was better than $\pm 20''$. Larger ^{13}CO 3–2 and 2–1 maps are presented in SBS2006. Due to their lower spatial resolution, we did not consider the $J=2-1$ data here.

The ^{12}CO and ^{13}CO 6–5 observations were presented in Köster (1998). The typical DSB receiver temperatures in the high frequency channel was between 160 – 220 K. The ^{13}CO 6–5 observations cover a region of $\sim 3' \times 3'$ centered on DR 21 while the ^{12}CO observations extend further to the north covering a region of $\sim 3' \times 6'$.

Atmospheric calibration was done by measuring the atmospheric emission at the reference position to derive the opacity. Sideband imbalances were corrected using standard atmospheric models (Cernicharo 1985) assuming a receiver sideband gain of 0.5 . Spectra taken simultaneously in the two frequency bands near 492 and 810 GHz were calibrated with an averaged water vapor value. The [C I] 2–1 and CO 7–6 lines, observed simultaneously in two sidebands, were also corrected for sideband imbalance. Their pointing is identical and the relative calibration error between these two transitions is very small. We scaled the antenna temperature data to the T_{mb} scale using main beam efficiencies η_{eff} as listed in Table 1. The HPBWs ($40''$ to $80''$) and η_{mb} were determined using continuum




Fig. 3. Integrated intensity map of [C I] 1–0 (*left*) and of [C I] 2–1 (*right*). The integration interval is -10 to 20 km s^{-1} . The contours for [C I] 1–0 run from $12(2\sigma)$ to 60 K km s^{-1} in steps of 2σ , and from $20(1\sigma)$ to 108 K km s^{-1} in steps of σ for [C I] 2–1.

scans on Jupiter. From observations of standard calibration sources, we estimate that the absolute calibration is accurate to within $\sim 15\%$. Most spectra were taken under excellent weather conditions with typical values of the atmospheric opacity along the line-of-sight better than 1.0 at 350 and 600 μm , and better than 0.5 above 870 μm .

2.2. ISO LWS Observations and Data Reduction

ISO Long Wavelength Spectrometer (LWS, Clegg et al. (1996)) $43 - 197$ μm grating scans (AOT L01) were obtained for 5 positions in the DR21 region from the ISO Data Archive (IDA) (see Table 2 and Fig. 7). We removed duplicate scans, overlapping positions, or data with a very poor spectral resolution from the sample.

The grating scans contain the FIR continuum, the atomic fine structure lines of [O I] at 63 and 145 μm , ionised atomic lines of [C II] at 158 μm , [O III] at 52 and 88 μm , [N II] at 122 μm , and the high- J CO molecular lines $J=14-13$ through $J=17-16$. All spectra were processed with the ISO Spectral Analysis Package (ISAP, v. 2.1). In ISAP, the data were deglitched by hand, defringed (detectors 4–9), and corrected for flux clipping (the extended source correction). Some artefacts remain after the processing (see Fig. 7), and were not corrected. Line strengths were measured within ISAP. See Table 4 for further details.




Fig. 4. Integrated intensity map of CS 7–6. The integration interval is -10 to 4 km s^{-1} . Contours run from $1.8(3\sigma)$ to 13.0 K km s^{-1} (in steps of 3σ).

2.3. FCRAO

We use ^{13}CO and C^{18}O 1–0 data from a large scale mapping project of the whole Cygnus X region using the FCRAO 14m telescope. These data were obtained be-




Fig. 5. Spectra of all observed species at the five ISO positions. *Grey filled spectra outlined in black:* observed lines, *thin grey lines:* line profiles modeled with the radiative transfer code SIMLINE (see Sect. 4.2). All spectra are shown at a common spatial resolution of $80''$ and on a T_{mb} scale. The baselines are shifted in steps of 30 K. [C I] was multiplied by a factor of 2, CS 7–6 and C^{18}O 1–0 by a factor of 4. $J = 1-0$ transition data are FCRAO observations. The dashed lines indicate the central velocity of -3 km s^{-1} , and the dotted lines the secondary velocity component at about 9 km s^{-1} , which was excluded from the fitting.

tween 2003 December and 2006 January and will be presented in more detail in upcoming papers (Simon et al., in prep.). The data were observed using the 32 pixel array SEQUOIA in an On-The-Fly mapping mode (at an angular resolution of $\sim 50''$). In this paper, we use smoothed data to have the same angular resolution of $80''$ as the KOSMA data.

3. Results

Figure 1 gives an overview centered on DR21 (OH) (designated as W75 S by some authors). The Spitzer $8 \mu\text{m}$ and $4.5 \mu\text{m}$ data by Marston et al. (2004) show the small scale structure of this region at angular resolutions between $1.3''-3''$. Several bright IR objects, in particular near DR21 C¹, delineate the dense ridge of gas and dust in both wavelength bands. The map of $8 \mu\text{m}$ emission is dominated by many streamers of emission, which appear to connect to the ridge, and diffuse emission. This is mainly emission from Polycyclic Aromatic Hydrocarbons (PAHs), which are the prime absorbers of far-ultraviolet (FUV) radiation from the surrounding and embedded massive stars. Bright

¹ We will use the ISO/LWS position names throughout the paper (cf. Table 2).

$8 \mu\text{m}$ emission therefore highlights the photon-dominated surface regions of molecular clouds (PDRs). The molecular outflow along DR21 E, C, and W is well traced by $4.5 \mu\text{m}$ emission, mainly due to strong lines of shock excited or UV-pumped $\text{H}_2 \nu=0-0 \text{ S}(9)$. No ro-vibrational lines of $\text{CO } \nu=1-0$ and only weak Br α have been found in this band (Smith et al. 2006).

3.1. Line-integrated maps

3.1.1. CO and ^{13}CO maps

Figure 1 (contours on right panel) and Fig. 2 show the KOSMA maps of integrated emission of the rotational transitions CO 3–2, 4–3, and 7–6 which preferentially trace the warm and dense molecular gas around DR21 and along the north-south orientated molecular ridge. Apart from the ridge, the most obvious feature seen in the maps is the line emission along the outflow axis oriented NE-SW. The main axis of the strong 3–2, 4–3 and 7–6 emission (FWHM $\sim 230''$, resp. $\sim 1.9 \text{ pc}$ ($D/1.7 \text{ kpc}$)) is tilted by 20° with respect to the right ascension axis and correlates nicely with extended H_2 emission lobes (Garden et al. 1986). The red-shifted CO line-wing emis-

Table 3. Line integrated intensities in [K km s^{-1}] (upper table) and in [$10^{-7} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$] (lower table) of the low- and mid- J CO lines observed (including the ^{13}CO and C^{18}O isotopomers) with KOSMA and FCRAO, and CS 7–6. In order to prevent the $+9 \text{ km s}^{-1}$ velocity component from adding to the integrated intensity, the listed intensities were integrated only over the -3 km s^{-1} component. As the ^{12}CO lines merge with the second velocity component, these spectra were fitted using a 2-component gaussian with the $+9 \text{ km s}^{-1}$ velocity component masked out. All line intensities are reported for $80''$ angular resolution (higher angular resolution data have been smoothed).

Position	CO	CO	CO	CO	^{13}CO	^{13}CO	^{13}CO	C^{18}O	[C I]	[C I]	CS
Units in [K km s^{-1}]	3–2	4–3	6–5	7–6	1–0	3–2	6–5	1–0	1–0	2–1	7–6
DR 21 E	348.7	283.4	247.2	265.7	22.38	69.4	43.1	1.66	21.9	28.2	3.0
DR 21 C	440.0	462.6	434.4	477.5	70.48	112.2	80.9	8.26	46.6	58.4	13.8
DR 21 W	230.5	221.9	177.7	223.1	41.02	47.6	33.9	3.67	26.9	28.5	2.9
DR 21 (OH)	283.2	197.4	162.0	153.2	67.92	91.5	24.9	9.03	39.5	30.2	13.6
DR 21 FIR1	223.8	193.2	130.2	107.3	59.49	61.9	26.0	9.16	33.4	33.0	3.9
Position	CO	CO	CO	CO	^{13}CO	^{13}CO	^{13}CO	C^{18}O	[C I]	[C I]	CS
in [$10^{-7} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$]	3–2	4–3	6–5	7–6	1–0	3–2	6–5	1–0	1–0	2–1	7–6
DR 21 E	147.8	284.6	837.6	1429.1	0.307	25.7	127.6	0.023	26.8	153.4	1.3
DR 21 C ^a	186.5	464.6	1471.9	2568.6	0.967	41.5	239.8	0.112	56.9	317.3	5.7
DR 21 W	97.7	222.9	602.1	1200.2	0.563	17.6	100.3	0.050	32.5	154.7	1.2
DR 21 (OH)	120.0	198.3	548.9	824.1	0.932	33.9	73.8	0.122	48.3	164.7	5.6
DR 21 FIR1	94.8	194.0	441.2	577.0	0.816	22.9	77.0	0.124	40.8	179.5	1.6

^a Boreiko & Betz (1991) give CO (^{13}CO) 9–8 intensities of $2.3 \pm 0.08 \cdot 10^{-4}$ ($0.45 \pm 0.1 \cdot 10^{-4}$) $\text{erg s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$ for a $80''$ ($83''$) beam and a telescope coupling efficiency of 0.6.




Fig. 6. Cut along the DR21 outflow axis (*top*) and along the ridge (*bottom*) centered on DR21 C in CO 3–2, 4–3, and 7–6 integrated emission. The $80''$ resolution is indicated as horizontal line. The dashed vertical lines indicate the positions of the ISO/LWS observations.

emission is stronger toward the west side, whereas the blue-shifted emission is more pronounced to the east indicating that this side is closer to the observer.

Morphologically, the CO 3–2 map shows a very similar shape in comparison to the CO 4–3 data. Detailed differences can be better seen if the maps are smoothed to the same resolution. In Fig. 6, we present two cuts along the outflow and along the main ridge from these data. At DR21, all CO line integrated intensities are very similar. Toward position DR21 E, the emission of CO 3–2 and 4–3

stays at a high level while the 7–6 emission drops. Toward the west, all three line intensities decline in a similar fashion. Away from DR21, along the N-S cut, the CO 3–2 emission is dominant, whereas CO 4–3 and CO 7–6 are sequentially weaker. Strong CO 7–6 emission is limited to the DR21 core. DR21 (OH) is located $180''$ north of DR21 and clearly separated from the DR21 region by a gap in the ridge. DR21 (OH) is bright in the lower J transitions, but appears weaker in the $J=7-6$ transition.

^{13}CO 3–2 (contours on left panel of Fig. 1) was selected to map the column density in the region. This transition traces the molecular ridge showing two maxima of emission at DR21 C and DR21 (OH). The total mass of the ridge is $3.5 \times 10^4 M_{\odot} (\text{D}/1.7 \text{ kpc})^2$ (SBS2006). ^{13}CO weakly follows the outflow from DR21 E to W. Following the molecular ridge further to the north, an elongated emission band roughly coincides in position with three far-infrared sources. The southernmost source, DR21 FIR1, was observed by ISO/LWS. Marston et al. (2004) find an extremely red object (ERO 3) here, illuminating a shell-like halo in the $4.5 \mu\text{m}$ band. The other two, FIR2 and FIR3, coincide with MSX point sources. As the FIR sources, this northern region is slightly tilted to the east, indicating flux contribution to mainly low- J CO emission from all three embedded far-IR sources. ^{13}CO reaches up to FIR3 which marks the edge of the dense ridge to the north.

3.1.2. [C I] maps

C I 1–0 and 2–1 appear in good agreement with the shape of the molecular ridge as traced by ^{13}CO . Even though the emission around DR21 in [C I] (see Fig. 3) appears more centrally peaked than in the CO maps, the main axis between positions E and W is clearly visible. The molecular ridge is seen to extend from the south east through DR21 and through DR21 (OH) 3' northwards. DR21 (OH) is visible as elongated, less compact region linking-up to the far-infrared sources further north. The DR21 (OH) region appears to harbour multiple sources as there is no central emission peak. The [C I] 1–0 map reveals a southern peak and extended weaker emission in the north. This substructure may not be significant because the variation is only about 2σ . Chandler et al. (1993) identified five compact sources in the DR21 (OH) region (indicated in Fig. 1) using 1.3-mm thermal dust emission at $11''$ resolution. C I 2–1 is, compared to 1–0, relatively weak toward DR21 (OH) and suggests a lower temperature than for DR21 C (see below). North of DR21 (OH), the carbon emission stretches over the three far-IR spots FIR1, FIR2, and FIR3. Here, the upper atomic carbon line is in good agreement with the overall shape of optically thin low- J CO isotopomers, i.e., the C^{18}O 2–1 map by Wilson & Mauersberger (1990), and to some extent our ^{13}CO 3–2 map. A more distinct morphology analysis is not possible due to the remaining noise and some self-chopping residuals.

3.1.3. CS 7–6 map

The map in Fig. 4 reveals two condensations at DR21 and around DR21 (OH). The southern peak is centered on the H II-region and follows the direction of the outflow to the west. The FWHM is $133'' \times 102''$. The northern complex (FWHM $108'' \times 156''$) is extended and may contain several unresolved cores (see above). The emission extends up to the FIR1 position, but does not cover all three FIR spots. The lines of CS 7–6 and C^{18}O 1–0 are the only lines in our sample that show brighter emission toward DR21 (OH) than toward the southern molecular cloud/H II-region DR21. For CS, this has also been observed by Shirley et al. (2003).

3.2. Line profiles

In the following, we concentrate on the 5 ISO/LWS positions (as indicated in Fig. 1). In Fig. 5, we present spectra of ^{12}CO , ^{13}CO , and C^{18}O towards these positions together with spectra of C I, and CS 7–6 between -30 to 30 km s^{-1} . We present all observations at a common resolution of $80''$. In Table 3, we give the corresponding line fluxes.

The ^{12}CO emission from the DR21/H II-region shows broad wing emission due to the outflowing molecular gas (with line widths of more than 15 km s^{-1} and peak temperatures well above 20 K). A self-absorption dip at the

velocity of -3 km s^{-1} is most pronounced in mid- J CO lines at DR21 C where the peak intensity drops by a factor of two at the line center. Toward DR21 (OH), CO shows blue-skewed, double-peaked lines with self-absorption. In both cases, the velocity of the line reversal nicely matches the corresponding emission in ^{13}CO and C I, and is likely caused by the same cold material in the dust lane.

With widths of 4 to 6 km s^{-1} , the line shapes of ^{13}CO are similar to the [C I] profiles. The ^{13}CO 6–5 emission ($E_u/k_B = 79.3 \text{ K}$) at all five positions indicates the presence of warm and dense gas. At DR21 C, ^{13}CO 6–5 shows broad symmetric wing emission that is underlying a narrow component. The velocity extent of the wings is similar to that in mid- J CO lines. A better signal-to-noise ratio would allow a more accurate estimate. Only weak line wings are visible in the lower ^{13}CO 3–2 line since it traces primarily the quiescent gas.

DR21 is associated with molecular gas centered at a velocity of -3 km s^{-1} . In addition to this, weaker emission between ~ 7 and 14 km s^{-1} is associated with the W75N complex approximately 11 pc (D/1.7 kpc) to the north-west. Some authors (e.g., Dickel et al. 1978) conclude that both clouds are actually interacting. SBS2006 shows channel maps revealing a link between both clouds. Diffuse emission in the ^{13}CO $J = 1-0$ and both [C I] transitions, and absorption in the red wing of the low- J CO lines (up to $J = 4-3$) suggest that this material is located in front of the DR21 complex. We do not further discuss this contribution here and refer to SBS2006.

3.3. ISO archive data

Figure 7 shows the ISO/LWS continuum fluxes together with spectrally unresolved lines of [C II], [O I], [O III], [N II], and rotational high- J CO transitions. The following Sections discuss the dust and line emission at the individual positions.

3.3.1. Grey body fit to the continuum

In all five cases, the far-infrared dust-continuum flux – with strong emission lines masked out – was χ^2 -fitted with a single isothermal grey-body model $F(T_{\text{dust}}, \tau_{\text{dust}})_\lambda = \Omega B_\lambda(T_{\text{dust}})(1 - e^{-\tau_{\text{dust}}})$ with fixed solid angle ($\Omega = 1.5 \cdot 10^{-7} \text{ sr}$, assuming that the emission fills the $80''$ aperture) and a dust opacity $\tau_{\text{dust}} \sim \tau_v \lambda^{-\beta}$ with $\beta = 1.5$. Spectral indices are typically found between 1.0 and 2.0 (see Goldsmith et al. (1997) for a discussion of observations and grain models). Although the spectral index is known to vary with the cloud type and optical depth, we assume a fixed β for all 5 positions (but see discussion in Sect. 4.2.3). A fixed β of 1.5 is compatible with grain models of Preibisch et al. (1993), Pollack et al. (1994), and observations, e.g., by Walker et al. (1990). The total infrared dust continuum flux and bolometric luminosity per position is derived by integrating the fit-model F_λ . The integration interval ($1 - 1000 \mu\text{m}$) covers the expected in-

Table 4. The ISO/LWS high- J CO and atomic FIR lines at $80''$ resolution. The transformation between flux densities and surface brightnesses was done using the beam sizes given in Table 5.10 of the ISO-LWS handbook v. 2.1. The beam solid angle varies between 8×10^{-8} sr and 1.4×10^{-7} sr. Because the lines are unresolved, the line widths were held to the instrumental line width: 0.29 or $0.60 \mu\text{m}$ for detectors 0 – 4 and 5 – 9, respectively. Correction factors between 0.37 to 0.85 for extended sources have been applied. We assume a 30% absolute photometry uncertainty for LWS. For DR21 (OH), numbers in parentheses are more uncertain (S/N between 1 – 3) due to a poorer frequency resolution. At DR21 FIR1, no detection of CO was possible, so we give 3σ upper limits. [N II] was clearly found only toward the outflow positions E and W. At the other positions we give 3σ upper limits.

Position in [$10^{-4} \text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]	CO 14–13	CO 15–14	CO 16–15	CO 17–16	[C II] 158 μm	[O I] 63 μm	[O I] 145 μm	[O III] 52 μm	[O III] 88 μm	[N II] 122 μm
DR 21 E	0.25	0.72	0.51	0.33	8.2	48.2	2.9	3.53	3.08	0.435
DR 21 C	1.75	2.46	0.99	0.63	8.1	49.4	6.15	11.5	4.64	< 0.75
DR 21 W	0.44	0.69	0.42	0.29	4.7	10.1	1.136	1.78	2.52	0.306
DR 21 (OH)	(0.45)	(0.99)	(0.40)	(0.50)	2.3	2.94	0.845	(1.27)	1.56	< 2.7
DR 21 FIR1	< 0.28	< 0.3	< 0.28	< 0.25	3.6	1.78	0.45	0.54	1.66	< 0.6

Table 5. Results of dust continuum single-component fit. β was fixed at 1.5, and Ω is $1.5 \cdot 10^{-7}$ sr. The extinction and the opacity are linked through $A_v = 2.5\tau_v \log_{10} e \approx 1.085\tau_v$. The total infrared flux is calculated from $I_{\text{IR}} = \int F_\lambda d\lambda$. The luminosity L and mass M are given for 1.7 kpc distance.

Position	τ_v	$A_v(\text{ISO})$	N(H ₂) [10^{22}cm^{-2}]	Mass [$M_\odot(\text{D}/1.7 \text{kpc})^2$]	T _{dust} [K]	I_{IR} [$\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$]	L [$10^3 L_\odot(\text{D}/1.7 \text{kpc})^2$]
DR 21 E	20	21.2 ± 1.9	2.1	148	41.6 ± 0.6	0.55 ± 0.05	7
DR 21 C	112	121 ± 11	12.1	850	42.5 ± 0.7	3.4 ± 0.3	45
DR 21 W	34	36.9 ± 3.5	3.69	259	30.7 ± 0.4	0.18 ± 0.02	2
DR 21 (OH)	238	259 ± 43	25.8	1814	31.2 ± 0.8	1.3 ± 0.2	17
DR 21 FIR1	45	48.4 ± 4.3	4.84	340	33.3 ± 0.5	0.37 ± 0.03	5

tensity maxima between $50\text{--}80 \mu\text{m}$ and corresponds to the range used, e.g., by Calzetti et al. (2000). Errors of the fit parameters were estimated as described in the following: we assume (i) the optical depths τ_{dust} and total fluxes are accurate to about $\sim 10\%$, mainly due to the remaining calibration artefacts within the LW1–5 bands. (ii) the temperature fit error is less than 2% but the absolute error also depends on the quality of SW1 detector data (leftmost data points in Figure 7).

We find extinctions between $20^m \leq A_v \leq 260^m$ and temperatures between 31–43 K (cf. Table 5). In contrast to the extinction, the variation in dust temperature is small. Considering that we use a single-temperature model, these temperatures are probably biased by the warmer dust. Evidence of cold dust peaking above $100 \mu\text{m}$ is visible at DR21 W, E, (OH) and FIR1 in Fig. 7 but disentangling this component would require longer wavelength data and the contribution to the integrated grey-body flux is relatively small. The cold dust is studied in detail by, e.g., Chandler et al. (1993) and Motte et al. (2005). We derive the molecular hydrogen column density from the conversion $N(\text{H}_2)/A_v \approx 0.94 \cdot 10^{21} \text{cm}^{-2} \text{mag}^{-1}$ (Bohlin et al. 1978, cf. Table 5) and if placed at the distance of 1.7 kpc, the molecular gas masses vary between $150\text{--}1800 M_\odot$.

The peak of the beam-averaged dust temperature is ~ 43 K at DR21 C. With $A_v > 100^m$, the strongest continuum sources are DR21 C and DR21 (OH). Due to the high

temperature towards DR21 C, the flux density is higher at short wavelengths, but drops below the flux density of DR21 (OH) above $\sim 200 \mu\text{m}$. For DR21 (OH), we estimate a higher mass than for the DR21 E–W complex (1814 , resp. $1257 M_\odot$). Chandler et al. (1993) assumed temperatures between 25 K and 40 K and derived a slightly lower total mass of $1650 M_\odot(\text{D}/1.7 \text{kpc})^2$ for the sources at DR21 (OH).

3.3.2. ISO/LWS far-IR lines: Carbon Monoxide and Ionised Carbon

The LWS band covers many important far-IR cooling lines and tracers of the photon dominated regime of molecular clouds. Table 4 gives the integrated line intensities of [C II] ($158 \mu\text{m}$), high- J CO, and [O I] (63 and $145 \mu\text{m}$) derived from the ISO observations quoted above. In this paragraph, we focus on the carbon bearing species needed for Sect. 4 and postpone the discussion of the other lines to the end of Sect. 4.

At DR21, highly excited CO emission is visible at all three positions, but most prominently around the H II-region at DR21 C. The center position with more than $2.4 \cdot 10^{-4} \text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$ in mid- and high- J CO is also the brightest continuum source in the mapped region. The emission stems from large columns of warm quiescent gas




Fig. 7. ISO/LWS Spectral Energy Distribution (SED) of the DR 21 region (*upper panel*), and the DR21 (OH) and DR21 FIR1 region (*lower panel*). Curves indicate the best greybody fit corresponding to the values given in Table 5.

that is heated from an embedded object. Toward the East and West positions, where vibrationally excited H_2 outlines the molecular outflow (Garden et al. 1986), the FIR continuum is somewhat weaker and more importantly, the continuum flux does not correlate with the unusually strong FIR line emission. This gas is visible in the line wings of mid- J CO lines (Fig. 5) and in bright H_2 line emission at $4.5 \mu\text{m}$ in the IRAC map (Fig. 1). Lane et al. (1990) conclude that neither a single C- nor a single J-shock can account for the observed far-IR line emission alone and additional heating through FUV radiation is required. Although DR21 (OH) has the second strongest continuum, the sources associated with this position are surprisingly inactive in far-IR line emission. We find four high- J CO line detections, but due to a lower scanning resolution, the values given in Table 4 have higher uncertainty. The far-IR source FIR1 is the faintest with respect to the line fluxes, although the continuum is stronger than towards DR21 W. High- J CO was not detected at FIR1, so we give upper limits in the table.

[C II] is detected at all five positions. The emission peaks at DR21 E and toward the West, the flux decreases by almost 50%. The low [C II] flux of $2.3 \cdot 10^{-4} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$ at DR21 (OH) may indicate a lack of ionised carbon or that the temperature is too low for excitation in the upper energy level ($E_u/k_B = 91 \text{ K}$ above ground). At DR21 FIR1, [C II] is again slightly enhanced compared to DR21 (OH).

4. The physical structure of DR 21

In this section we analyse the line properties of the observed carbon species from ISO/LWS data and combine these results with the KOSMA observations. We start with an LTE approach and then model the line emission with full radiative transfer.

4.1. Comparison of CO, C, and C^+

We estimate the composition of the molecular gas with respect to the carbon abundance in the main gas-phase species CO, C, and C^+ . This is done by assuming LTE and optically thin line emission. We therefore use the less abundant isotopomers ^{13}CO and C^{18}O . We list the results for excitation temperature, column density and abundance in Table 6.

4.1.1. Atomic Carbon

The C I excitation temperature needed to produce the observed line ratios can be estimated since both C I lines are likely to be optically thin (cf. line profiles in Fig. 5). The line temperature ratio $R_{\text{C I}}$ falls between 0.8 at DR21 (OH) and 1.3 around and east of DR21. The temperature is determined from this ratio using $T_{\text{ex}} = 38.8 \text{ K} / \ln(2.11/R_{\text{C I}})$ (Eq. (C.5)). The range (38 K at (OH) to 79 K at E) indicates the origin of the atomic carbon lines in a warm environment. Zmuidzinas et al. (1988) derived a temperature of 32 K from the two C I lines at DR21 (OH) which is close to our value at this position. Our results for T_{ex} are lower compared to other Galactic high mass star forming regions (e.g., W3 Main and S 106 with typically $T_{\text{ex}} \geq 110 \text{ K}$, Kramer et al. 2004; Schneider et al. 2003), but substantially higher compared to the average temperature in the inner Galactic disk of $\sim 20 \text{ K}$ (Fixsen et al. 1999). The beam averaged column density, $3.2 - 6.8 \cdot 10^{17} \text{ cm}^{-2}$ (Eq. (C.6)), shows variations by a factor of 2 or less, peaking at the position DR21 C and around (OH), i.e., towards the densest condensations along the ridge. This is remarkable since the H_2 column density derived from the dust was found to vary by a factor of 12. The column density for DR21 (OH) of $6.3 \cdot 10^{17} \text{ cm}^{-2}$ as determined by Zmuidzinas et al. is only slightly higher than our result.

4.1.2. Carbon Monoxide

The amount of ^{13}CO is determined in two ways assuming again optically thin emission. Firstly, by using $T_{\text{ex}, ^{13}\text{CO}, \text{high}} = 79.4 \text{ K} / \ln(4/R_{^{13}\text{CO}, \text{high}})$ (Eq. (C.9)) with the ^{13}CO 6–5 to 3–2 integrated intensity line ratio $R_{^{13}\text{CO}, \text{high}}$ and by Eq. (C.10). This ratio is a sensitive temperature measure from $\sim 20 \text{ K}$ up to a few 100 K. The excitation temperatures vary between 30 and 46 K and represent upper limits because of possible self-absorption due to a high optical thickness in the lower transition. The temperatures agree with values from Wilson & Mauersberger (1990) who find 20 – 65 K from NH_3 emission, and are

Table 6. C I and ^{13}CO line ratio R (in K km s^{-1} ; values taken from Table 3), excitation temperature (with lower and upper 1σ limits), and C I, ^{13}CO (with relative upper and lower errors), and C^{18}O column densities (bracketed by T_{ex} of ^{13}CO). The C II column density was estimated using a method described in Sect. 4.1. All abundances X are relative to the H_2 column densities (first line; taken from Col. 4 of Table 5). The last row lists the relative abundance of carbon in the three major gas phase species C II, C I, and CO. The CO column density is derived from the lower C^{18}O column density assuming a relative $[\text{CO}]:[\text{C}^{18}\text{O}]$ abundance of 490.

Line/Lineratio	Source Property	DR21 E	DR21 C	DR21 W	DR21 (OH)	DR21 FIR1
$\text{H}_{2,\text{dust}}$	$\text{N}(\text{H}_2)$ [10^{22} cm^{-2}]	2.1	12.1	3.69	25.8	4.84
C I	$R_{\text{C I}}$ (C I 2–1/1–0)	1.29 ± 0.36	1.25 ± 0.35	1.06 ± 0.30	0.77 ± 0.22	0.99 ± 0.28
	$T_{\text{ex},[\text{C I}]}$ [K]	79_{47}^{157}	74_{50}^{140}	56_{41}^{88}	38_{31}^{51}	51_{39}^{76}
	$\text{N}(\text{C I})$ [10^{17} cm^{-2}]	$3.2_{-0.2}^{+0.3}$	$6.8_{-0.4}^{+0.6}$	$3.8_{-0.2}^{+0.3}$	$5.2_{-0.1}^{+0.2}$	$4.6_{-0.2}^{+0.3}$
	$\text{X}(\text{C I})$ [10^{-5}]	1.5	0.56	1.03	0.20	0.10
^{13}CO 6–5/3–2	$R_{13\text{CO},\text{high}}$ (^{13}CO 6–5/3–2)	0.62 ± 0.26	0.72 ± 0.31	0.71 ± 0.30	0.27 ± 0.12	0.42 ± 0.18
	$T_{\text{ex},^{13}\text{CO},\text{high}}$ [K]	43_{36}^{53}	46_{38}^{58}	46_{38}^{58}	30_{26}^{34}	35_{30}^{41}
	$\text{N}(^{13}\text{CO}$ 6–5) [10^{15} cm^{-2}]	$33.0_{+11.1}^{-9.0}$	$53.8_{+20.4}^{-12.8}$	$22.6_{+8.5}^{-5.4}$	$39.6_{+21.3}^{-10.7}$	$27.9_{+13.4}^{-7.2}$
^{13}CO 3–2/1–0	$R_{13\text{CO},\text{low}}$ (^{13}CO 3–2/1–0)	3.10 ± 1.30	1.59 ± 0.67	1.16 ± 0.49	1.35 ± 0.57	1.04 ± 0.44
	$T_{\text{ex},^{13}\text{CO},\text{low}}$ [K]	25_{16}^{37}	15_{12}^{19}	13_{10}^{16}	14_{11}^{17}	12_{10}^{15}
	$\text{N}(^{13}\text{CO}$ 1–0) [10^{15} cm^{-2}]	$33.4_{-8.9}^{+12.3}$	$75.6_{-8.6}^{+12.2}$	$40.0_{-4.6}^{+5.1}$	$69.1_{-8.0}^{+8.6}$	$56.5_{-4.3}^{+7.2}$
C^{18}O 1–0	$\text{N}(\text{C}^{18}\text{O})$ [10^{15} cm^{-2}]	2.4 – 3.8	8.7 – 20.0	3.5 – 8.9	9.0 – 15.1	8.5 – 17.7
	$\text{X}(\text{C}^{18}\text{O})$ [10^{-7}]	1.2 – 1.8	0.72 – 1.7	0.95 – 2.4	0.35 – 0.59	1.7 – 3.7
C II	$\text{N}([\text{C II}])$ [10^{17} cm^{-2}]	5.25	5.18	3.00	1.47	2.30
	$\text{X}(\text{C II})$ [10^{-5}]	2.5	0.43	0.81	0.057	0.48
	$[\text{C}^+]:[\text{C}^0]:[\text{CO}]$ [%]	26:16:58	10:12:78	16:12:72	03:10:87	05:09:85

always below the C I temperatures from above indicating an origin from different gas. The ^{13}CO column density, $23 - 54 \cdot 10^{15} \text{ cm}^{-3}$, varies by a factor of 2 as the C I column density. The same scatter is found if we estimate the column density from ^{13}CO 1–0 rather than ^{13}CO 6–5 (Eq. (C.11)) and derive temperatures by Eq. (C.8). Those column densities tend to be higher, between $33 - 76 \cdot 10^{15} \text{ cm}^{-3}$, indicating that a larger amount of gas traced by ^{13}CO is cold.

Considering the optically thin emission of C^{18}O , we estimate the total molecular gas column density from Eq. (C.12) using both ^{13}CO T_{ex} estimates as limits. We find high C^{18}O column densities of about $1 - 2 \cdot 10^{16} \text{ cm}^{-2}$ along the ridge and $2 - 4 \cdot 10^{15} \text{ cm}^{-2}$ toward E and W.

4.1.3. Ionised Carbon

In order to get a rough estimate of the C^+ column density from the ISO data, we assumed $\text{N}(\text{C II}) = 6.4 \cdot 10^{20} \times I([\text{C II}])$ [cm^{-2}] ($\text{erg s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$) $^{-1}$ (Crawford et al. 1985), which is a good approximation for an excitation temperature of $> 91 \text{ K}$, and a density $> n_{\text{cr}} = 5 \cdot 10^3 \text{ cm}^{-3}$. The peak column density derived this way, $\text{N}(\text{C II}) \sim 5.2 \cdot 10^{17} \text{ cm}^{-3}$, is observed towards DR21 C and E, while only about 28 % of the peak value is found towards DR21 (OH).

4.1.4. Abundances

The comparison between C^{18}O and ^{13}CO column density (for the lower T_{ex} limit in Table 6) shows that the abundance ratio $[\text{C}^{18}\text{O}] : [\text{C}^{18}\text{O}] = 6.6 - 11.4$ is typically near the canonical local abundance ratio $[490] : [65] = 7.5$ (Langer & Penzias 1990).

The atomic carbon abundances $[\text{C I}]/[\text{H}_2]$ fall in the range of $0.1 - 1.5 \cdot 10^{-5}$. The high values are only observed at the E and W positions. In the cooler northern positions the abundance is reduced by a factor up to 10. This is in contrast to the values of $1 - 2 \cdot 10^{-5}$ by Keene et al. (1997) found in cold dark clouds.

Except for DR21 FIR1, there is a good correlation between $\text{X}(\text{C I})$ and $\text{X}(\text{C}^{18}\text{O})$. Toward DR21 (OH), the C^{18}O and $[\text{C I}]$ abundance are systematically lower than toward DR21 by factors 2 – 3. Although, the position with the lowest $[\text{C I}]$ abundance (10^{-6}), FIR1, has the highest C^{18}O abundance (up to $3.7 \cdot 10^{-7}$).

Mookerjea et al. (2006) present a compilation of the C/CO abundance ratios versus H_2 column densities for Galactic star forming regions and diffuse clouds. The ratio at the positions DR21 E, W, and FIR1, 0.07–0.17, agrees well with the relation $\log[\text{N}(\text{C})/\text{N}(\text{CO})] = -0.94\text{N}(\text{H}_2) + 19.9$ given there for Cepheus B. DR21 and DR21 (OH) deviate from this relation but still are within the observed scatter. We therefore underline their conclusion that C I

is no straightforward tracer of total H_2 column densities and total masses.

The amount of carbon in the carbon bearing species C^+ , C^0 , and CO can be compared using their relative abundances (last row of Table 6). The neutral atomic carbon relative abundance is rather homogeneous at about 9–16% of the total carbon gas content at all five positions. The rest is shared between CO and C II . Ionised carbon has the highest abundance toward DR21 E (26%). Here, the gas is likely exposed to a high UV radiation field and carbon species are photodissociated to C I and ionised further to C II . High fractional CO abundances (>85%), such as for DR21 (OH) and FIR1 were also reported for S106 (>86%, Schneider et al. 2003) and W3 Main (>60%, Kramer et al. 2004). The ratios in other sources are, e.g., 37:07:56 for IC 63 (Jansen et al. 1996), NGC2024 with 40:10:50 (Jaffe & Plume 1995), or η Carina with 68:15:16 (Brooks et al. 2003).

4.2. Radiative transfer modeling

In a self-consistent approach, the temperature, column and number density, total molecular mass, volume filling and mass composition were determined using a 1-D radiative transfer code (SIMLINE, Ossenkopf et al. 2001). In order to describe these physical conditions, we created heterogeneous models with internal clumping and turbulence in a spherically symmetric geometry for each of the 5 observed regions. The physical parameters of a shell are defined by their values at the inner radius and the radial dependence is described by a power-law.

A model with two nested shells provided in most cases reasonable results: (i) a hot and dense molecular *interface zone* wrapped around a central H II -region, (ii) enclosed by a massive cold cloud – the *envelope region*. While in the inner component primarily the mid- and high- J CO emission lines are excited, the envelope is responsible for self-absorption and the low lying lines of ^{13}CO and C^{18}O . We set the abundance of species with respect to molecular hydrogen to standard values (see Table 7). The source size is a crucial parameter that is not well confined by the line profiles or the map we obtained. A small cloud model can easily produce results similar to those for a bigger cloud with a lower volume filling. We therefore approximated the radii using high resolution line and continuum observations (Wilson & Mauersberger 1990; Lane et al. 1990; Jaffe et al. 1989; Chandler et al. 1993; Vallée & Fiege 2006). The inner cut-off was assumed to be 0.04 pc ($< 10^4$ AU), except for DR21 C where we estimated ~ 0.12 pc from the 14.7 GHz continuum (Roelfsema et al. 1989). The H II -region is considered to be free of molecular gas and we neglect the continuum contribution by ionised gas. Gradients within the shells in any of the parameters are initially set to zero and are adjusted during the fit if required.

We used the following *recipe* to find appropriate model parameters: (i) The linewidth is approximated by adjust-

ing the turbulent velocity field until both the low- J and mid- J line wings match the observed profile. As line emission from the inner warm gas is shielded by the colder envelope gas closer to the surface, the width of the CO absorption dip in the line center constrains the turbulent velocity dispersion of this envelope component. (ii) The H_2 -density is defined as a local clump density parameter and varied between 10^4 and 10^7 cm^{-3} independently for both shells. The high densities are a pre-requisite for high- J CO excitation. (iii) In the optical thin line wings we see predominantly material from the inner shell where the simulated macroturbulent motion leads to the higher line width which is only partly shielded by the outer shell. This allows us to fit the temperatures of both the inner and the outer shell from the line shape. (iv) The model allows to differentiate between radial and turbulent gas motion. Where lines of high opacity toward the line center show a blue-skewed profile, we introduced radial infall as a plausible explanation. If possible, we tried to keep the radial component close to zero.

The full line profiles of CO , ^{13}CO , and C^{18}O from KOSMA and FCRAO are taken into account in a reduced χ^2 fit in ~ 1900 degrees of freedom and using 20 – 28 free parameters. The second velocity component is masked out by a window centered on $9 \pm 3 \text{ km s}^{-1}$. For the ISO lines, we fit the integrated flux of three lines. $\text{CO } 14-13$ is not considered because no consistent fit was possible in conjunction with the other CO lines. In total, we include 11 lines weighted by the σ^2 determined from a baseline region not including any lines. The spectra do not contain any obvious baseline ripples above the system noise.

4.2.1. Line Modelling: CO , ^{13}CO , and C^{18}O

We independently optimized the model parameters for each position. Fig. 5 shows the derived line profiles. At all five positions, we obtain solutions of better than $\chi_0^2 = 4.1$ which we consider – given the number of fitted lines – a very good result, and definitely much better than for a single-component model.

Within the interface-zone, a temperature range from 80 K (at DR21 FIR1) to 148 K (at DR21 C) for gas masses of 15 – 60 M_\odot is found. This temperature is much higher than the typical LTE excitation temperature of the mid- J CO line peak. The fit converges toward solutions with a high density of about 10^6 cm^{-3} and with a volume filling of a few percent, indicating a non-homogeneous distribution. The warm gas dominates the emission in the line wings and therefore is explained by the steep turbulent velocity gradient with velocities up to 30 km s^{-1} . Because the envelope is much colder and more massive (cf. Sect. 4.2.3), the average temperature in the regions DR21 C with 36 K and DR21 (OH) with 31 K is rather low compared to the E, W, and FIR1 position. There, the gas temperature rises up to 60 K but the gas masses are only a few hundred solar masses. Besides the mass fraction, the steepness of the gradient in excitation condition and line width between

Table 7. The physical parameters for the observed ISO/LWS positions. The radii listed in column 2 give the inner and outer edge of each shell. Column 3 to 7 lists the temperature, clump-averaged number density of H₂, volume filling, column density of H₂, and the gas mass. For these columns, the first entry is the mass-averaged value for the whole cloud, values in parentheses show averages within each shell. The last column lists the reduced χ^2 fit result for 11 lines (6 KOSMA, 2 FCRAO & 3 ISO).

Position	Radii [pc]	T _{gas} [K]	n(H ₂) [10 ⁴ cm ⁻³]	ϕ_v volume filling ^b	N(H ₂) [10 ²¹ cm ⁻²] ^a	Mass [M _⊙ (D/1.7 kpc) ²] ^c	χ_0^2
DR 21 E	0.04/0.2/0.3	62 (87/55)	92 (300/4.8)	0.04 (0.01/0.89)	46 (9/37)	243 (46/197)	2.2
DR 21 C	0.12/0.2/0.6	36 (148/32)	110 (40/110)	0.024 (0.08/0.02)	65 (2/63)	1379 (51/1328)	3.5
DR 21 W	0.04/0.1/0.3	54 (118/52)	12 (100/9)	0.59 (0.06/0.8)	88 (3/85)	489 (15/474)	2.8
DR 21 (OH)	0.04/0.2/0.7	31 (99/29)	161 (220/160)	0.01 (0.01/0.01)	49 (1/48)	1421 (34/1387)	3.9
DR 21 FIR1	0.04/0.2/0.5	41 (82/37)	16 (220/2)	0.1 (0.01/1)	47 (4/43)	700 (59/641)	4.1

^a Column densities are source averaged and not normalized to 80''.

^b The product of the volume filling and the abundance of the species is one of the free parameters. We fixed the abundance [X/H₂] of ¹²CO to $2 \cdot 10^{-4}$ (resp. $3.1 \cdot 10^{-6}$ for ¹³CO, and $4 \cdot 10^{-7}$ for C¹⁸O).

^c The molecular gas mass is derived from the cloud-averaged density. This density is the product of local H₂ density of the clumps and volume filling correction ϕ_v .

inner and outer component determines whether we see a distinct self-absorption (e.g., DR21 C) or a flat-top (e.g., DR21 W) line profile. Table 7 compiles a list of the derived physical parameters. A detailed discussion of the individual positions is given in App. A.

4.2.2. Line Modelling: [C I] and CS

As a consistency check, we computed the CS and C I profiles using the same model parameters as derived from CO. This implies a homogeneous distribution of CS and C I with CO, only affected by the abundances of the species.

For both species we also obtained excellent fit results. The resulting line profiles are shown in Fig. 5. CS 7–6 is only excited where the H₂ density and temperatures are sufficiently high. Hence, we find CS emission in the dense regions DR21 C and (OH) only. The models fit best to the observations for an abundance of X(CS)= $4 \cdot 10^{-9}$ (Hatchell et al. 1998), whereas the range $1 \cdot 10^{-10}$ to $2.3 \cdot 10^{-10}$ given by Shirley et al. (2003) is clearly too low for this region.

Instead of a carbon abundance variation across the positions (as found in Sect. 4.1), an abundance X(C I) of $1 \cdot 10^{-5}$ nicely fits our [C I] observations. The C I excitation temperatures as derived in Sect. 4.1.1 fall in the range covered by the two-component radiative transfer model.

4.2.3. Masses and Volume Filling

The volume filling, ϕ_v , is a model parameter capable of increasing the clump density without increasing the overall mass as well. The determined ϕ_v is usually low between 1% to 8% (Col. 5 in Table 7). At DR21 C, this is confirmed by our observations of the CS 7–6 transition which is sensitive to densities of several 10^6 cm⁻³: Given the high H₂ column density (up to $2 \cdot 10^{23}$ cm⁻²) this

translates to a column-to-volume density ratio of about 0.006 pc. If we assume that the visible extent of the DR21 source (~ 0.69 pc in CS, after deconvolving the beam) is similar to the size along the line-of-sight, a volume filling of ~ 0.01 is derived. A high degree of small scale spatial and velocity structure is also needed to explain the hyper-fine-structure anomalies observed in ammonia data toward DR21 (Stutzki 1985). Shirley et al. (2003), who studied CS 5–4 in a sample of high-mass star-forming regions, derived a volume filling with a median of 0.14. This is one order of magnitude higher than our results for the dense regions DR21 and DR21 (OH).

The total mass for all 5 positions is $4.2 \cdot 10^3 M_{\odot}$, and mostly shared between DR21 C and DR21 (OH) with $\sim 1.4 \cdot 10^3 M_{\odot}$ each. In direct comparison to the masses derived from dust observation in Sect. 3.3.1, we find that our model gives higher masses ($\times 1.6 - \times 1.8$), except for the DR21 (OH) model mass. To compare our mass estimate with masses from the literature, often derived for a different distance, we have scaled the mass in Table 7 with the square of the distance, basically assuming that the column density, computed from the observed lines, is independent from the distance. This approach is consistent with a local radiative transfer treatment, such as LTE or escape probability analyses, where the molecular excitation does not depend on the total gas mass. For the self-consistent radiative transfer code, this assumption is, however, not completely justified, but deviations from this simple scaling relation are to be expected. When comparing the values from Table 7 to results obtained for much larger distances our quadratically scaled mass estimates are therefore always expected to be too high and when comparing them to results for lower distances they should be systematically lower. Shirley et al. estimated virial masses of $561 M_{\odot}(D/1.7 \text{ kpc})$ for DR21 (resp. $714 M_{\odot}(D/1.7 \text{ kpc})$ for DR21 (OH)). These estimates are




Fig. 8. Cooling curves of the best-fit model line fluxes (marked by open symbols) of CO, ^{13}CO , and C^{18}O . Observed line intensities are indicated as filled symbols (see Table 3). At position DR21 C, we included two observations by Boreiko & Betz (1991) of CO and ^{13}CO $J = 9-8$. Since the ^{13}CO 6–5 map does not cover the positions DR21 (OH) and DR21 FIR1 on a fully sampled grid, we expect a larger error. The high- J CO emission at DR21 FIR1 are upper limits.

lower than our masses but were calculated over a much smaller radius of 0.17 pc ($D/1.7$ kpc) while our region covers up to 0.7 pc. Ossenkopf et al. (2001) estimated the DR21 mass using a radiative transfer analysis on CS data. They estimated $578 M_{\odot} (D/1.7 \text{ kpc})^2$ at a (cloud-averaged) density of $1.6 \times 10^3 \text{ cm}^{-3}$. Their lower density is partly explained by the larger outer radius, that was chosen to be 1.0 pc ($D/1.7$ kpc). It is probably also due to the larger physical cloud size at the assumed distance of $D = 3$ kpc which prevents the simple quadratic distance scaling of the mass in a self-consistent radiative-transfer treatment.

The total mass of DR21 (OH) corresponds to a cloud-averaged column density of $\sim 5 \cdot 10^{22} \text{ cm}^{-2}$. The value of $5.7 \cdot 10^{23} \text{ cm}^{-2}$ reported by Richardson et al. from $1100 \mu\text{m}$ data and our value given in Table 5 based on the far-IR dust-emission are higher by a factor of 5–10. A flatter spectral index β may explain why dust and line observation analyses result in different conclusions. Values between 1.0 (Vallée & Fiege 2006) and 2.0 (Chandler et al. 1993) were discussed for DR21 (OH) in the literature. If β is lower than the assumed 1.5, this will have a significant effect on the derived A_{ν} . For instance, with $\beta = 1.25$, the column density is reduced to about one fourth of our value at $\beta = 1.5$, then is consistent with the CO model results.

An alternative explanation for the discrepancy could be depletion as a result of freeze-out of CO onto grains. Some authors (e.g., Vallée & Fiege 2006, and references therein) use a depletion factor of 10 or more ($X(\text{CO}) = 2.5 \cdot 10^{-6}$ at DR21 (OH)) as appropriate for star-forming clouds. In Jørgensen et al. (2004), the abundance is reduced by a factor of 10 for the conditions $T < 30$ K and $n > 3 \cdot 10^4 \text{ cm}^{-2}$. The DR21 (OH) envelope and the resulting column density would comply with these conditions. However, the CO line data analysis does not allow us to draw conclusions toward depletion.

4.2.4. Problems of the fit

As noted before, a precise determination of the outer radii of the model clouds, and hence their mass, is quite uncertain and a small increase may already more than double

the mass. In the models of DR21 C and DR21 (OH), we found no reasonable solutions for radii much smaller than 0.6 pc matching the ^{13}CO and C^{18}O 1–0 observations. As possible indication of a systematic error in the column density we typically see that the observed peak temperature of the C^{18}O 1–0 line is weaker than predicted by the models (see Fig. 5) while the ^{13}CO 1–0 line is well reproduced. This can either be a result of an inaccurate modeling of density and temperature conditions, or our assumption about the fixed abundances for CO, ^{13}CO , or C^{18}O is incorrect.

The models overestimate the ^{13}CO 3–2 peak temperature. Except for DR21 E, this transition is optically thick ($\tau > 2$) and reflects the kinetic temperature of the envelope gas. As the CO 3–2 to 7–6 lines also become optically thick in the envelope, the brightness temperature at the self-absorbing line center of these mid- J lines is governed by the same material. The very deep (~ 5 K) absorption dip in CO 7–6 observed by Jaffe et al. (1989) indicates very cold foreground gas (< 20 K) which we cannot reproduce with only one component for the outer shell. A better match of the observed intensities may be possible by adding a third colder layer outside of the envelope.

Since the cold and warm components certainly are intermixed to some extent rather than separated in two nested shells, the focus during the modeling was put on a qualitative description of the different components and should not be seen as a geometric representation of the real cloud structure. We demonstrated, however, that the simplified assumption of a spherical cloud is adequate to estimate physical properties, such as mass, density, filling factors, and temperature.

4.3. Oxygen and Nitrogen

4.3.1. ISO/LWS far-IR lines: [O I], [O III], and [N II]

In addition to the high- J CO lines (see Sect. 3.3.2), ISO provided the integrated line intensities of [O I] (63 and $145 \mu\text{m}$) (Table 4). Their relative strength and cooling efficiency depends on the temperature and density regime: At low temperatures, rotational mm- and submm-lines of

CO dominate the cooling of the molecular and neutral gas. At temperatures above $T \sim 150$ K, pure rotational transitions of OH, H₂O, H₂, and the $63 \mu\text{m}$ [O I] line become more important. [O I] at $63 \mu\text{m}$ and the atomic and ionic fine-structure lines of [C II] and [C I] are diagnostic lines for PDRs. Due to the nearby H II-regions, the ionic lines [O III] and [N II] are also expected to be strong emitters and may serve as diagnostics to discriminate the fraction of [C II] emission coming from the neutral and ionised gas. For completeness, we included the [O III] (52 and $88 \mu\text{m}$) and [N II] ($122 \mu\text{m}$) lines in the table.

Since DR21 E, C & W line up along the outflow axis, we compare the emission in the outflow lobes with the central position: The strong [O I] emission at $63 \mu\text{m}$ decreases rapidly toward the west (cf., Lane et al. 1990). This line easily becomes optically thick and Poglitsch et al. (1996) indeed found a deeply absorbed line toward DR21. The $63 \mu\text{m}$ emission extends toward the E position at almost the same strength as toward the center. As the A_v at the same time drops down, the gas traced by O I must be a separate, necessarily warm component. [O I] at $145 \mu\text{m}$ is peaked at DR21 C, showing only a slight east-west asymmetry.

If we assume thermalized optically thin [O I] emission, we can derive the excitation temperature from $T_{\text{ex}} = 227.7 \text{ K} / \ln(2.83 / (12.22 \cdot R_{[\text{O I}]}))$ (Eq. (C.13)), where $R_{[\text{O I}]}$ is the ratio of [O I] $145/63 \mu\text{m}$ emission (cf. Table 4). The ratios of observed intensities, 0.06–0.12, result in temperatures between 170 to 370 K and are to be seen as strict upper limits since $R_{[\text{O I}]}$ decreases for optically thick [O I] $63 \mu\text{m}$ line emission. The [O I] line ratio at DR21 (OH) of $R_{[\text{O I}]} = 0.29$ requires a temperature just below 100 K. We detect the weakest emission of [O I] at DR21 FIR1. Errors in $R_{[\text{O I}]}$ become too large to estimate a temperature here.

As expected for the position of the H II-region, the ionised [O III] lines both are strong at DR21 C and decreases to the sides. Furthermore, the [O III] $88/52 \mu\text{m}$ line ratio is a good discriminant of the electron density in the range $10^2 - 10^5 \text{ cm}^{-3}$, while being insensitive to abundance and temperature variations (Rubin et al. 1994). At DR21 C, a ratio of about 0.4 indicates an electron density of $\langle N_e \rangle \sim 10^3 \text{ cm}^{-3}$. In the east, the ratio rises to 0.8 indicating an extension of the H II-region to this side. To the west, the ratio is above 1.4. An electron density below 10^2 cm^{-3} , consistent with the observed [O III] line ratio at DR21 (OH), indicates a low degree of ionisation. No H II-region has been identified in this area, e.g., in the VLA 6-cm continuum (Palmer et al. 2004). Thus, the finding of [O III] emission is a bit surprising. The [O III] ratio $> 3 \pm 1.3$ found at DR21 FIR1 is not covered by the Rubin et al. model but corresponds to a very low $\langle N_e \rangle$.

[N II] $122 \mu\text{m}$ was detected at the E and W positions only. At the position of the H II-region DR21 C, the 3σ noise limit still permits centrally peaked emission. The fraction of [C II] stemming from the H II-region can be estimated from the intensity ratio of [C II] to [N II]. The Galactic abundance ratio C/N in dense H II-regions leads to an expected ratio of $[\text{C II}]/[\text{N II}] = 1.1$

(Kramer et al. 2005, and references therein). With the observed $[\text{C II}]/[\text{N II}]$ ratio between 11–19 for DR21, less than 10 % of the [C II] emission are estimated to come from the ionised gas.

4.3.2. Line Modelling: [O I] and [C II]

A determination of the O I and C II abundances may be difficult. If we assume that their abundance is closely connected to the dense molecular gas, we can calculate and fit the emission of these two species by adjusting the abundance of the two-shell radiative transfer models (Sect. 4.2). At DR21 C, we find abundances of $6 \cdot 10^{-5}$ for O I and $2.5 \cdot 10^{-4}$ for C II. For both C II and the O I $63 \mu\text{m}$ line, the opacity toward the line center is about 9 and the line profile is deeply self-absorbed. The high energy above the ground level of O I leads to almost complete absorption of the center emission leaving signatures of warm gas only in the wings. The $145 \mu\text{m}$ transition ($\tau \sim 0.5$) is unaffected and is a better measure of the abundance. As C II has only one transition and also shows some absorption in the model profiles, the abundance may be less accurate. At DR21 (OH), the abundance of O I is $\sim 1.2 \cdot 10^{-5}$ and $9 \cdot 10^{-5}$ for C II. The abundances at all five observed positions vary between $1 - 6 \cdot 10^{-5}$ for O I, and $1 - 4 \cdot 10^{-4}$ for C II.

Compared to our LTE estimate in Sect. 4.1, the C II amount derived here is higher by factors of 13 – 160. Because of recombination reactions, it is rather unlikely to have a large amount of C II *hidden* in the cold gas, this can be considered as an upper limit.

5. Line cooling efficiency

The relative importance of the various line tracers among each other and with respect to the FIR continuum provides a detailed signature of the heating and cooling processes in the DR21 molecular cloud. Table 8 lists the cooling intensities at the five selected ISO positions E, C, W, OH, and FIR1.

The total line cooling intensity, I_{tot} , derived by summing the contributions of CO, C I, C II, and O I, peaks at position C, dropping to less than 15% of this peak value at FIR1.

The relative contributions to I_{tot} vary strongly at the 5 positions. [O I] is the strongest coolant at positions E and C, while CO is strongest at W, (OH), and FIR1, tying up more than 50 % of the total cooling. The relative contribution of CO to I_{tot} varies between 26 % at E and 76 % at (OH) while the relative contribution of [C II] is lower, varying between 7 % at C and 21 % at FIR1. The two [C I] fine structure lines contribute generally less than 2 %. The ratio of CO cooling relative to [C I] cooling, peaks at C at 114 and drops to less than 50 at FIR1.

We also compare I_{tot} with the FIR continuum to derive the gas cooling efficiency ϵ_{tot} as a measure of the fraction of FUV photon energy which goes into gas heating. ϵ_{tot}

Table 8. Total cooling intensities of CO, [C I], [C II], and [O I]. For CO we use the best fitting model (Sec. 4.2) and sum over all rotational lines up to $J=20$. For C I and O I, we sum the observed intensities of both fine structure lines respectively. The values in parentheses show the relative contribution to the sum I_{tot} . I_{IR} is the total observed infrared intensity (Table 5). The total cooling efficiency ϵ_{tot} is defined by $I_{\text{tot}}/I_{\text{IR}}$. For comparison, we also list the ϵ' defined by $(I_{[\text{C II}]} + I_{[\text{O I}]})/I_{\text{IR}}$.

Intensity in [10^{-4} erg s $^{-1}$ cm $^{-2}$ sr $^{-1}$]	DR21 E	DR21 C	DR21 W	DR21 (OH)	DR21 FIR1
$I_{\text{CO}} (= \Sigma \text{CO}_{1 \leq J_u \leq 20})$	24.1 (28.8 %)	51.3 (44.4 %)	18.7 (53.7 %)	20.0 (75.9 %)	13.4 (68.7 %)
$I_{[\text{C I}]} (= \Sigma [\text{C I}])$	0.22 (0.3 %)	0.45 (0.4 %)	0.23 (0.7 %)	0.25 (0.9 %)	0.27 (1.4 %)
$I_{[\text{O I}]} (= \Sigma [\text{O I}])$	51.1 (61.1 %)	55.6 (48.2 %)	11.2 (32.1 %)	3.78 (14.4 %)	2.23 (11.4 %)
$I_{[\text{C II}]}$	8.2 (9.8 %)	8.1 (7.0 %)	4.7 (13.5 %)	2.3 (8.8 %)	3.6 (18.5 %)
I_{tot}	83.62 (100 %)	115.45 (100 %)	34.83 (100 %)	26.33 (100 %)	< 19.5 (100 %)
I_{IR}	5500	33900	1810	12700	3680
ϵ_{tot} [%]	1.5	0.34	1.9	0.21	0.52
$I_{[\text{C II}]+[\text{O I}]} ([\text{C II}]+[\text{O I}]_{63 \mu\text{m}})$	56.4	57.5	14.8	5.24	5.38
ϵ' [%]	1.0	0.17	0.82	0.04	0.15

varies by one order of magnitude, between 0.2% at (OH) and 1.5 – 2% at the two outflow positions E and W.

Table 8 also lists the cooling efficiency ϵ' just taking into account the 158 μm [C II] and the 63 μm [O I] lines, ignoring CO, as is often done in the literature. ϵ' varies between 0.04% at (OH) and 1% at E.

Several studies in the literature have determined the gas heating efficiency ϵ' in Galactic star forming regions and in external galaxies. Vastel et al. (2001) find values of typically 0.2 – 0.4% in Galactic star forming regions, 1.1% at the Orion Bar, and less than 0.035% toward W49N. For W3 Main/IRS5, Kramer et al. (2004) report ϵ' of $\sim 0.1\%$ similar to the value found in DR21 C. Malhotra et al. (2001) find efficiencies between 0.05 and 0.3% in 60 unresolved normal galaxies. Kramer et al. (2005) find efficiencies between 0.21 and 0.36% in the centers and spiral arm positions of M83 and M51.

PDR models, e.g., by Kaufman et al. (1999), predict the heating efficiency ϵ' of a cloud illuminated from all sides between 0.01% to 1.5% and our values are within these theoretical limits.

A remarkable point is that only a few percent of the mass (up to 20% at DR21 E) is efficiently heated. Considering only the cold envelope component of our radiative transfer models, i.e., removing the warm interface zone, shows that the heated gas accounts for most of the CO emission (80% to 94%, summing over $J_{\text{up}} = 1...20$) in terms of radiated energy.

Since the total line power of all lines that we consider, I_{tot} , does not include the contributions from other cooling lines (e.g., of H₂, H₂O, OH, and PAHs) we may miss some of the cooling energy.

6. Summary

We presented the first comprehensive mapping of sub-mm lines in the whole DR21/DR21 (OH) region. The line emission of [C I] at 492 GHz and 809 GHz, CO 3–2, 4–3, 6–5, and 7–6, ¹³CO 3–2 and 6–5, and of CS 7–6 reveals contributions from different gas components: (i) A very wide winged emission in mid- J CO transitions having a line width of more than 15 km s $^{-1}$ and self-absorption, and (ii) narrow (4 to 6 km s $^{-1}$) emission in the other lines. Based on these lines, and supported by mm- and FIR-lines from FCRAO and ISO, we confined temperature, density, masses, and volume filling for five positions toward DR21, DR21 (OH) and DR21 FIR1 using a radiative transfer code by modeling the spectral line profiles.

1. We found that 2-component models consisting of a warm inner region and a cold envelope are needed to explain the observed strengths of the energetically low and high lying transitions of CO and to explain the self-absorption.

2. The bulk of molecular gas at DR21 & DR21 (OH) is cold (~ 30 K) and concentrated in the ridge. The clump volume filling is typically a few percent and the density within a clump is of the order 10⁶ cm $^{-3}$.

3. By fitting the dust emission SED, we constrain the H₂ column density to 2–5 10²² cm $^{-3}$ for DR21 E, W and toward FIR1. At DR21/DR21 (OH) it rises to above 10²³ cm $^{-3}$. The dust temperatures vary between 31–43 K and are close to the temperature of the cold gas.

4. Warm gas (80–150 K) was found at all five positions. The warm gas mass is typically a few percent of the total mass but accounts for most of the total CO emission (80% to 94%, mainly in the mid- and high- J transitions). DR21 (OH), as the coldest source, still shows a significant amount of high- J CO emission.

5. CS 7–6 emission can be reproduced if the local density in the envelope region is sufficiently high. Therefore, high densities are expected at DR21 and DR21 (OH) supporting again the low volume filling factors.

6. Supported by the model results, we derived the total intensity of the strongest emission lines ([O I], CO, [C II], and [C I]). Line emission provides up to 2% to the total cooling of the atomic and molecular gas and by dust grains. At the dense cores DR21 and DR21 (OH), the cooling efficiency is less than 0.5%. At DR21, O I and CO are equally strong, whereas at DR21 (OH) the CO lines dominate the cooling with more than 75%. C II contributes generally less than 20% and C I not more than 1.5% to the line cooling. The [O I] emission shows a large variation from < 15% at DR21 (OH) and FIR1 to 61% at E.

Acknowledgements. The KOSMA 3m submillimeter telescope at the Gornegrat-Süd is operated by the University of Cologne in collaboration with the Radio Astronomy Department of the Argelander-Institute for Astronomy (Bonn), and supported by special funding from the Land NRW. The observatory is administrated by the International Foundation Gornegrat & Jungfrauoch.

References

- Black, J. H. 2000, in *Astrochemistry: From Molecular Clouds to Planetary*, ed. Y. C. Minh & E. F. van Dishoeck, 81–+
- Bohlin, R. C., Savage, B. D., & Drake, J. F. 1978, *ApJ*, 224, 132
- Boreiko, R. T. & Betz, A. L. 1991, *ApJ*, 369, 382
- Bradford, C. M., Nikola, T., Stacey, G. J., et al. 2003, *ApJ*, 586, 891
- Bradford, C. M., Stacey, G. J., Nikola, T., et al. 2005, *ApJ*, 623, 866
- Brooks, K. J., Cox, P., Schneider, N., et al. 2003, *A&A*, 412, 751
- Calzetti, D., Armus, L., Bohlin, R. C., et al. 2000, *ApJ*, 533, 682
- Campbell, M. F., Nilas, D., Nawfel, R., et al. 1982, *ApJ*, 261, 550
- Cernicharo, J. 1985, ATM: A program to compute theoretical atmospheric opacity for frequencies < 1000 GHz, Tech. rep., IRAM
- Chandler, C. J., Gear, W. K., & Chini, R. 1993, *MNRAS*, 260, 337
- Clegg, P., Ade, P., Armand, C., & Baluteau, J.-P. 1996, *A&A*, 315, 38
- Crawford, M. K., Genzel, R., Townes, C. H., & Watson, D. M. 1985, *ApJ*, 291, 755
- Cyganowski, C. J., Reid, M. J., Fish, V. L., & Ho, P. T. P. 2003, *ApJ*, 596, 344
- Davis, C. J., Kumar, M. S. N., Sandell, G., et al. 2006, *MNRAS*
- Dickel, H., Lubenow, A., Goss, W., Forster, J., & Rots, A. 1983, *A&A*, 120, 74
- Dickel, J. R., Dickel, H. R., & Wilson, W. J. 1978, *ApJ*, 223, 840
- Downes, D. & Rinehart, R. 1966, *ApJ*, 144, 937
- Fixsen, D. J., Bennett, C. L., & Mather, J. C. 1999, *ApJ*, 526, 207
- Garden, R., Geballe, T. R., Gatley, I., & Nadeau, D. 1986, *MNRAS*, 220, 203
- Genzel, R. & Downes, D. 1977, *A&A*, 30, 145
- Giannini, T., Nisini, B., & Lorenzetti, D. 2001, *ApJ*, 555, 40
- Giannini, T., Nisini, B., Lorenzetti, D., et al. 2000, *A&A*, 358, 310
- Goldsmith, P. F., Bergin, E. A., & Lis, D. C. 1997, *ApJ*, 491, 615
- Graf, U. U., Heyminck, S., Michael, E. A., et al. 2002, in *Millimeter and Submillimeter Detectors for Astronomy*, ed. T. G. Phillips & J. Zmuidzinas (Proceedings of SPIE Vol. 4855)
- Habart, E., Abergel, A., Walmsley, C. M., Teyssier, D., & Pety, J. 2005, *A&A*, 437, 177
- Hatchell, J., Thompson, M. A., Millar, T. J., & MacDonald, G. H. 1998, *A&A*, 338, 713
- Jaffe, D. & Plume, R. 1995, in *Airborne Symposium on the Galactic Ecosystem*, ed. M. Haas, J. Davidson, & E. Erickson (ASP Conf.Ser., Vol.73)
- Jaffe, D. T., Genzel, R., Harris, A. I., et al. 1989, *ApJ*, 344, 265
- Jansen, D. J., van Dishoeck, E. F., Keene, J., Boreiko, R. T., & Betz, A. L. 1996, *A&A*, 309, 899
- Jørgensen, J. K., Schöier, F. L., & van Dishoeck, E. F. 2004, *A&A*, 416, 603
- Kaufman, M., Wolfire, M., Hollenbach, D., & Luhman, M. 1999, *ApJ*, 527, 795
- Keene, J., Lis, D., Phillips, T., & Schilke, P. 1997, in *Molecules in Astrophysics: Probes and Processes*, ed. E. F. van Dishoeck, IAU Symposium No. 178 (Dordrecht: Kluwer), 129
- Köster, B. 1998, Ph.D. thesis, Universität zu Köln
- Kramer, C., Jakob, H., Mookerjea, B., et al. 2004, *A&A*, 424, 887
- Kramer, C., Mookerjea, B., Bayet, E., et al. 2005, *A&A*, 441, 961
- Lane, A., Haas, M., Hollenbach, D., & Erickson, E. 1990, *ApJ*, 361, 132
- Langer, W. D. & Penzias, A. A. 1990, *ApJ*, 357, 477
- Leung, H. O. & Thaddeus, P. 1992, *ApJS*, 81, 267
- Malhotra, S., Kaufman, M., Hollenbach, D., et al. 2001, *ApJ*, 561, 766
- Mangum, J. G., Wootten, A., & Mundy, L. G. 1992, *ApJ*, 388, 467
- Marston, A. P., Reach, W. T., Noriega-Crespo, A., et al. 2004, *ApJS*, 154, 333
- Mauersberger, R., Wilson, T. L., Walmsley, C. M., Henkel, C., & Batrla, W. 1985, *A&A*, 146, 168
- Mookerjea, B., Kramer, C., Roellig, M., & Masur, M. 2006, *ArXiv Astrophysics e-prints*
- Motte, F., Bontemps, S., Schilke, P., et al. 2005, in *IAU Symposium*, 151–156
- Oka, T., Iwata, M., Maezawa, H., et al. 2004, *ApJ*, 602, 803

- Ossenkopf, V., Trojan, C., & Stutzki, J. 2001, *A&A*, 378, 608
- Palmer, P., Goss, W. M., & Whiteoak, J. B. 2004, *MNRAS*, 347, 1164
- Poglitsch, A., Herrmann, F., Genzel, R., et al. 1996, *ApJ*, 462, L43+
- Pollack, J. B., Hollenbach, D., Beckwith, S., et al. 1994, *ApJ*, 421, 615
- Preibisch, T., Ossenkopf, V., Yorke, H. W., & Henning, T. 1993, *A&A*, 279, 577
- Richardson, K. J., Sandell, G., Cunningham, C. T., & Davies, S. R. 1994, *A&A*, 286, 555
- Roelfsema, P. R., Goss, W. M., & Geballe, T. R. 1989, *A&A*, 222, 247
- Rubin, R. H., Simpson, J. P., Lord, S. D., et al. 1994, *ApJ*, 420, 772
- Schneider, N., Bontemps, S., Simon, R., et al. 2006, *A&A*, SBS2006
- Schneider, N., Simon, R., Kramer, C., et al. 2003, *A&A*, 406, 915
- Schroder, K., Staemmler, V., Smith, M. D., Flower, D. R., & Jaquet, R. 1991, *Journal of Physics B Atomic Molecular Physics*, 24, 2487
- Shirley, Y. L., Evans, N. J., Young, K. E., Knez, C., & Jaffe, D. T. 2003, *ApJS*, 149, 375
- Smith, H. A., Hora, J. L., Marengo, M., & Pipher, J. L. 2006
- Sternberg, A. & Dalgarno, A. 1989, *ApJ*, 338, 197
- Stutzki, J. 1985, Ph.D. Thesis
- Tielens, A. & Hollenbach, D. 1985, *ApJ*, 291, 722
- Vallée, J. P. & Fiege, J. D. 2006, *ApJ*, 636, 332
- van Zadelhoff, G.-J., Dullemond, C. P., van der Tak, F. F. S., et al. 2002, *A&A*, 395, 373
- Vastel, C., Spaans, M., Ceccarelli, C., Tielens, A. G. G. M., & Caux, E. 2001, *A&A*, 376, 1064
- Walker, C. K., Adams, F. C., & Lada, C. J. 1990, *ApJ*, 349, 515
- Wendker, H. 1984, *A&AS*, 58, 291
- Williams, J. A., Dickel, H. R., & Auer, L. H. 2004, *ApJS*, 153, 463
- Wilson, T. & Mauersberger, R. 1990, *A&A*, 239, 305
- Zmuidzinas, J., Betz, A. L., Boreiko, R. T., & Goldhaber, D. M. 1988, *ApJ*, 335, 774

Appendix A: Results at individual positions

Table 7 lists the derived physical parameters (shell- and model-averaged) for temperature, H₂ number density, volume filling factors, H₂ column density, and molecular gas mass. The resulting spectral profiles are compared with the observed lines in Fig. 5. The plots in Figure 8 show the fit of the integrated emission for CO, ¹³CO, and C¹⁸O up to $J = 20$ in comparison to the observed lines. In the following, we discuss details of the individual positions:

DR21 C: This position covers the DR21 H II-region. Its surrounding molecular cloud (within a radius of ~ 0.6 pc)

forms the southern part of the molecular ridge. Our best-fit model indicates that the envelope at a temperature of 40 K comprises roughly 94 % of the total mass, leaving 6 % for the warm (about 150 K) and dynamically active gas (v_{turb} between 4.5 and 29 km s⁻¹) subject to shocks and UV-heating. Jaffe et al. already found in 1989 indications of a warm component with a mass of $55 M_{\odot} (D/1.7 \text{ kpc})^2$, which is fully consistent with our result from Table 7. The by far more massive, colder gas component was only indirectly traced by the CO $J=7-6$ line reversal in their study. In our model, the properties of this gas component in the envelope shell is much better constrained by the C¹⁸O and ¹³CO $J=1-0$ transitions which are close to optically thin ($\tau \sim 0.1$ resp. 0.8). The turnover at $J_{\text{upper}} \sim 12$ (cf. Figure 8) results from the increasing critical density leading to sub-thermal excitation of the high- J transitions at the clump density of about $3 \cdot 10^6 \text{ cm}^{-3}$. Boreiko & Betz (1991) observed CO and ¹³CO 9–8 at comparable resolution onboard the Kuiper Airborne Observatory (KAO). We indicate their integrated fluxes in Fig. 8 for DR21 C. There is a good agreement with ¹³CO, but CO $J=9-8$ is weaker than the here presented $J=7-6$ line. This may indicate a stronger flattening of the CO rotation curve that could only be explained by a discontinuous density or temperature structure.

In support of a shock driven excitation, Lane et al. (1990) gave an upper limit to CO $J=22-21$ of $2 \cdot 10^{-4} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$ in a $44''.8$ beam. Considering beam dilution, this limit is still high compared to the intensity of CO 17–16, the highest rotational line considered here. Our prediction with the radiative transfer model of $2 \cdot 10^{-7} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}$ can explain why this detection attempt failed, albeit we cannot fully exclude the presence of > 200 K warm gas.

DR21 E: The east side of the DR21 H II-region is not located on the molecular ridge, but enriched with gas ejected from the ridge. The lack of cold gas gives rise to an average temperature of 62 K for a gas mass of $\sim 240 M_{\odot}$. From the Spitzer/IRAC image it appears that a fraction of gas in the east is undergoing a blister phase. Unlike the western outflow side, it is not stopped by a dense cloud and expands unhampered. We find that 20 % of the gas is heated to about 90 K and accelerated to velocities up to $v = 32 \text{ km s}^{-1}$. As the radiative transfer code is not designed for outflow geometries, we applied a turbulent velocity gradient from 32 to 5 km s⁻¹ to simulate the outflow. For the envelope gas, the asymmetry of the CO profiles indicates a motion away from the core at $\sim 0.7 \text{ km s}^{-1}$. Because of the steep drop of high- J CO emission, the local clump density was fitted below $n(\text{H}_2) = 10^5 \text{ cm}^{-3}$.

DR21 W: Molecular line emission and radiative transfer models of DR21 W are similar to position DR21 E: The gas is warmer than in the ridge (> 50 K) but not at densities above 10^5 cm^{-3} . The warm (~ 120 K) gas contributes

with a small fraction of about 3.5% to the total derived mass of $\sim 430 M_{\odot}$. A turbulent velocity gradient between 30 and 11 km s^{-1} is responsible for line wing emission. The CO lines show only weak self-absorption although the opacity is high at the line center. This is plausible because the cloud surface is illuminated by the cluster at DR21 C. Hence, we had to implement a positive temperature gradient from 44 K to 55 K across the cloud envelope.

DR21 (OH): The position of DR21 (OH) is dominated by cold gas from the ridge at a temperature below 30 K over a size of up to 0.7 pc. For the modeling, we simplified the scenario of several embedded cores as reported by, e.g., Mangum et al. (1992) (at possibly different systemic velocities), to a single clumpy core defined by the warm interface region with an outer radius of 0.2 pc. The significant mid- and high- J CO emission is a clear indication for warm and dense gas. We find only a few percent (2%, or $34 M_{\odot}$) of the gas is heated to ~ 100 K and at densities of 10^6 cm^{-3} or more. There is no evolved outflow associated with this position, but Garden et al. (1986) and Richardson et al. (1994) see indications of a very young outflow associated with high temperatures and shocks. In our model, half of the warm gas mass is at relative velocities $5 < v_{\text{turb}} < 32 \text{ km s}^{-1}$ creating the line wing emission. The observed blue-skewed CO line profiles are compatible with radial infall motion of the envelope gas of $1.0 - 1.5 \text{ km s}^{-1}$.

DR21 FIR1: The northern FIR-region contains at least three sub-sources. Besides FIR1, the $80''$ beam partly covers the second source FIR2, so we have to consider this as contamination. The modeled outer radius overlaps slightly with the neighbouring source, but is well within the beam. Because high- J CO is not detected at DR21 FIR1, we can constrain the temperature to ≤ 82 K. Given the CO 7–6 flux at this position, the lower limit of the temperature in the inner region is $T > 50$ K. The relatively weak self-absorption of the low- J lines is probably due to the presence of moderately warm gas (up to 40 K) in the outer surface layers, supporting the flat density profile used in the model.

Appendix B: Correction for Reference-Beam emission

Before the implementation of an On-The-Fly-mode (OTF), Dual-Beam-Switch (DBS) was the only efficient observing mode to use SMART at KOSMA. Therefore, observations in 2003 and 2004 of CO 7–6 and [C I] had to be performed in DBS-mode. In this mode, a chopping secondary mirror is used for fast periodic switching between an ON and an OFF position (Signal- and Reference phase). After some time, usually 20 sec, determined by the stability of the system, the telescope is moved so that the Reference phase is now the ON, and the new OFF position falls exactly on the opposite side. In the data reduction,




Fig. B.1. On-The-Fly map of the ^{13}CO 2–1 emission (SBS2006). An emission-free position is indicated by the triangle on the right and was used as OFF position for the OTF observations. The box indicates the map size of DBS observations in Fig. 2 and 3. The circle in the top left corner indicates the $120''$ beam size. The smaller circles labeled ON1&ON2, OFF1, and OFF2 show an exemplary DBS measurement (cf. Fig. B.2) with a field rotation angle $\alpha = 10^\circ$.

each OFF position is weighted equally. Slight differences in the optical path lengths usually lead to strong standing waves in the spectra when using a simple BS-mode. The period of these waves corresponds to the optical length to the subreflector. These standing waves are very efficiently suppressed in DBS-mode. However, mapping of extended sources is rather difficult if the chopper-throw (between $5' - 6'$) is small compared to the size of the source and the chopper movement is fixed to one direction (azimuth for KOSMA). Under those conditions, mapping needs careful planning to prevent self-chopping within the source. Observing time is thus not only constrained by source visibility but also by sky rotation. For an hour angle of 0^{h} , the secondary throw is only in right ascension, while for other angles there is a component in declination. The ^{13}CO 3–2 Position-Switched OTF-map in Fig. B.1 shows that the DR21 ridge is N-S orientated and field rotation angles α of less than $\pm 30^\circ$ are preferred in order to prevent chopping into the ridge of emission with both OFF beams.

To reduce the impact of artefacts remaining after the standard DBS reduction pipeline, all DBS data were corrected in the following way (see Fig. B.2 for an example): (i) Regardless whether the DBS-spectra (Fig. B.2 (a)) show self-chopping or not, the emission in the *difference* (Fig. B.2 (b)) of the two OFF positions is analysed. Any significant signal here is considered as pollution. This assumption holds as long as at least one of the two OFF positions is free of emission and no absorption against the continuum is present. In the example (Fig. B.2 (c) and (d)) only the OFF-position of Beam 1 (OFF1) is contaminated. (ii) If a signal in the difference is detected above the noise




Fig. B.2. This plot demonstrates the method to correct self-chopping in Dual-Beam-Switch observing mode. The uppermost spectrum (a) shows a standard DBS spectrum of C I 1–0 with a self-chopped absorption line (ON–0.5(OFF1+OFF2)), and (b) shows the difference signal (OFF2–OFF1). In the middle, (c) and (d), both beams of the DBS spectrum are displayed (Beam 1 or ON1–OFF1, and resp. for Beam 2). The result (e) (after removing standing waves) shows a faint emission line.

level, the algorithm tries to reconstruct the emission signal (Fig. B.2 (d)) by removing Beam 1 from the final spectrum (Fig. B.2 (e)), i.e. we use only ON2–OFF2 here. This scheme works well for situations where standing waves are negligible. In cases where standing-wave patterns are persistent, we transform the spectra to fourier-space, mask-out the standing wave frequencies, and transform the data back to user-space. Due to the shorter integration time, this procedure increases the noise level by a factor of $\sqrt{2}$. In case of DR21, this method successfully recovered the emission because one OFF position was always nearly free of emission. If, however, the source extends over more than twice the chop-throw in all directions this method does not work.

Appendix C: LTE Methods

A simple determination of the excitation temperature and the column density of C I can be done if we assume the emission is optically thin and the level population is given by thermal excitation. The column density of a material in an excited state u is given by:

$$N_u = \frac{8\pi k_B \nu^2}{hc^3 A_u} \int \frac{T_{\text{mb}} dv}{\text{K km s}^{-1}}. \quad (\text{C.1})$$

With Einstein A coefficients for atomic carbon $A_1 = 7.9 \times 10^{-8} \text{s}^{-1}$ and $A_2 = 2.7 \times 10^{-7} \text{s}^{-1}$ (Schroder et al. 1991) the column densities for each transition are

$$N_1 = 5.96 \cdot 10^{15} \int \frac{T_{\text{mb}}(\text{C I } ^3\text{P}_1 - ^3\text{P}_0) dv}{\text{K km s}^{-1}} [\text{cm}^{-2}] \quad (\text{C.2})$$

and

$$N_2 = 4.72 \cdot 10^{15} \int \frac{T_{\text{mb}}(\text{C I } ^3\text{P}_2 - ^3\text{P}_1) dv}{\text{K km s}^{-1}} [\text{cm}^{-2}]. \quad (\text{C.3})$$

To derive the excitation temperature, we assume LTE and substitute N_1 and N_2 using Boltzmann statistics:

$$\frac{N_u}{N_l} = \frac{g_u}{g_l} e^{-\frac{(E_u - E_l)/k_B}{T_{\text{ex}}}} \quad (\text{C.4})$$

$$T_{\text{ex}} = \frac{\frac{h\nu_2}{k_B}}{\ln\left(\frac{N_1 g_2}{N_2 g_1}\right)} = \frac{38.8 \text{ K}}{\ln\left(\frac{1.26 \cdot 5}{R_{\text{C I}} \cdot 3}\right)} = \frac{38.8 \text{ K}}{\ln(2.11/R_{\text{C I}})}, \quad (\text{C.5})$$

$R_{\text{C I}}$ is the ratio of integrated intensities (defined in (C.2) and (C.3)) and the statistical weight is $g_J = 2J + 1$. Although the temperature of the 3-level state of [C I] is very sensitive to small variations in $R_{\text{C I}}$, the total column density depends only weakly on the temperature:

$$\begin{aligned} N(\text{C I}) &= \sum_{J=0}^2 N_J = N_1 \frac{Z}{3e^{-\frac{23.6 \text{ K}}{T_{\text{ex}}}}} \\ &= 5.96 \cdot 10^{15} \frac{1 + 3e^{-\frac{23.6 \text{ K}}{T_{\text{ex}}}} + 5e^{-\frac{62.4 \text{ K}}{T_{\text{ex}}}}}{3e^{-\frac{23.6 \text{ K}}{T_{\text{ex}}}}} \times \\ &\int \frac{T_{\text{mb}}(\text{C I } ^3\text{P}_1 - ^3\text{P}_0) dv}{\text{K km s}^{-1}} [\text{cm}^{-2}] \end{aligned} \quad (\text{C.6})$$

where $Z = \sum_{J=0}^2 g_J e^{-\frac{E_J}{k_B T_{\text{ex}}}}$ is the partition function.

To estimate the excitation temperature from the two transitions $^{13}\text{CO } J=3-2$ and $J=1-0$ we used Eq. (C.4) which then translates into

$$T_{\text{ex,low}} = -\frac{(E_3 - E_1)/k_B}{\ln\left(\frac{N_3 g_1}{N_1 g_3}\right)} = \frac{26.4 \text{ K}}{\ln(9/R_{^{13}\text{CO,low}})}. \quad (\text{C.8})$$

The transition $J=6-5$ and $J=3-2$ gives accordingly

$$T_{\text{ex,high}} = \frac{79.4 \text{ K}}{\ln(4/R_{^{13}\text{CO,high}})}. \quad (\text{C.9})$$

$R_{^{13}\text{CO,low}}$ (resp. $R_{^{13}\text{CO,high}}$) is the integrated intensity ratio of 3–2/1–0 (resp. 6–5/3–2).

The ^{13}CO and C^{18}O column density can then be approximated by these equations

$$\begin{aligned} N(^{13}\text{CO}) &= 1.268 \cdot 10^{12} (T_{\text{ex}} + 0.88 \text{ K}) e^{111.1 \text{ K}/T_{\text{ex}}} \times \\ &\int \frac{T_{\text{mb}}(^{13}\text{CO}(6-5)) dv}{\text{K km s}^{-1}} [\text{cm}^{-2}], \end{aligned} \quad (\text{C.10})$$

$$\begin{aligned} N(^{13}\text{CO}) &= 4.71 \cdot 10^{13} (T_{\text{ex}} + 0.88 \text{ K}) e^{5.27 \text{ K}/T_{\text{ex}}} \times \\ &\int \frac{T_{\text{mb}}(^{13}\text{CO}(1-0)) dv}{\text{K km s}^{-1}} [\text{cm}^{-2}], \end{aligned} \quad (\text{C.11})$$

$$\begin{aligned} N(\text{C}^{18}\text{O}) &= 4.60 \cdot 10^{13} (T_{\text{ex}} + 0.88 \text{ K}) e^{5.27 \text{ K}/T_{\text{ex}}} \times \\ &\int \frac{T_{\text{mb}}(\text{C}^{18}\text{O}(1-0)) dv}{\text{K km s}^{-1}} [\text{cm}^{-2}]. \end{aligned} \quad (\text{C.12})$$

Using Eq. (C.4), an upper limit to the excitation temperature of O I can be derived from the [O I] 145/63 μm ratio (fluxes in $\text{erg s}^{-1} \text{cm}^{-2} \text{sr}^{-1}$) by:

$$T_{\text{ex}} = \frac{227.7 \text{ K}}{\ln(2.83/(12.22 \cdot R_{[\text{O I}]})}. \quad (\text{C.13})$$