

# *Flavour physics now and in the LHC era*


Gino Isidori

[ *Scuola Normale Superiore - Pisa & INFN - Frascati* ]

- ▶ Introduction
- ▶ What we learned so far
  - Model-independent fits
  - The MFV hypothesis
- ▶ What we could still hope to learn
  - The most interesting observables in the MSSM with MFV
  - Other observables
- ▶ Conclusions


► Introduction

I. In most realistic BSM scenarios, the new degrees of freedom of the model are heavier than B or K mesons (usually heavier also than W & Z bosons) and contribute to meson decays only via virtual effects → indirect sensitivity to NP



► Introduction

**I.** In most realistic BSM scenarios, the new degrees of freedom of the model are heavier than B or K mesons (usually heavier also than W & Z bosons) and contribute to meson decays only via virtual effects → indirect sensitivity to NP



**II.** If the new degrees of freedom respect the  $SU(2)_L \times U(1)$  gauge symmetry (very reasonable/general assumption) → NP effects at low energies decouple as  $1/\Lambda^2$  ( $\Lambda$  = energy scale of the new degrees of freedom)

trivial kinematical factors

$$A = A_0 \left[ c_{SM} \frac{1}{M_W^2} + c_{NP} \frac{1}{\Lambda^2} \right]$$

(adimensional) effective couplings

Detailed description: The equation shows the amplitude A as a product of a factor A\_0 and a bracketed sum of two terms. The first term is c\_SM / M\_W^2, and the second is c\_NP / Λ^2. Blue dashed arrows point from the text 'trivial kinematical factors' to A\_0, and from '(adimensional) effective couplings' to the bracketed sum. Another blue dashed arrow points from the text 'trivial kinematical factors' to the entire expression.

## ► Introduction

$$A = A_0 \left[ c_{\text{SM}} \frac{1}{M_{\text{W}}^2} + c_{\text{NP}} \frac{1}{\Lambda^2} \right]$$

$\Lambda$  = energy scale of the new particles


$c_{\text{SM(NP)}}$  = eff. couplings

- The sensitivity to the energy scale grows slowly with the statistics or the luminosity of the experiment (  $\sigma \sim 1/N^{1/4}$  )
- The interest of a given observable depends on the magnitude of  $c_{\text{SM}}$  vs.  $c_{\text{NP}}$  (*loop-induced observables usually more interesting because of small  $c_{\text{SM}}$ , but other type of suppressions, such as the helicity suppression, can make specific tree-level processes particularly interesting*)  
and on the theoretical error of  $c_{\text{SM}}$   
(*CKM + hadronic uncertainties → important role of auxiliary observables*)
- There is no way of disentangling the information on  $\Lambda$  and  $c_{\text{NP}}$ , but the combined information which can be extracted is fully complementary to the direct searches performed at high- $p_{\text{T}}$ : key role of (low-energy) flavour physics in determining the flavour symmetry structure of NP

► What we learned so far

The SM is very successful in describing quark-flavour mixing

This is quite clear by looking at the consistency of the exp. constraints appearing in the so called CKM fits, and is confirmed by the absence of significant deviations from the SM in clean rare decays such as  $B \rightarrow X_s \gamma$




$$V_{CKM} V_{CKM}^+ = I$$



triangular relations:


$$V_{i1} (V^+)_{1j} + V_{i2} (V^+)_{2j} + V_{i3} (V^+)_{3j} = 0$$



► What we learned so far

The SM is very successful in describing quark-flavour mixing

This is quite clear by looking at the consistency of the exp. constraints appearing in the so called CKM fits, and is confirmed by the absence of significant deviations from the SM in clean rare decays such as  $B \rightarrow X_s \gamma$




$$V_{CKM} V_{CKM}^+ = I$$



triangular relations:

$$V_{i1} (V^+)_{1j} + V_{i2} (V^+)_{2j} + V_{i3} (V^+)_{3j} = 0$$






*New physics effects in quark-flavour mixing can only appear as small corrections to the leading CKM mechanism*

## Model-independent fits of $\Delta F=2$ amplitudes

Present data allow us to determine the CKM unitarity triangle using only tree-level dominated amplitudes



General fit of NP in  $\Delta F=2$  amplitudes




## Model-independent fits of $\Delta F=2$ amplitudes

These general results are quite instructive if interpreted as bounds on the scale of new physics:

$$M(B_d - \bar{B}_d) \sim \frac{(V_{tb}^* V_{td})^2}{16 \pi^2 M_w^2} + \left( c_{\text{NP}} \frac{1}{\Lambda^2} \right)$$

contribution of the new heavy degrees of freedom

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{SM}} + \sum_{d \geq 5} \frac{c_n}{\Lambda^{d-4}} \mathcal{O}_n^d$$



## Model-independent fits of $\Delta F=2$ amplitudes

These general results are quite instructive if interpreted as bounds on the scale of new physics:

$$M(B_d - \bar{B}_d) \sim \frac{(V_{tb}^* V_{td})^2}{16 \pi^2 M_w^2} + \underbrace{c_{\text{NP}} \frac{1}{\Lambda^2}}_{\text{contribution of the new heavy degrees of freedom}}$$
  

$c_{\text{NP}}$	$\sim 1$	tree/strong + generic flavour	$\Lambda \gtrsim 2 \times 10^4 \text{ TeV [K]}$
	$\sim 1/(16 \pi^2)$	loop + generic flavour	$\Lambda \gtrsim 2 \times 10^3 \text{ TeV [K]}$
	$\sim (V_{ti}^* V_{tj})^2$	tree/strong + MFV	$\Lambda \gtrsim 5 \text{ TeV [K \& B]}$
	$\sim (V_{ti}^* V_{tj})^2 / (16 \pi^2)$	loop + MFV	$\Lambda \gtrsim 0.5 \text{ TeV [K \& B]}$

If you don't think this is an accident of  $\Delta F=2$ ...  $\Rightarrow$  MFV (Minimal Flavour Violation)

A rigorous definition of the Minimal Flavour Violation hypothesis:


The flavour structure of the SM is quite constrained:

- a large global symmetry in the gauge sector

$$U(3)^5 = \text{SU}(3)_Q \times \text{SU}(3)_U \times \text{SU}(3)_D \times \dots$$

- broken only by the Yukawa couplings

$$Y_D \sim \bar{3}_Q \times 3_D \quad Y_U \sim \bar{3}_Q \times 3_U \quad (Y_E \sim \bar{3}_L \times 3_E)$$



$$\mathcal{L}_{\text{SM}} = \mathcal{L}_{\text{gauge}} + \mathcal{L}_{\text{Higgs}}$$

$$\rightarrow \bar{Q}_L^i Y_U^{ij} U_R^j \phi + \bar{Q}_L^i Y_D^{ij} D_R^j \phi_c$$

This specific symmetry + symmetry-breaking

pattern is responsible for the GIM suppression of FCNCs,

the suppression of CPV, ... *the successful SM predictions in the quark flavour sector*

A rigorous definition of the Minimal Flavour Violation hypothesis:

The flavour structure of the SM is quite constrained:

- a large global symmetry in the gauge sector

$$U(3)^5 = \text{SU}(3)_Q \times \text{SU}(3)_U \times \text{SU}(3)_D \times \dots$$

- broken only by the Yukawa couplings



$$Y_D \sim \bar{3}_Q \times 3_D \quad Y_U \sim \bar{3}_Q \times 3_U \quad (Y_E \sim \bar{3}_L \times 3_E)$$



A natural mechanism to reproduce the SM successes in flavour physics -without fine tuning- is the MFV hypothesis:

*Yukawa couplings = unique sources of flavour symmetry breaking also beyond SM*

General principle which can be applied to any TeV-scale NP model



A rigorous definition of the Minimal Flavour Violation hypothesis:

*basic MFV:*

- global symmetry

$$U(3)^5 = \text{SU}(3)_Q \times \text{SU}(3)_U \times \text{SU}(3)_D \times \dots$$

- broken only by the Yukawa couplings

$$Y_D \sim \bar{3}_Q \times 3_D \quad Y_U \sim \bar{3}_Q \times 3_U \quad (Y_E \sim \bar{3}_L \times 3_E)$$


*Interesting extension/variation in case of more than one Higgs doublet:*

- With two Higgs doublets we can change the relative normalization of  $Y_U$  &  $Y_D$  (controlled by  $\tan\beta = \langle H_U \rangle / \langle H_D \rangle$ )

$$\mathcal{L}_{\text{q-Yukawa}} = \bar{Q}_L Y_D D_R H_D + \bar{Q}_L Y_U U_R H_U + \text{h.c.}$$

$$y_u = m_u / \langle H_U \rangle$$

$$y_d = m_d / \langle H_D \rangle = \tan\beta m_d / \langle H_U \rangle$$



A few important comments:

I) There is still room for non-MFV effects



A few important comments:

I) There is still room for non-MFV effects

According to a recent analysis by the UTfit collaboration [based on recent CDF & D0 results on  $B_s \rightarrow \psi\phi$ ], there is even a hit of a deviation from the SM in the CPV phase of  $B_s$  mixing that -if confirmed- would rule out both SM and MFV

$$C e^{2i\phi} = \frac{\langle M | H^{\text{SM+NP}} | \bar{M} \rangle}{\langle M | H^{\text{SM}} | \bar{M} \rangle}$$

Bona et al. arXiv:0803.0659






A few important comments:

I) There is still room for non-MFV effects


According to a recent analysis by the UTfit collaboration [based on recent CDF & D0 results on  $B_s \rightarrow \psi\phi$ ], there is even a hit of a deviation from the SM in the CPV phase of  $B_s$  mixing that -if confirmed- would rule out both SM and MFV

$$C e^{2i\phi} = \frac{\langle M | H^{\text{SM+NP}} | \bar{M} \rangle}{\langle M | H^{\text{SM}} | \bar{M} \rangle}$$

Caution needed given non-Gaussian errors (remember lesson from  $B_d \rightarrow \psi K$ )



Bona et al. arXiv:0803.0659



A few important comments:

- I) There is still room for non-MFV effects
- II) Even if we forget about  $B_s$  mixing, MFV is far from being “verified”

To prove MFV from data we would need to

- observe some deviation from the SM in FCNCs
- observe the CKM pattern predicted by MFV [within same type of FCNCs]

$$A_{\text{FCNC}} [b \rightarrow d(s)] \sim V_{td(s)} \left[ c_{\text{SM}}^{(0)} \frac{1}{M_W^2} + c_{\text{NP}}^{(0)} \frac{1}{\Lambda^2} \right]$$

$\Delta F = 2$  processes are in principle good candidates to prove MFV, but so far we are limited by theoretical (Lattice) uncertainties

Some  $\Delta F = 1$  rare decays could provide more useful infos to proof (or disproof) the MFV hypothesis from data (very interesting candidates:  $B_{d,s} \rightarrow l^+ l^-$ )

A few important comments:

- I) There is still room for non-MFV effects
- II) Even if we forget about  $B_s$  mixing, MFV is far from being “verified”
- III) Even within the “pessimistic” MFV hypothesis we can still expect sizable deviations from the SM in various B physics observables

Typical examples:

$B_{d,s} \rightarrow l^+ l^-$  up to order of magnitude enhancements if  $\tan\beta$  is large

$A_{FB}(B \rightarrow K^* l^+ l^-)$  up to O(1) deviations from the SM

A few important comments:

- I) There is still room for non-MFV effects
- II) Even if we forget about  $B_s$  mixing, MFV is far from being “verified”
- III) Even within the “pessimistic” MFV hypothesis we can still expect sizable deviations from the SM in various B physics observables



Typical examples:

$B_{d,s} \rightarrow l^+ l^-$	up to order of magnitude enhancements if $\tan\beta$ is large
$A_{\text{FB}}(B \rightarrow K^* l^+ l^-)$	up to O(1) deviations from the SM



- IV) The fact we have not observed yet a significant deviation from the SM in a few rare B decays (in particular  $B \rightarrow X_s \gamma$ ) puts significant constraints on the parameter space of NP models, even if they respect the MFV hypothesis

**E.g.:** The  $B \rightarrow X_s \gamma$  constraint in the CMSSM (after imposing Dark-matter conditions)




► What we could still hope to learn

Long list of potentially interesting observables, if we take into account that MFV has not been clearly established yet, and if we consider all the “auxiliary measurements” which could help in reducing SM uncertainties.


However, the list of interesting observables in “pessimistic” but very realistic scenarios, such as the MSSM with MFV, is rather limited.

The most interesting observables in the MSSM with MFV:




$$B^\pm \rightarrow l^\pm \nu$$

(  $B \rightarrow D \tau \nu$  )




$$B_{s,d} \rightarrow l^+ l^-$$



$$B \rightarrow X_s \gamma$$

The most interesting observables in the MSSM with MFV:



In models with 2 Higgs doublets (such as the MSSM) the  $H^\pm$  exchange appear at the tree-level in charged-current amplitudes. The effect is usually negligible (suppression of Yukawa couplings), except for helicity suppressed observables ( $B \rightarrow l \nu$ ) or  $\tau$  final states ( $B \rightarrow D \tau \nu$ )

Simple  $M_H$  &  $\tan\beta$  dependence

[mild dependence on other parameters]:

$$B(B \rightarrow l \nu) = B_{SM} \left( 1 - \frac{m_B^2 \tan^2\beta}{M_H^2 (1 + \epsilon_0 \tan\beta)} \right)^2$$

- O(10-30%) effect in  $B \rightarrow l \nu$
- ~ 3 times smaller in  $B \rightarrow D \tau \nu$
- ~ 100 times smaller in  $K \rightarrow l \nu$

The most interesting observables in the MSSM with MFV:

$$B(B \rightarrow \tau \nu) = (1.43 \pm 0.43) \times 10^{-4}$$

Babar+Belle '07

$$B(B \rightarrow \tau \nu)_{SM} = B_0 F_B^2 V_{ub}^2 \approx 1.2 \times 10^{-4}$$

sizable theoretical  
(parametric) error

$$B(K \rightarrow \mu \nu) = (63.66 \pm 0.17)\%$$


KLOE '06

+  $f_K/f_\pi$  @ 0.7%    MILC, UKQCD '07  
 +  $V_{us}$  @ 0.5%    KLOE, NA48, KTeV '06-'07

$B(B \rightarrow D \tau \nu)$  @ 10% potentially competitive


$F_{B \rightarrow D}$  known better than  $F_B$

$V_{cb}$  has a negligible error




Improving th. and exps.  
 on these channels can  
 lead to very valuable  
 infos on  $M_H$  &  $\tan\beta$  !

The most interesting observables in the MSSM with MFV:



There are no tree-level FCNC couplings of the neutral Higgses in MFV models; however, effective couplings can appear at the one loop level and they are potentially quite large in the MSSM



Crucial dependence on  $\mu$  and  $A_U$  [ +  $M_H$  &  $\tan\beta$ ]

$$A(B \rightarrow ll)_H \sim \frac{m_b m_l}{M_A^2} \frac{\mu A_U}{\tilde{M}_q^2} \tan^3\beta$$

Possible large enhancement over the SM, but the magnitude of the effect can vary a lot in different SUSY-breaking scenarios

## The most interesting observables in the MSSM with MFV:

$$B(B_s \rightarrow \mu\mu)_{\text{SM}} \approx 3.5 \times 10^{-9}$$

$$B(B_d \rightarrow \mu\mu)_{\text{SM}} \approx 1.3 \times 10^{-10}$$

$e$  channels suppressed by  $(m_e/m_\mu)^2$

$\tau$  channels enhanced by  $(m_\tau/m_\mu)^2$


Most interesting bound set by:

$$B(B_s \rightarrow \mu\mu) < 5.8 \times 10^{-8} \text{ (95\%CL)}$$



CDF+D0 '07

Significant constraint, but a good fraction of the parameter space is still allowed



**N.B.:** the  $B(B_d \rightarrow \mu\mu)/B(B_s \rightarrow \mu\mu)$  ratio is a key observable to proof or falsify MFV



The most interesting observables in the MSSM with MFV:






Most complicated observable with several, naturally competitive, contributions:





- positive
- decreasing with  $\tan\beta$
- sign  $\sim \text{sgn}(\mu, A)$
- increasing with  $\tan\beta$

The most interesting observables in the MSSM with MFV:



Most complicated observable with several, naturally competitive, contributions:



- positive
- decreasing with  $\tan\beta$

- sign  $\sim \text{sgn}(\mu, A)$
- increasing with  $\tan\beta$

One of the most significant constraint of the MSSM (even at small  $\tan\beta$ )

$$B(B \rightarrow X_s \gamma)^{\text{exp}} = (3.55 \pm 0.26) \times 10^{-4}$$

HFAG '06


$$B(B \rightarrow X_s \gamma)^{\text{SM}} = (3.15 \pm 0.23) \times 10^{-4}$$

Misiak et al. '06

E.g.: combined constraints in a general MFV-MSSM assuming heavy squarks:

Flavour physics  
 +  $(g-2)_\mu$   
 + dark matter  
 (A-funnel region)

$M_{sq} = 1.5 \text{ TeV}$     $M_{sl} = 0.5 \text{ TeV}$   
 $A_u = -1.0 \text{ TeV}$     $\mu = 0.5 \text{ TeV}$




## Other observables (a very incomplete list):

### I. Improved CKM fits


Improving the determination of the CKM matrix from tree-level processes offer a valuable tool to improve constraints on NP (including MFV models).

Key measurements:

▶  $\gamma$  from various  $B_{(s)} \rightarrow D(\bar{D})$  modes



good prospects of improvements from **LHCb**




All relevant hadronic parameters extracted from data with no theoretical assumptions

Other observables (a very incomplete list):


II. Time-dependent CPV

Is there still some hope to observe *significant* NP effects in time-dep. CPV asymmetries in  $b \rightarrow s$  hadronic-penguin modes ?

E.g.:



$\sin(2\beta^{\text{eff}}) \equiv \sin(2\phi_1^{\text{eff}})$  **HFAG**  
LP 2007  
PRELIMINARY



Other observables (a very incomplete list):

II. Time-dependent CPV

Is there still some hope to observe *significant* NP effects in time-dep. CPV asymmetries in  $b \rightarrow s$  hadronic-penguin modes ?


*Personally I'm quite skeptical...*

- Best observables [high stat. + full Dalitz Plot analysis] show no significant effect
- We are already close to the level of irreducible th. errors [remember the  $\epsilon'/\epsilon$  lesson...]

$$\sin(2\beta^{\text{eff}}) \equiv \sin(2\phi_1^{\text{eff}})$$



LP 2007  
PRELIMINARY




Other observables (a very incomplete list):

**II. Time-dependent CPV**

The time-dependent CPV asymmetry which is definitely worth to improve in the LHC era is the **phase of  $B_s$  mixing** from  $B_s \rightarrow \psi\phi \Rightarrow$  Tevatron/LHCb

New theoretically clean observable which could allow to falsify MFV [or to constraint viable non-MFV models] in the  $\Delta F = 2$  sector



## ► Conclusions

We learned a lot about flavour physics in the recent past...

*..but what is still to be discovered is more !*

TeV-scale NP models must have a rather sophisticated flavour structure (not to be excluded by present data) but we have not clearly identified this structure yet



Important to continue high-statistics / high-precision B physics in the LHC era

In realistic models there is only a limited set of particularly interesting observables [*theoretically-clean leptonic/semileptonic final states*]

but these observables play a key role in determining the flavour symmetry structure of NP